

THE MIGRANT



A QUARTERLY JOURNAL
DEVOTED TO TENNESSEE BIRDS

PUBLISHED BY
THE TENNESSEE
ORNITHOLOGICAL
SOCIETY

DECEMBER 2018
VOL. 89, No. 4

THE MIGRANT
A QUARTERLY JOURNAL OF ORNITHOLOGY
PUBLISHED BY

THE TENNESSEE ORNITHOLOGICAL SOCIETY
The TOS is a non-profit, educational, scientific and conservation organization.

EDITORIAL STAFF

Editor: Bob Ford, 808 Hatchie, Brownsville, TN 38012 <editorthemigrant@gmail.com>

Associate Editors:

Susan McWhirter, 274 Beech Grove Road, McMinnville, TN 37110 <snmcwhirter@gmail.com>
and
Martha Waldron, 1014 Murray Hill Lane, Memphis, TN 38120 <martha.waldron@gmail.com>

State Count Compiler: Ron Hoff, 282 Hackworth Lane, Clinton, TN 37716 <aves7000@bellsouth.net>

Season Editor: Richard L. Knight, 804 North Hills Dr., Johnson City, TN 37604 <rknight8@earthlink.net>

Regional Season Editors:

Western Coastal Plain: Dick Preston, 261 Sassafras Circle, Munford, TN 38058
<dickpreston@rittermail.com>

Highland Rim and Basin: Stephen C. Zipperer, 3105 Chapel Hills Drive, Murfreesboro, TN 37129
<stczipperer@aol.com>

Eastern Ridge and Valley: Richard L. Knight, 804 N. Hills Rd., Johnson City, TN 37604
<rknight8@earthlink.net>

Eastern Mountain: Richard P. Lewis, 407 V.I. Ranch Rd., Bristol, TN 37620 <mountainbirds@email.com>

OFFICERS FOR 2018

President: Dr. Steve Routledge, 1515 N. Willow Bend Court, Clarksville, TN 37043 <eyerout@bellsouth.net>

Vice Presidents:

East Tenn. - Rack Cross, 300 River Edge Drive, Kingsport, TN 37660 <rackcross@gmail.com>

Middle Tenn. - Danny Shelton, 705 Countrywood Drive, Franklin, TN 37064 <dashelt@comcast.net>

West Tenn. - Martha Waldron, 1040 Murray Hill Lane, Memphis, TN <martha.waldron@gmail.com>

Treasurer: Pam Lasley, 5886 Wilshire Drive, Nashville, TN 37215 <plasley@comcast.net>

Secretary: Cyndi Routledge, 1515 N. Willow Bend Court, Clarksville, TN 37043 <routledges@bellsouth.net>

Directors-at-Large:

East Tenn. - Susan Hubley, 128 John Sevier Circle, Rogersville, TN. 37857 <shubley@msn.com>

Middle Tenn. - Daniel Moss, 395 Janet Drive, Pleasant View, TN 37146 <dmos@earthlink.net>

West Tenn. - Allan Trently, 580 East Lafayette Street, Jackson, TN 38301 <Allan.Trently@tn.gov>

Curator and Webmaster: Charles P. Nicholson, P.O. Box 402, Norris, TN 37828 <cpnichol@bellsouth.net>

Assistant Curator: Dr. Stefan Woltmann, Austin Peay State University, 601 College St., Clarksville, TN 30740
<woltmanns@apsu.edu>

The Tennessee Warbler (TOS Newsletter) Editor: Theresa Graham, P.O. Box 366, Oakland, TN 38060
<t2graham@bellsouth.net>

The TOS website can be found at: www.tnbirds.org

Send subscriptions & address changes to:

Tennessee Ornithological Society, 5886 Wilshire Drive, Nashville, TN 37215

Printed by: Russell Printing Options, 1800 Grand Ave., Knoxville, TN 37916

Copyright © 2014 by the Tennessee Ornithological Society — ISSN 0026-3575575

THE MIGRANT

Published by the Tennessee Ornithological Society
to Record and Encourage the Study of Birds in Tennessee
Issued in March, June, September and December

VOL. 89

DECEMBER 2018

NO. 4

The Migrant 89(4): 117-119, 2018

STATUS OF THE SWALLOW-TAILED KITE IN TENNESSEE

Richard L. Knight
Johnson City, Tennessee

Swallow-tailed Kites (*Elanoides forficatus*) breed in the southeastern United States as well as in southern Mexico and southward into South America. The U.S. population currently breeds on the lower coastal plain from South Carolina to eastern Texas and migrates to wintering grounds in South America (Alderfer 2006). Historically this species nested further north, including along the Mississippi River drainage as far north as Minnesota and Ohio and probably nested in West Tennessee, but vanished from these northern regions by the early 1900s (Robertson 1988). Despite a small, but apparently stable, breeding population in the U.S., this species has shown a recent trend of increasing extralimital sightings in states north of its breeding range (Alderfer 2006). Herein I summarize the increased frequency of observations in Tennessee of this spectacular raptor.

Robinson (1990) cited just 11 records of Swallow-tailed Kites in Tennessee up to 1986. A literature search of *The Migrant* and *North American Birds* produced a total of at least 56 records in the state through 2017, a rather dramatic increase during the last three decades. Of this total, five records occurred from 1810 to 1933, with two records in the 1960s, one record in the 1970s, three records in the 1980s, and three records in the early 1990s. The period of greatest increase has been since 1999 with at least 42 records through 2017, a span of 19 years. Indeed, Swallow-tailed Kites have occurred in Tennessee annually since 1999 except for the years 2003 and 2004. The number of sightings per year of occurrence since 1999 has ranged from one to seven, with the greatest number in 2016.

Slightly over half of all sightings have come from East Tennessee (57%), with the remainder split evenly between West and Middle Tennessee (Table 1). Sightings have occurred in 30 counties. The Sequatchie Valley portion of Bledsoe and Sequatchie counties has hosted these kites seven times, more than any other location in the state. Most reports (80%) in the state have been of a single bird. The highest counts in the state were ten kites in Bledsoe County in August 2012 (Knight 2012), eight to ten in Obion County in August

1886 (Yeatman 1965), and up to seven in August 2016 again in Bledsoe County (Knight 2017).

Sightings of Swallow-tailed Kites in Tennessee have occurred in two distinct seasons, spring and late summer to early autumn (combined here as “fall”). Spring reports have ranged from 10 April to 16 June, while fall reports have ranged from 6 July to 25 September. The majority of sightings have occurred in the fall period (80%, Table 1), with most in the first three weeks of August. Spring sightings generally are considered to be over-shooting migrants, while reports in fall represent post-breeding dispersal. Of Tennessee sightings since 1968, nearly three quarters were observed on just a single day. All nine modern spring sightings have been of a single kite seen on just one day, perhaps not surprising for individuals apparently still migrating. The longest stay by Swallow-tailed Kites at one location in the fall is for 25 days in August 2012 in Bledsoe County (Knight 2012), although there likely was some turnover of individuals during this span.

Most sightings in Tennessee have come from river valleys with a mix of woodlands and open fields where the kites can hunt for large aerial insects. However, these kites have been seen from mountain tops, too, such as Camp Creek Bald in Greene County (Lewis 2007). Therefore, early season hawk-watching from traditional ridge-top sites could produce additional sightings.

The growth in numbers of birders in recent decades and the greater awareness of kite distribution by the birding community may have contributed to the increase in the number of Swallow-tailed Kite sightings in Tennessee. Also, the distribution of birders within the state may have skewed the regional distribution of kite sightings somewhat. Nevertheless, it is evident from the data presented here that a genuine increase in Swallow-tailed Kites has occurred in the state.

Table 1. Regional and Seasonal Distribution of Swallow-tailed Kites in Tennessee.

	<u>Number of Records</u>	<u>Percent of Total</u>
West Tennessee	12	21
Middle Tennessee	12	21
East Tennessee	32	57

Spring	10	18
Fall	45	80
Unknown	1	2

LITERATURE CITED

- Alderfer, J. 2006. Complete Birds of North America. National Geographic Society, Washington, D.C.
- Knight, R.L. 2012. The season: Cumberland plateau / ridge and valley region. *The Migrant* 83:151-156.
- Knight, R.L. 2017. The season: Cumberland plateau / ridge and valley region. *The Migrant* 88:49-53.
- Lewis, R.P. 2007. The season: eastern mountain region. *The Migrant* 78:40-43.
- Robertson, W.B., Jr. 1988. American Swallow-tailed Kite. Pages 109-131 in *Handbook of North American Birds, Volume 4*. (R.S. Palmer, Editor). Yale University Press, New Haven, Connecticut.
- Robinson, J.C. 1990. *An Annotated Checklist of the Birds of Tennessee*. University of Tennessee Press, Knoxville.
- Yeatman, H.C. 1965. Swallow-tailed Kite in Franklin County, Tennessee. *The Migrant* 36:58-59.

LATE NESTING KILLDEER IN HAMILTON COUNTY

David A. Aborn
Chattanooga, Tennessee

Killdeer (*Charadrius vociferus*) are common breeding birds throughout Tennessee (Somershoe and Sloan 2015). Nesting in Tennessee typically commences in February, with most eggs laid in March and April, and breeding is usually completed by the end of the summer (Nicholson 1997). On 19 November 2017, I received a phone text of a Killdeer nesting at a church playground in Chattanooga. The text included a picture of what looked like four Killdeer eggs in a shallow scrape in the gravel, as well as video of a bird doing the broken-wing distraction display. On 20 November, I went to the church to verify the nest, which was indeed a Killdeer nest (Figs. 1). I cordoned off an area around the nest with red flagging tape so that the nest would not get stepped on by anyone using the playground. I monitored the nest one to two times per week thereafter. When I checked on 3 December one egg was missing, but the other three were present and intact, and the adult was doing the distraction display. When I checked the nest on 12 December, there were two eggs in the nest and about 50 m away the adult was brooding a 1-2 day old chick. The eggs were cold and were presumably either abandoned by the adult soon after the chick hatched or were infertile. Adults and chicks typically remain at the nest until all chicks have hatched and their plumage has dried (Jackson and Jackson 2000). I returned on 15 December and I saw neither the adult nor the chick. Assuming the clutch was completed on 19 November and the one egg hatched on 11 December that gives a 23-day incubation period, which is in the typical 22-28 day incubation range for this species (Jackson and Jackson 2000).

The average high temperature during this period was 16^o C (61^o F) and the average low temperature was 2^o C (37^o F), however there was a significant cooling in the days prior to hatching. From 6 – 11 December, the average high was 3^o C (47^o F) and the average low was 0^o C (32^o F). Bergstrom (1989) found a strong correlation between ambient temperature and egg temperature in a Killdeer nest in Virginia, and since egg temperature affects embryonic development, it is likely that the parents compensated for the cooler temperatures by spending more time incubating the eggs.

There is only one other record of a Killdeer nesting that late in Tennessee; in 2008 a nest with three eggs was found in Shelby County (Knight 2009). The nest was not monitored, so it is unknown whether any of the eggs hatched. Other November nests have been found in Mississippi, Arkansas, and South Carolina (Jackson and Jackson 2000). This account constitutes the latest successful nesting of Killdeer in Tennessee.

ACKNOWLEDGEMENTS

I thank Sherry Ferguson of St. Stephen Catholic Church for alerting Happinest Wildlife Rescue to the presence of the nest, and Sherry Teas from Happinest who passed the information along to me.

LITERATURE CITED

- Bergstrom, P.W. 1989. Incubation temperatures of Wilson's Plovers and Killdeers. *The Condor* 91:634-641.
- Jackson, B.J. and J.A. Jackson. 2000. Killdeer (*Charadrius vociferus*), version 2.0. *The Birds of North America* (P. G. Rodewald, editor). Cornell Lab of Ornithology, Ithaca, New York, USA. <https://doi.org/10.2173/bna.517>. Accessed 12 December 2017.
- Knight, R.L. 2009. The fall season. *The Migrant* 80:22.
- Nicholson, C.P. 1997. *Atlas of the Breeding Birds of Tennessee*. University of Tennessee Press, Knoxville, Tennessee.
- Somershoe, S.G., and C.A. Sloan. *Birds of Tennessee: a new annotated checklist*. CreateSpace Independent Publishing Platform



Figure 1. Killdeer nest with 4 eggs in Chattanooga. Photo taken by David Aborn, 20 November 2017.

ROUND TABLE NOTES

FIRST SWAINSON'S HAWK RECORD FOR MIDDLE TENNESSEE - On the afternoon of 9 March 2018 I participated in a prescribed burning operation on Fort Campbell Military Installation (Montgomery County, TN; 36° 36' 28.740, -87° 33' 43.019). While riding an ATV I saw a buteo flying low over the freshly burned field. It came towards me and landed in a nearby sweetgum tree (*Liquidambar styraciflua*). I expected it to be a Red-tailed Hawk (*Buteo jamaicensis*) as they typically frequent the fields when we are burning. Because of the reduced visibility from my goggles and helmet, I drove the ATV to within 20 m of the tree to get a better look. The hawk did not flush and was very tolerant of my presence; the bird remained perched when another burn crew member drove up next to me on an ATV.

I saw that the hawk did not have the distinctive belly band of a Red-tailed Hawk, but instead had a dark rufous patch on its breast. When I realized this was an interesting buteo and I did not have binoculars, I immediately took out my iPhone 6s Plus phone and snapped a few photos in the zoom mode. The long, pointed wings and slender body of the perched hawk were noticeable. When it did eventually fly from the tree, I noticed the dark remiges contrasting with the light wing linings. I did not see the bird again after it flew from the tree. Looking at the photographs (Figure 1) later provided convincing identification of a light morph Swainson's Hawk (*Buteo swainsoni*).

This sighting is the first Swainson's Hawk documented in middle Tennessee and the earliest spring record for the state, according to Somershoe and Sloan (2015). Swainson's Hawk is listed as a very rare transient on the Official List of Birds of Tennessee (Tennessee Birds Record Committee). The closest known record to this observation occurred in 1989 in Calloway County, Kentucky (Brainard Palmer-Ball, pers. comm.).

Swainson's Hawks, like many raptor species, are attracted to fire and prescribed burns. Migrating Swainson's Hawks were the most abundant raptor observed using prescribed fires in the Great Plains (Hovick et al. 2017). Hovick et al. (2017) concluded Swainson's Hawks, because they are opportunistic feeders that covered large distances per day in migration, developed a behavioral attraction to smoke plumes allowing them to take advantage of foraging opportunities. Swainson's Hawks have also been observed hunting over freshly burned fields in California (Estep 1989) and Texas (Baker 1940).

The field consisted primarily of native grasses, forbs, blackberry (*Rubus fruticosus*), and some hardwood saplings prior to burning. There were a few scattered mature hardwood and loblolly trees (*Pinus taeda*) in the field.

LITERATURE CITED

- Baker, R. H. 1940. Effects of burning and grazing on rodent populations. *Journal of Mammalogy* 21:223.
- Estep, J. A. 1989. Biology, movements, and habitat relationships of the Swainson's Hawk in the Central Valley of California, 1986-87. California Department of Fish and Game, Nongame Bird and Mammal Sec. Rep., 52pp.
- Hovick, T.J., D.A. McGranahan, R.D. Elmore, J.R. Weir, S.D. Fuhlendorf. Pyric-carnivory:

Raptor use of prescribed fires. *Ecol Evol.* 2017;7:9144–9150. <https://doi.org/10.1002/ece3.3401>.

Somershoe, S. G., and C. A. Sloan. 2015. *Birds of Tennessee: A New Annotated Checklist*. CreateSpace Independent Publishing.



Figure 1. A light morph Swainson's Hawk photographed with an iPhone 6s Plus camera during a prescribed burning operation on Fort Campbell Military Installation (Montgomery County) 9 March 2018.

Daniel Moss, Fort Campbell, Tennessee

MINUTES OF THE FALL TOS BOARD OF DIRECTORS MEETING

6 October 2018
Manchester, Tennessee

The annual Fall Meeting was held 5-7 October 2018 at The Ambassador Inn in Manchester, Tennessee and hosted by the Highland Rim Chapter. Field trips were offered to Tims Ford Lake and Woods Reservoir.

President Steve Routledge called the meeting to order at 2:40 pm. Secretary Cyndi Routledge determined there was a quorum. The minutes of the 2018 Annual Spring Board of Directors and Spring Meeting were approved as read.

REPORTS OF OFFICERS AND DIRECTORS -AT-LARGE

TREASURER: Mac McWhirter reported the revenues and expenses are running as expected with no unexpected surprises in 2018. At this time the Vanguard investment portfolio is valued at approximately \$372,000. This is an increase of \$19,000 from the 31 December 2017 value of \$353,232. We have built a sizable investment account over the years in anticipation of future partnerships for land acquisition by the Tennessee Division of Environment and Conservation, NAWCA and TWRA.

COMMITTEE REPORTS

TENNESSEE BIRD RECORDS COMMITTEE: Secretary Ruben Stoll sent a report of the actions of the committee as of 30 September 2018. This report in its entirety may be found in *The Migrant*, Volume 89, No. 3, (September 2018). Current members of the committee are Phillip Casteel, Ron Hoff, Daniel Jacobson, David Kirschke (alternate) Chris Sloan, Ruben Stoll (secretary) and Terry Witt.

CONSERVATION POLICY COMMITTEE: Co-chairs of the committee are Dick Preston and Melinda Welton; members are Cyndi Routledge, Kellie Clelland, Ashley Heeney, Laura Cook, and Dev Joslin. Dick Preston reported the following action of the Committee.

- Drafted a letter to the developers of the Parkside Development in Memphis requesting incorporating bird-friendly design into the windows of the boutique hotel to reduce bird kills. This development is immediately adjacent to Shelby Farms, a major bird oasis in Shelby County. Signed by President Steve Routledge.
- Recommended TOS join with other organizations in sign-on letters to oppose harmful riders to the Farm Bill and addressing threats to the Endangered Species Act. Both approved and signed by President Steve Routledge.
- Drafted a letter for President Steve Routledge to the Secretary of the Interior. The letter

urges support for the U.S. Geographical Survey (USGS) Bird Banding Lab (BBL). The BBL faces a severe shortage of staff and funding.

- Urged the support of the House version of the Recovering America's Wildlife Act, which provides for continuing funding, while the Senate version does not. This act would provide Tennessee with more than \$20 million per year, primarily for non-game species.

PUBLICATIONS AND WEBSITE: Committee members are Chuck Nicholson, Ashley Heeney, Melinda Welton, and Rick Blanton. Ashley Heeney sent the following report that was presented by President Steve Routledge: The committee proposes building a new and modern/user-friendly website for TOS to launch at the beginning of 2019. The new site will feature current offerings, historical resources, plus highlight current chapter news and integrate with social media. The website will remain tnbirds.org. Mac McWhirter made a motion that the Executive Committee continue research, contract negotiations, and approve costs for redesign of website. Motion carried.

TNBird: President Steve Routledge reported that Rob Biller will be replace Andy Jones as a moderator of the site.

OLD BUSINESS

Discover Birds Activity Book coordinator Cyndi Routledge reported that Tennessee Wildlife Resources Agency (TWRA) has offered up to \$3000 annually for reprinting of the books. She requested that the Society match the funds set forth by TWRA so printing and distribution of the book could continue. Ron Hoff moved TOS spend up to \$3000 annually to match the funds provided by TWRA in like amount until rescinded for printing of the *Discover Birds Activity Book*. Motion carried.

NEW BUSINESS

NOMINATING COMMITTEE: Members are Susan Hubley, Anita Gaddy, and Chair Susan McWhirter. Susan McWhirter placed in nomination Pamela Lasley of Nashville for Treasurer effective 1 January 2019, and asked that she be authorized to transact all financial matters for the Society, including, but not limited, to being the signatory on our current banking account with Regions Bank. Chris Welsh made a motion to accept the nomination and the motion passed.

The meeting adjourned at 4:40 p.m.

Respectfully submitted,

Cyndi Routledge, Secretary

2018 TENNESSEE FALL BIRD COUNTS

Ron Hoff
Loudon, Tennessee

There were eight Fall Bird Counts conducted this year. Participants logged just over 412 party hours in the field to find 189 species and 32,352 individuals (lowest individuals total in 11 years). The weather was mild during the count season with no major rain or wind events having an effect on the counts. Temperatures ranged from a brisk 50° F in Perry County to a very warm 92° F in Shelby County.

White Ibis, Purple Gallinule, Black-legged Kittiwake, and Purple Finch were all new species bringing the species total for the Fall Bird Count compilation to 255. Count highlights also included Gadwall, Least Bittern, Osprey (highest total in 11 years), “Traill’s” Flycatcher spp, 33 species of warblers (including a Brewster’s hybrid), and 843 Rose-breasted Grosbeaks. Loggerhead Shrike was missed for the first time on a fall count in 11 years.

Counts are placed in the table geographically from West to East. Taxonomy follows the latest revision of the AOS Checklist of North American Birds.

COUNTY SUMMERIES (F/Y = feeder/yard watcher)

Anderson County -23 September; 0659-1424. Weather: cloudy to partly cloudy, some light rain; 68-77° F. Red-headed Woodpecker was a highlight. Observers: Eric Bodker, Doug Bruce, Chuck Estes, Melinda Fawver (compiler), Bill Keeler, Toby Koosman, Stephen J. Stedman, and Dean Turley.

Blount County – 15 September; 0600-2030. Weather: cloudy to partly cloudy; wind 0-8 mph; 65-87° F. Yellow-crowned Night-Heron, Sora, and Olive-sided Flycatcher were notable. Observers: Jean J. Alexander (compiler), Tiffany Beachy, Suzanne Bielenberg, Warren Bielenberg, Doug Bruce, Chuck Estes, Lola Estes, Melinda Fawver, Marian D. Fitzgerald, Tom E. Fitzgerald, Kim J. Henry, Stephen P. Henry, Thomas D. Howe, Susan Hoyle, James R. Human, David M. Johnson, Evan Kidd, Julie Labhart, Karen J. Petrey, Mary Tankersley, Harumi Tsuruoka, and Randy Winstead.

Elizabethton – 29 September; 0415-1900. Weather: partly cloudy; wind light and variable; 57-82° F. Two highlights were Purple Gallinule and Black-legged Kittiwake. Observers: Fred Alsop, Jim Anderson, Betty Bailey, Gary Bailey, Jerry Bevins, Rob Biller, Aleesha Broyles, Debbie Campbell, J. G. Campbell, Ron Carrico, Diane Draper, Glen Eller, Harry Lee Farthing, Bambi Fincher, Dave Gardner, Bill Grigsby, Paul Haynes, Jacki Hinshaw, Don Holt, Mike Hubley, Susan Hubley, Connie Irick, David Irick, David Kirschke, Rick Knight (compiler), Roy Knispel, Richard Lewis, Vern Maddux, Larry McDaniel, Joe McGuinness,

Cathy McNeil, Tom McNeil, Eric Middlemass, Charles Moore, Harry Norman, Brookie Potter, Jean Potter, Sherrie Quillen, Pete Range, Judi Sawyer, Helen Sirett, Chris Soto, Michele Sparks, Amber Stanley, Bryan Stevens, Kim Stroud, Gary Wallace, Charles Warden, Mary Anna Wheat, and Wendy Williams.

Knoxville – 29 September; 0630-1915. Weather: clear to partly cloudy; wind NW 0-5 mph; 60-85° F. Highlights were Sora, Marsh Wren, and Wilson's Warbler. Observers: Mark Armstrong, Sam Bee, Doug Bruce, Billie Cantwell, Gail Clendenen, Steve Clendenen, Howard Cox, Ellie Doughty, Melinda Fawver, Paul Hartigan, Susan Hoyle, James Human, Kristine Johnson, Jane Kading, Susie Kaplar, Jerry Ledbetter, Collin Leonard, Morton Massey, Stephany McNew, Stephanie Mueller, Talissa Ralph, Doug Raybuck, Kelly Roy, Martha Rudolph, Michael Ryon, Kurt Sickafus, Bailey Smith, Colin Sumrall, Douglas Sumrall, Michael Tenbus, Harumi Tsuruoka, Jimmy Tucker, Merikay Waldvogel, Keith Watson, and Eddy Whitson. Non-participating compiler was K. Dean Edwards.

Montgomery County – 15 September; no times given. Weather: fair; wind N 2-8 mph; 68-86° F. Highlights were Little Blue Heron, American Woodcock, and Marsh Wren. Observers: Elaine Faust, Daniel Moss (compiler), Rick Shipkowski, and Sue Shipkowski.

Nashville – 28-30 September; 0500-1800. Weather: partly cloudy to clear; 58-78° F. The count takes place over a 3 day period. Notable finds included Least Bittern, Black-billed Cuckoo, Olive-sided Flycatcher, and Brewster's Warbler. Observers: Rachael Anderson, Sharon Arnold, Susan Bradfield, Trae Bradfield, Jean Buchanan, Ed Byrne, Phillip Casteel, Richard Connors, Tammy Devine, Jerry Drewry, Graham Gerdeman, Steve Goodbred, Chris Guerin, Joe Hall, Barbara Harris (compiler), Charles Newman, Giovanni Ortiz, Amy Potter, Sarah Scott, Jan K. Shaw, Shelia Shay, Chris Sloan, Michael Smith, Joe Stone, Edwin Warner, Percy Warner, LinnAnn Welch, Melinda Welton, and Mary Zimmerman.

Perry County - 29 September; 0400-1920. Weather: clear; 50-80° F. Highlights included Herring Gull, Black-billed Cuckoo (new for the count), and Sedge Wren. The total for Barred Owl was a new high count for the state. Observers: Adam Beachy, Evan Beachy, Mikey Beachy, Seth Beachy, Joseph Byler, David Kinsinger, Ruben Stoll (compiler), and Victor Stoll.

Shelby County – 15 September; 0730-1830. Weather: partly cloudy; wind S 3-17 mph; 75-92° F. White Ibis and Buff-breasted Sandpiper were notable. Observers: Kenneth Bishop, Betty Blaylock, David Blaylock, Thomas Blevins, Chad Brown, Judy Dorsey, Margaret Jefferson, Georges McNeil, Mary Schmidt, Norman Soskel, Ed Thomas, Jim Varner, Jay Walko, and Jacob Wessels. Non-participating compiler was Dick Preston.

2018 Fall Bird Counts	Shelby Co.	Perry Co.	Montgomery Co.	Nashville	Anderson Co.	Blount Co.	Knox Co.	Elizabethton	Totals
Count Dates	15-Sep	9/29/2018	9/15/2018	9/28-30/2018	9/23/2018	9/15/2018	9/30/2018	9/29/2018	
Species									
Black-bellied Whistling-Duck	58	-	-	-	-	-	-	-	58
Canada Goose	14	76	31	108	179	64	286	781	1539
Wood Duck	18	8	28	17	-	13	19	79	182
Blue-winged Teal	114	27	-	3	-	7	27	110	288
Northern Shoveler	2	-	-	5	-	-	-	-	7
Gadwall	-	-	-	1	-	-	-	-	1
Mallard	46	1	-	121	51	14	136	345	714
Common Merganser	-	-	-	-	-	-	-	2	2
Northern Bobwhite	-	1	-	-	-	4	15	-	20
Ruffed Grouse	-	-	-	-	-	-	-	2	2
Wild Turkey	-	3	20	61	3	62	41	56	246
Pied-billed Grebe	-	-	-	9	-	2	-	32	43
Rock Pigeon	139	15	-	31	107	55	67	597	1011
Eurasian Collared-Dove	-	5	-	3	-	5	-	2	15
Mourning Dove	48	185	38	104	48	246	154	330	1153
Yellow-billed Cuckoo	1	-	3	4	-	1	2	4	15
Black-billed Cuckoo	-	1	-	1	-	-	-	-	2
Common Nighthawk	-	-	2	-	-	2	-	1	5
Chimney Swift	4	32	15	100	67	49	72	481	820
Ruby-throated Hummingbird	59	34	10	27	11	31	11	31	214
Virginia Rail	-	-	-	-	-	-	-	1	1
Sora	-	-	-	-	-	1	1	-	2
Purple Gallinule	-	-	-	-	-	-	-	1	1
American Coot	-	-	-	8	-	-	-	-	8

2018 Fall Bird Counts	Shelby Co.	Perry Co.	Montgomery Co.	Nashville	Anderson Co.	Blount Co.	Knox Co.	Elizabethton	Totals
Black-necked Stilt	18	-	-	-	-	-	-	-	18
Killdeer	35	25	12	31	16	6	41	43	209
Least Sandpiper	980	-	-	-	1	-	-	-	981
White-rumped Sandpiper	2	-	-	-	-	-	-	-	2
Buff-breasted Sandpiper	1	-	-	-	-	-	-	-	1
Pectoral Sandpiper	16	-	-	-	1	-	-	-	17
Semipalmated Sandpiper	20	-	-	-	-	-	-	-	20
Western Sandpiper	1	-	-	-	-	-	-	-	1
American Woodcock	-	-	1	4	-	-	-	-	5
Wilson's Snipe	-	-	-	-	1	-	2	-	3
Spotted Sandpiper	1	-	-	1	-	-	1	1	4
Solitary Sandpiper	-	-	-	-	1	-	-	-	1
Lesser Yellowlegs	52	-	-	-	-	-	-	-	52
Greater Yellowlegs	8	-	-	-	-	-	-	-	8
Black-legged Kittiwake	-	-	-	-	-	-	-	1	1
Ring-billed Gull	-	-	-	24	-	-	-	-	24
Herring Gull	-	1	-	-	-	-	-	-	1
Caspian Tern	-	-	-	-	-	-	-	1	1
Common Tern	-	-	-	-	-	-	-	2	2
Double-crested Cormorant	3	7	1	78	13	12	29	78	221
American White Pelican	-	-	-	1	-	-	-	-	1
Least Bittern	-	-	-	1	-	-	-	-	1
Great Blue Heron	12	15	1	48	8	13	26	49	172
Great Egret	14	28	1	26	1	1	4	2	77
Little Blue Heron	-	-	2	-	-	-	-	-	2
Green Heron	-	6	2	-	2	3	2	2	17

2018 Fall Bird Counts	Shelby Co.	Perry Co.	Montgomery Co.	Nashville	Anderson Co.	Blount Co.	Knox Co.	Elizabethhton	Totals
Black-crowned Night-Heron	-	-	-	-	-	3	2	5	10
Yellow-crowned Night-Heron	5	-	-	-	-	3	-	-	8
White Ibis	68	-	-	-	-	-	-	-	68
Black Vulture	2	112	52	195	27	17	90	77	572
Turkey Vulture	6	178	31	81	15	51	49	203	614
Osprey	-	4	-	5	1	2	6	27	45
Northern Harrier	-	1	2	2	-	-	1	1	7
Sharp-shinned Hawk	-	2	-	1	-	-	3	3	9
Cooper's Hawk	2	1	1	4	1	2	3	9	23
Bald Eagle	-	9	2	5	1	2	4	8	31
Red-shouldered Hawk	2	13	1	17	3	7	4	6	53
Broad-winged Hawk	-	70	1	427	1	4	75	321	899
Red-tailed Hawk	4	9	2	14	1	12	6	19	67
Buteo sp.	-	-	-	-	-	-	1	-	1
Eastern Screech Owl	-	3	-	5	-	2	3	24	37
Great Horned Owl	-	6	1	5	-	2	5	10	29
Barred Owl	2	25	2	6	-	-	1	5	41
Northern Saw-whet Owl	-	-	-	-	-	-	-	1	1
Belted Kingfisher	36	14	1	16	5	7	23	32	134
Red-headed Woodpecker	10	9	3	4	1	2	9	5	43
Red-bellied Woodpecker	48	50	17	81	33	38	100	92	459
Yellow-bellied Sapsucker	-	4	-	4	-	-	1	5	14
Downy Woodpecker	22	35	11	57	16	20	32	59	252
Hairy Woodpecker	-	10	6	15	3	2	4	12	52
Northern Flicker	2	26	-	48	4	5	31	67	183

2018 Fall Bird Counts	Shelby Co.	Perry Co.	Montgomery Co.	Nashville	Anderson Co.	Blount Co.	Knox Co.	Elizabethton	Totals
Purple Martin	-	-	2	-	-	-	-	-	2
Tree Swallow	-	1	-	5	-	3	91	465	565
Northern Rough-winged Swallow	6	18	-	47	-	-	151	3	225
Cliff Swallow	2	-	-	-	-	-	-	3	5
Barn Swallow	2	5	30	6	-	-	6	1	50
Carolina Chickadee	39	73	29	126	48	76	119	177	687
Tufted Titmouse	29	75	20	100	48	59	80	133	544
Red-breasted Nuthatch	-	-	-	3	-	2	-	4	9
White-breasted Nuthatch	22	14	8	55	32	18	18	71	238
Brown-headed Nuthatch	-	-	-	-	2	10	5	-	17
Brown Creeper	-	-	-	-	-	-	-	1	1
House Wren	-	1	-	9	1	1	4	4	20
Winter Wren	-	-	-	-	-	-	-	1	1
Sedge Wren	-	1	-	-	-	-	-	-	1
Marsh Wren	-	14	1	3	-	-	2	-	20
Carolina Wren	45	110	4	118	107	79	182	218	863
Blue-gray Gnatcatcher	1	8	1	5	-	5	-	-	20
Golden-crowned Kinglet	-	-	-	-	-	-	-	8	8
Ruby-crowned Kinglet	-	2	-	2	-	-	1	-	5
Eastern Bluebird	26	168	17	65	51	94	112	187	720
Veery	-	6	-	7	1	1	-	-	15
Gray-cheeked Thrush	-	55	-	14	3	-	-	12	84
Swainson's Thrush	3	77	2	30	13	7	2	46	180
Hermit Thrush	-	-	-	1	-	-	-	-	1
Wood Thrush	3	20	2	15	6	6	-	9	61
American Robin	51	146	8	102	11	91	88	591	1088

2018 Fall Bird Counts	Shelby Co.	Perry Co.	Montgomery Co.	Nashville	Anderson Co.	Blount Co.	Knox Co.	Elizabethhton	Totals
Gray Catbird	6	38	-	39	9	4	13	64	173
Brown Thrasher	1	36	3	22	23	11	27	23	146
Northern Mockingbird	15	52	3	57	31	50	156	111	475
European Starling	514	113	33	530	206	250	1172	1226	4044
Cedar Waxwing	-	28	14	68	231	71	11	294	717
House Sparrow	18	50	5	28	-	13	13	69	196
House Finch	6	18	5	22	6	33	76	41	207
Purple Finch	-	-	-	13	-	-	-	-	13
Pine Siskin	-	-	-	-	-	-	-	11	11
American Goldfinch	8	33	37	57	39	49	72	220	515
Eastern Towhee	2	-	7	45	9	-	36	66	165
Chipping Sparrow	15	164	26	15	2	32	2	94	350
Field Sparrow	-	35	2	23	11	19	60	16	166
Savannah Sparrow	-	2	-	-	-	-	-	-	2
Song Sparrow	-	-	-	7	4	13	51	97	172
Swamp Sparrow	-	2	-	2	-	11	-	-	15
White-throated Sparrow	-	-	-	1	-	-	-	-	1
Dark-eyed Junco	-	-	-	-	-	29	-	42	71
Yellow-breasted Chat	-	-	1	-	-	1	-	-	2
Bobolink	-	-	2	-	-	-	-	2	4
Eastern Meadowlark	-	9	-	8	-	-	16	7	40
Orchard Oriole	-	1	-	-	-	-	-	-	1
Baltimore Oriole	1	1	-	2	-	-	-	-	4
Red-winged Blackbird	8	-	5	3	5	365	145	132	663
Brown-headed Cowbird	-	-	-	4	-	-	83	5	92


2018 Fall Bird Counts	Shelby Co.	Perry Co.	Montgomery Co.	Nashville	Anderson Co.	Blount Co.	Knox Co.	Elizabethhton	Totals
Common Grackle	2	8	15	324	-	5	13	8	375
Ovenbird	-	9	-	-	-	-	1	-	10
Worm-eating Warbler	-	1	-	-	-	-	-	1	2
Northern Waterthrush	-	5	-	-	-	-	-	1	6
Golden-winged Warbler	-	2	-	2	-	-	-	-	4
Blue-winged Warbler	1	1	-	3	-	-	-	-	5
Brewster's Warbler	-	-	-	1	-	-	-	-	1
Black-and-white Warbler	-	15	-	16	1	1	-	7	40
Prothonotary Warbler	-	4	-	-	-	-	-	-	4
Tennessee Warbler	-	154	1	51	2	2	14	42	266
Orange-crowned Warbler	-	1	-	2	-	-	1	2	6
Nashville Warbler	-	1	-	3	-	-	-	-	4
Mourning Warbler	-	2	-	-	-	-	-	-	2
Kentucky Warbler	-	1	-	-	-	-	-	2	3
Common Yellowthroat	-	53	5	26	2	4	39	9	138
Hooded Warbler	6	6	-	1	8	3	1	7	32
American Redstart	15	32	-	46	6	7	8	16	130
Cape May Warbler	-	-	-	-	-	-	3	6	9
Cerulean Warbler	1	-	-	-	-	-	-	-	1
Northern Parula	32	32	3	8	-	-	-	8	83
Magnolia Warbler	1	125	-	92	11	1	12	18	260
Bay-breasted Warbler	-	2	-	16	1	-	5	15	39
Blackburnian Warbler	-	2	-	6	-	-	1	8	17
Yellow Warbler	-	-	-	-	1	-	-	1	2
Chestnut-sided Warbler	-	10	-	9	2	1	1	1	24
Black-throated Blue Warbler	-	-	-	-	-	1	-	4	5

2018 Fall Bird Counts	Shelby Co.	Perry Co.	Montgomery Co.	Nashville	Anderson Co.	Blount Co.	Knox Co.	Elizabethton	Totals
Palm Warbler	-	53	1	8	-	8	103	63	236
Pine Warbler	1	10	3	4	1	9	4	11	43
Yellow-rumped Warbler	-	3	-	11	-	-	1	-	15
Yellow-throated Warbler	1	2	-	6	-	-	5	1	15
Prairie Warbler	-	1	-	-	-	1	-	1	3
Black-throated Green Warbler	3	9	-	13	1	-	-	7	33
Canada Warbler	-	-	-	1	-	-	-	-	1
Wilson's Warbler	-	1	-	2	-	-	1	-	4
Warbler sp.	-	-	-	-	-	1	-	-	1
Summer Tanager	25	49	12	32	5	4	3	2	132
Scarlet Tanager	-	33	-	3	6	7	-	11	60
Northern Cardinal	59	232	62	164	87	65	165	149	983
Rose-breasted Grosbeak	-	598	-	85	8	1	26	125	843
Blue Grosbeak	-	5	1	4	-	1	3	-	14
Indigo Bunting	7	309	30	52	2	7	76	14	497
Total individuals	2886	5012	791	5089	1710	2621	4924	9283	32316
Total species	88	125	74	125	81	101	106	127	188
Observers	14	8	3	29	8	20	35	50	167
Parties	8	5	2	10	6	13	13	13	70
Party hours	29.5	48	16.5	61.35	20.04	67.42	65.28	104	412.09
Hours by car	12	18	11	15.8	6.67	18.09	14.8	-	96.36
Hours by foot	17.5	30	5.5	45.55	13.37	49.33	50.48	-	211.73
Hours by boat or other	-	-	-	-	-	-	-	-	-
Miles by car	34.5	101	116.5	309.7	78.6	194	121	-	955.3
Miles by foot	6.75	21.5	1.5	30.09	13.02	39.5	49.82	-	162.18

2018 Fall Bird Counts	Shelby Co.	Perry Co.	Montgomery Co.	Nashville	Anderson Co.	Blount Co.	Knox Co.	Elizabethton	Totals
Miles by canoe or other	-	-	-	-	-	-	-	-	-
Hours owling	-	4.5	0.5	3.17	-	2.17	0.67	9	20.01
Miles owling	-	6.5	-	14.1	-	1	5	-	26.6
Feeder/yard watch observers	-	-	2	2	-	8	-	-	12
Feeder/yard watch hours	-	-	1.25	-	-	5.5	2.75	-	9.5

THE SUMMER SEASON

Richard L. Knight, Editor



1 June - 31 July 2018

Heat was the main weather theme this summer with temperatures above average statewide. Rainfall, however, varied somewhat by region, although many areas were on the dry side.

A diverse group of waterbirds from ducks to spoonbills made news this summer. Common Mergansers had a productive breeding season in East Tennessee, as another brood of young in June adds to three separate broods found during spring. This large duck appears to be reclaiming historic breeding range. With only three previous summer records of Horned Grebe in Tennessee, three sightings across three regions this summer was intriguing.

Three other waterbirds once considered vagrants to Tennessee have increased dramatically in recent years. Neotropic Cormorants at Cross Creeks National Wildlife Refuge and Duck River Unit of the Tennessee National Wildlife Refuge apparently were continuing from spring. First found in the state in 1993, this species now has been reported 31 times, of which 24 records (77%) have occurred since 2004. Neotropic Cormorants have shown expansion inland from Texas and Louisiana. How soon til they nest in Tennessee? Four Brown Pelicans this summer, scattered across three regions, constitute the most ever found in any year in Tennessee. There are now 24 records in the state, with 20 (83%) having occurred since 2002. This increase coincides with the species' recovery along the Gulf of Mexico and Atlantic coasts. Similarly, three sightings of Roseate Spoonbill this summer bring the state total to about 32 records, of which 27 (84%) have occurred since 2000. Both of these distinctive coastal species have shown a recent trend of increased wandering to inland areas.

Standard Abbreviations

ad - adult	max - maximum count
Co - County	m.ob. - many observers
Cr - Creek	Mtn - Mountain
ers - earliest reported sighting	NWR - National Wildlife Refuge
et al. - and others	R - River
im - immature	SP - State Park
L - Lake	WMA - Wildlife Management Area
lrs - latest reported sighting	yg - young

WESTERN COASTAL PLAIN REGION - - Few folks seemed to venture out in the heat this summer. Despite this, the season's highlights included a lingering Fulvous Whistling-Duck, a White Ibis, continued breeding by Western Kingbirds, and a trio of Bell's Vireos.

Duck - Coot: **Black-bellied Whistling-Duck:** thru season (max 129) Ensley (m.ob.). **Fulvous Whistling-Duck:** from May into Aug (1) Ensley (m.ob.). **Gadwall:** 4 Jun (pair) Phillipy Pits (MAG). **Blue-winged Teal:** 4 Jun (2 males) Phillipy Pits (MAG). **Northern Shoveler:** thru 10 Jun (1) Ensley (m.ob.). **Green-winged Teal:** thru 7 Jul (1) Ensley (m.ob.). **Hooded Merganser:** 27 Jun (1 hen, 3 yg) Obion Co (MAG). **American Coot:** 13 Jun (2) Beech L, Henderson Co (VS).

Plover - Tern: **Piping Plover:** 27 Jul (1) Fritz Landing (MAG). **Upland Sandpiper:** 28 Jul (1) Ensley (DDP). **Ruddy Turnstone:** 27 Jul (4) Fritz Landing (MAG). **Stilt Sandpiper:** 10 Jul (1) Ensley (MCT, GG), ers. **Sanderling:** 23 / 27 Jul (1 / 8) Tiptonville Bar, Lake Co (MAG); 27 Jul (3) Fritz Landing (MAG). **Dunlin:** 24 Jul (1) Ensley (DDP). **Least Sandpiper:** 19 Jun (1) Phillipy Pits (MAG); 30 Jun (2) Ensley (MCT). **White-rumped Sandpiper:** 2 Jun (5) Ensley (Virginia Reynolds), lrs. **Pectoral Sandpiper:** 10 Jul (1) Ensley (MCT, GG), ers. **Semipalmated Sandpiper:** 19 Jun / 10 Jul (2 / 4) Ensley (Tom McNeil, Cathy McNeil / MCT, GG), lrs / ers. **Western Sandpiper:** 10 Jul (3) Ensley (MCT, GG), ers. **Short-billed Dowitcher:** 28 Jul (2) Ensley (DDP), ers. **Willet:** 27 Jul (1) Fritz Landing (MAG). **Ring-billed Gull:** 18 / 28 Jul (23 / 260) Paris Landing SP, Henry Co (JH, SA). **Least Tern:** 27 Jul (325 +) Tiptonville Bar, Lake Co (MAG), max. **Caspian Tern:** 13 Jun (1) Beech L, Henderson Co (VS); 21 Jun (1) Paris Landing SP, Henry Co (JH, DR); 16 Jul (1) Gibson Co L (MAG).

Anhinga - Falcon: **Anhinga:** 30 Jun (8) Ensley (MCT). **American White Pelican:** 9 Jun (17) Britton Ford, Henry Co (JH); 3 Jul (2) Reelfoot L, Lake Co (MAG); 21 Jul (2) Eagle Cr, Henry Co (SA, JH). **White Ibis:** 24 Jun (1 im) Hog Cr Bottoms, Decatur Co (RS). **Mississippi Kite:** 19 Jun (32) Hwy 641, Benton Co (Shawna Ellis); 27 Jun (15) Maness Swamp Refuge, Gibson Co (MAG). **Peregrine Falcon:** 9 Jun (1) Henry Co (JH).

Flycatcher - Bunting: **Willow Flycatcher:** 5 Jun (1) T.O. Fuller SP, Shelby Co (Bob Foehring); 8 Jun (1) Milan, Gibson Co (Maureen Ellis); 1 Jul (1) Shelby Co (Andrew Meiborg). **Western Kingbird:** thru Jul (4+) President's Island, Shelby Co (m.ob.), with 1 ad + 4 yg on 25 Jun (Bob Zaremba); thru Jul (2-3) Ensley (m.ob.), with a pair + 1 yg on 10 Jul (MCT, GG); 7 Jul (5) North Treatment Plant, Shelby Co (RS). **Scissor-tailed Flycatcher:** 9 Jun (1) Ensley (David Grant). **Bell's Vireo:** 14 Jun (3) Bogota WMA, Dyer Co (RS, VS), continuing from May. **Lark Sparrow:** thru Jun and Jul (1-2 at several sites) Gibson Co (MAG); 4 Jun (2) Fayette Co (Rob Harbin); 6 / 29 Jun (1) Crockett Co (Damien Simbeck); 6 Jun (2) Haywood Co (Damien Simbeck). **Painted Bunting:** 10 Jul (2 males, 1 female) Ensley and (1 male) TVA Lake, Shelby Co (MCT, GG).

Locations: Ensley - Shelby Co; Fritz Landing - Lake Co; Phillipy Pits - Lake Co.

HIGHLAND RIM AND BASIN - - Weather this summer has been hot and relatively dry. June temperatures averaged 4 degrees above normal, while July was 2.3 degrees above normal. Precipitation amounts were slightly below normal both months. Highlights this summer included Neotropic Cormorant, Brown Pelican, White Ibis, Roseate Spoonbill, and Red Crossbill.

Duck - Coot: **Blue-winged Teal:** 20 Jul (1) Duck R Unit (AT), ers. **Gadwall:** thru 20 Jul (1-3) Duck R Unit (RS, AT). **Ring-necked Duck:** thru 15 Jul (1) Duck R Unit (RS, AT); thru 2 Jul (1) Pickwick L (AT et al.). **Lesser Scaup:** 13-22 Jun (1) Pickwick L (RS, AT). **Bufflehead:** thru 22 Jun (1) Pickwick L (AT, RS). **Hooded Merganser:** thru 2 Jul (1) Pickwick L (RS et al.); 5 Jul (1) Duck R Unit (RS). **Ruddy Duck:** 5-12 Jul (1) Duck R Unit (DR). **Northern Bobwhite:** 28 Jun (19) Fort Campbell, Stewart Co (DMo), max. **Pied-billed Grebe:** 30 May / 30 Jun (1-2) Old Hickory L, Davidson Co (Michael Smith, Phillip Casteel / JH); thru 2 Jul (1-4) Pickwick L (RS, AT et al.); 21 Jun (1) Cross Cr NWR (JH, DR). **Horned Grebe:** 17 Jul (1) Edgar Evins SP, DeKalb Co (Mark Taylor, photo). **American Coot:** 1 Jun / 27 Jul / 7 Aug (2 ad with 1-4 yg) Centennial Park, Davidson Co (Peter Kondrashov / Sandra Zabala / GG); thru 2 Jul (1-3) Pickwick L (AT, RS).

Plover - Tern: **Semipalmated Plover:** 3 Jul (1) Duck R Unit (RS), ers. **Upland Sandpiper:** 26 Jul (1) Duck R Unit (refuge staff). **Stilt Sandpiper:** 20 Jul (1) Duck R Unit (AT), ers; 21 Jul (1) Fowler Rd, Lincoln Co (DSn). **White-rumped Sandpiper:** 10 Jun (1) Duck R Unit (RS, AT, David Kinsinger), lrs. **Semipalmated Sandpiper:** 10 Jun (1) Duck R Unit (RS, AT, David Kinsinger), lrs. **Long-billed Dowitcher:** 14 Jul (1) Barkley WMA, Stewart Co (JH). **Wilson's Snipe:** 8 Jul (1) Cross Cr NWR (SR, CR), ers. **Spotted Sandpiper:** 3 Jul (1) Duck R Unit (RS), ers. **Solitary Sandpiper:** 4 Jul (1) Dickson Co (DR), ers. **Lesser Yellowlegs:** 3 Jul (25) Duck R Unit (RS), ers. **Greater Yellowlegs:** 3 Jul (1) Duck R Unit (RS), ers. **Laughing Gull:** 13-14 Jun (1 ad) Pickwick L (RS, AT). **Ring-billed Gull:** 30 Jun (4) Drake's Cr, Old Hickory L, Sumner Co (GG). **Herring Gull:** 13-14 Jun (1 im) Pickwick L (RS, AT). **Least Tern:** 9 Jun (1) Duck R Unit (RS). **Caspian Tern:** 9-10 Jun / 5 Jul (1-4) Duck R Unit (AT, RS / DR); 13 Jun (2) Pickwick L (RS); 21 Jul (2) Fowler Rd, Lincoln Co (DSn). **Black Tern:** 21 Jun (1) Cross Cr NWR (JH, DR).

Loon - Falcon: **Common Loon:** 13 Jun (4) Pickwick L (RS); 26 Jun (1) Center Hill L, DeKalb Co (Mark Taylor); 26 Jun (1) Gray's Landing, Kentucky L, Stewart Co (DR). **Neotropic Cormorant:** 18 / 23 Jul (1 / 2) Cross Cr NWR (SA, photo); 21 Jul (1) Duck R Unit (SA, JH, photo). **Anhinga:** 21 Jun (1, flying over) Perry Co (RS). **Brown Pelican:** 13 Jun (1) Pickwick Dam (RS, photo); 30 Jun (1) Drake's Cr, Old Hickory L, Sumner Co (Paul Latour, m.ob., photo). **American White Pelican:** 12 Jun - 2 Jul (3-18) Pickwick L (RS, AT); thru Jul (up to 60) Duck R Unit (RS, AT). **Least Bittern:** 16 Jun / 24 Jul (1) Duck R Unit (DR). **Great Egret:** 12 Jul (550) Duck R Unit (DR), max. **White Ibis:** 6 / 10 Jul (1 im) Duck R Unit (RS, DR). **Roseate Spoonbill:** 3 Jul (2) Cross Cr NWR (DR, photo). **Mississippi Kite:** 2 Jun (1) Lawrence Co (CR, SR); 2 Jun - 30 Jul (6 reports of 1-3 birds) Montgomery Co (CR, SR, JH, Stefan Woltmann); 9 Jun (1) Duck R Unit (RS); 19 Jun - 13 Jul (3 reports) Dickson Co (RS,

DR, Jim Peters); 20 Jul (1) Johnsonville SP, Humphreys Co (AT). **Peregrine Falcon**: 31 Jul (1) Duck R Unit (refuge staff).

Flycatcher - Dickcissel: **Scissor-tailed Flycatcher**: 9 Jun (3, nest) Lytle Cr Rd, Rutherford Co (Justin Nation); 13 Jun (2 pair) Clifton, Wayne Co (RS); 13 Jun (5) Savannah Bottoms, Hardin Co (RS); 17 Jun (1, nest) Bobo Rd, Wilson Co (MCT); 22 Jun (2) Pickwick L (RS); 21 Jul (15, ad + im) Wartrace, Bedford Co (Melissa Turrentine), apparently 2-3 family groups. **Loggerhead Shrike**: 1 Jun (pair, 4 yg) Lincoln Co (SR, CR). **Bell's Vireo**: 4 Jun (1) Fort Campbell, Stewart Co (DMo). **Red Crossbill**: 4 Jun (10) Fort Campbell, Stewart Co (DMo); 24 Jun (1) Perry Co (RS). **Bachman's Sparrow**: 28 Jun (1) Fort Campbell, Stewart Co (DMo). **Lark Sparrow**: 4-20 Jun (1-3) Rush Rd, Franklin Co (DSn); 9 Jun / 30 Jul (2 / 4) Lincoln Co (DSn / Evan Buck); 13 Jun (4) Savannah Bottoms, Hardin Co (RS); 24 Jun (1) Nashville Speedway, Wilson Co (EBo); 29 Jun (1) Lytle Cr Rd, Rutherford Co (DR); 15 Jul (1) Nolensville, Williamson Co (Hugh Barger). **Henslow's Sparrow**: 4 Jun (3) Fort Campbell, Stewart Co (DMo); 11 Jun (4) Fort Campbell (DMo). **Hooded Warbler**: 1 Jul (1) Rutherford Co (Daniel Estabrooks). **American Redstart**: 16 Jun (6) White Oak Rd, Houston Co (RS, VS). **Cerulean Warbler**: 16 Jun (3) White Oak Rd, Houston Co (RS, VS). **Yellow Warbler**: 23 Jun (pair, carrying food and fecal sacs) Shelton Ferry WMA, Montgomery Co (Stefan Woltmann). **Dickcissel**: 4 Jun (15) Fort Campbell, Stewart Co (DMo), max.

Locations: Cross Cr NWR - Stewart Co; Duck R Unit - unit of Tennessee NWR, Humphreys Co; Fort Campbell - Montgomery Co (unless specified otherwise); Pickwick L - Hardin Co.

STEPHEN C. ZIPPERER, Murfreesboro, TN stczipperer@gmail.com

CUMBERLAND PLATEAU / RIDGE and VALLEY REGION - - This summer was hotter than normal. Rainfall amounts varied; for example, Johnson City received nearly twice as much rain as Tri-cities airport, just 15 miles away. On the whole, precipitation was slightly above average across the region.

A brood of juvenile Common Mergansers on the Watauga River in Washington County was the fourth brood found in East Tennessee this year. A Horned Grebe photographed on Norris Lake was among few summer records in the state. Shorebird highlights included a late north-bound White-rumped Sandpiper and two south-bound Willets. Two separate Laughing Gulls were seen in the region. Both North American pelican species occurred at Chickamauga Lake. Two reports each of Tricolored Heron and Roseate Spoonbill were remarkable, while two reports of White Ibis were more expected.

Nesting Scissor-tailed Flycatchers were observed in Marion, Bledsoe, and Cumberland counties. The latter represents a first breeding record in that county. Common Ravens in Scott and Morgan counties were notable. Impressive numbers of Dickcissels occurred this spring and summer.

Goose - Coot: **Greater White-fronted Goose:** 22 Jun (1) John Sevier L (SHu, photo). **Gadwall:** 12 Jun (1) Gupton Wetland, Roane Co (Ron Hoff, Dollyann Myers). **Blue-winged Teal:** 3-27 Jun (1 male) Brainerd Levee, Hamilton Co (Michael Yablick, BD); 24 Jun (pair) South Mouse Cr, Bradley Co (Rick Houlk). **Redhead:** 8 Jul (1 male) Watts Bar L, Rhea Co (RSh, DSh, photo). **Ring-necked Duck:** 12 Jun (1) Nickajack L, Marion Co (Nancy Robb, photo). **Hooded Merganser:** 14 Jun (2 females) Chester Frost Park, Hamilton Co (S.K. Jones). **Common Merganser:** 23 Jun (female with 7 yg) Watauga R, at Austin Springs, Washington Co (David Kirschke); 22 Jul - 4 Aug (1 female-type) South Holston R, Kingsport, Sullivan Co (Rack Cross, m.ob.). **Pied-billed Grebe:** 23 Jul (1) Ponderosa L, Scott Co (Betty Dunn), ers. **Horned Grebe:** 17 Jun (1, breeding plumage) Norris L, Union Co (Laura Tappan, photo). **Black-billed Cuckoo:** 9 Jun (1) Upper Buffalo Cr Valley, Scott Co (Charles Nicholson). **American Coot:** 14 Jun (1) Chester Frost Park, Hamilton Co (S.K. Jones); thru season (1, injured) Gupton Wetland, Roane Co (RoK, EBo).

Plover - Osprey: **Semipalmated Plover:** 20 Jun (1) John Sevier L (SHu). **Stilt Sandpiper:** 12 Jul (2) John Sevier L (SHu); 18 Jul (1) Eagle Bend Fish Hatchery, Anderson Co (Bates Estabrooks). **White-rumped Sandpiper:** 20 Jun (1) John Sevier L (SHu), Mrs. **Willet:** 14 Jul (1) Tennessee R, Hamilton Co (BD); 31 Jul (1) Paddle Cr pond, Sullivan Co (Rob Biller, Ron Carrico et al.). **Greater Yellowlegs:** 26-27 Jun (1) Brainerd Levee, Hamilton Co (BD), ers. **Laughing Gull:** 4 Jun (1 im) Chickamauga L (BD); 3 Jul (1 ad) L Tansi, Cumberland Co (EKL). **Ring-billed Gull:** 11 Jun (1) Fort Loudoun L, Loudon Co (David Vogt). **Caspian Tern:** 17 Jun (1) Fort Patrick Henry L, Sullivan Co (Bambi Fincher, Sherrie Quillen). **Forster's Tern:** 4 Jun (1) Chickamauga L (BD). **Brown Pelican:** 6 Jun (1 im) Chickamauga L (BD). **American White Pelican:** 18 Jun (14, flying over) Chickamauga L (S. K. Jones). **Least Bittern:** 8 / 31 Jul (1) John Sevier L (SHu). **Great Egret:** 23 Jun (30) John Sevier L (SHu), max. **Little Blue Heron:** 28 Jun - 2 Jul (1) Amnicola Marsh, Hamilton Co (BD, m.ob.); 13 Jul (1 im) John Sevier L (SHu); 16 Jul (1 im) Snow Hill, Hamilton Co (Rick Houlk); 17 Jul (1 im) Heritage Center Greenway, Roane Co (RoK); 27 Jul (1 im) Nolichucky R mouth, Hamblen Co (Ron Shrieves). **Tricolored Heron:** 25 Jul into Aug (1 im) John Sevier L (SHu); 30 Jul into Aug (1 im) Paddle Cr pond, Sullivan Co (Adrianna Nelson, m.ob.). **Cattle Egret:** 29 Jul (1) John Sevier L (SHu). **Yellow-crowned Night-Heron:** thru season (2 pair, nest) Maryville Greenway, Blount Co (Evan Kidd, m.ob.); 3 Jun (2) Baum's L, Knox Co (Susan Hoyle); 14-18 Jul (1) Sevierville, Sevier Co (Ryan Trenkamp, m.ob.). **White Ibis:** 4 Jul (1 sub-adult) Candies Cr, Meigs Co (Jeffrey Schaarschmidt et al.); 21 Jul (3 im) Birchwood, Meigs Co (Charles Murray). **Roseate Spoonbill:** 7 Jun (1) Grasshopper Rd, Hamilton Co (Charles Murray); 17 Jun (4) French Broad R, about a mile downstream from Seven Islands SP, Knox Co (Karen Wilkinson, photo). **Barn Owl:** 30 Jul (2 ad, 2 yg) Roane Co (RoK). **Osprey:** 30 Jun (3 nests with yg) Kingsport (Bambi Fincher, Sherrie Quillen), still a scarce breeder in NE Tenn.

Flycatcher - Dickcissel: **Scissor-tailed Flycatcher:** 20-21 Jun (pair, nest) Ketner Mill Rd, Marion Co (RSh, DSh / BD); 20 Jun (3) Swafford Chapel, Bledsoe Co (Dawn Greenway); 22 Jun / 13 Jul (pair, nest / 2 ad, 3 yg) Ninemile Crossroad, Bledsoe Co (RSh, DSh / EKL); 14-23

Jul (1-2, active nest) Dunbar Rd, Cumberland Co (EKL, Pam Barrett, photo). **Loggerhead Shrike**: 10 Jun (2 ad, 2 yg) Ooltewah, Hamilton Co (BD); 13 Jul (2 ad, 2 yg) Bledsoe Co (EKL). **Warbling Vireo**: 7 Jun (1, audio recording) Roane Co (RoK); 21 Jun (1) Mossy Cr, Jefferson Co (EBo). **Fish Crow**: thru season (1-3) Johnson City and Kingsport (m.ob.). **Common Raven**: 6 Jun / 8 Jul (1) Johnson City (RLK); 16 Jun (1) Bays Mtn, Sullivan Co (Sherrie Quillen); 18 Jun (1) Hwy 297, eastern Scott Co (Betty Dunn); 25 Jun (2) Frozen Head SP (RoK). **Veery**: 6 Jun (2) Frozen Head SP (RoK). **Savannah Sparrow**: thru season (pair) / 11 Jul (2 ad, 2 yg) Conklin, Washington Co (RLK), regular breeding site; 14 Jun (8) Morgan / Fentress Co line (RoK). **Nashville Warbler**: 16 Jul (1, singing) UT campus, Knox Co (Colin Sumrall). **Cerulean Warbler**: 26 Jul (2) Kingsport (PGR), migrants. **Blackburnian Warbler**: 15 Jun (1) Cross Mtn, Campbell Co (Bates Estabrooks). **Chestnut-sided Warbler**: 15 Jun / 8 Jul (1) Cross Mtn, Campbell Co (Bates Estabrooks / EBo). **Rose-breasted Grosbeak**: 6 Jun (3) Frozen Head SP (RoK). **Dickcissel**: widely reported following a strong spring showing; reports from 12 counties, mainly in southern part of region (Bledsoe, Blount, Bradley, Cumberland, Greene, Loudon, Marion, Meigs, Polk, Rhea, Roane, and Washington Cos); 12 Jun (10 at 3 sites) Bradley Co (RSh, DSh), max.

Locations: Chickamauga L - Hamilton Co; Frozen Head SP - Morgan Co; John Sevier L - Hawkins Co.

RICHARD L. KNIGHT, Johnson City, TN. rknight8@earthlink.net

EASTERN MOUNTAIN REGION - - It was a hotter than normal summer with rainfall slightly below normal. The two broods of Common Mergansers found in the spring were observed throughout the summer. A regional first Brown Pelican was found. Several high elevation breeders are highlighted in this report.

Merganser - Sapsucker: **Common Merganser**: 17 Jun (10) Metcalf Bottoms, Blount Co (Janie Finch); 31 Jul (10) Townsend, Blount Co (Randy Winstead). **Horned Grebe**: 4 Jul (2) South Holston L, Sullivan Co (Mike Sanders, Gary Cooper). **Mourning Dove**: thru season (2) Carver's Gap on Roan Mtn (RLK), regular site, but unusual at high elevation. **BROWN PELICAN**: 16 Jun (1) Watauga L, Carter Co (Odes Roberson, photo), first regional record. **Northern Saw-whet Owl**: 29 Jun (1 im) Tricorner Knob, GSMNP (MGH). **Red-headed Woodpecker**: 19 Jul (pair with 2 recently fledged yg) Watauga Point, Carter Co (RLK), second year nesting at this site. **Yellow-bellied Sapsucker**: 6 / 9 Jun (1) Roan Mtn (RLK); 9 Jun (3) Ripshin L, Carter Co (Fred Alsop, Kevin Brooks); 16 Jun (1) Unaka Mtn (Glen Eller, PGR, Harry Lee Farthing, Jerry Bevins); 16 Jun (2) Iron Mtn Gap, Unicoi Co (Bryan Stevens).

Flycatcher - Warbler: **Alder Flycatcher**: thru season (2) Carver's Gap on Roan Mtn (RLK et al.). **Hermit Thrush**: thru season (2-4) Roan Mtn (RLK); early Jun (3) Roger's Ridge, Johnson Co (Joe McGuiness); 16 Jun (4) Unaka Mtn (PGR); fairly common thru

season along the Appalachian Trail in the GSMNP (MGH). **House Finch**: 9 Jun (2 pairs, at feeders) Roan Mtn at 4450 ft (RLK), unusual at high elevation. **Red Crossbill**: 7 Jun (2) Clingman's Dome, GSMNP (MGH); 9 / 22 Jun (1 / 8) Roan Mtn (RLK / Larry McDaniel, Roy Knispel et al). **Pine Siskin**: thru season (2-6, at feeders) Roan Mtn at 4450 ft elevation (RLK et al.); 7 Jun (3) Clingman's Dome, GSMNP (MGH). **Chipping Sparrow**: 9 Jun (1) Roan Mtn at 6100 ft (RLK), unusual at high elevation. **Vesper Sparrow**: 6 Jun thru season (1) Round Bald on Roan Mtn (RLK et al.). **Grasshopper Sparrow**: 9 Jun (1) Elizabethton airport, Carter Co (Brookie Potter, Jean Potter); 16 Jun (1) near town of Unicoi, Unicoi Co (Joe McGuinness); few recent records in these two Cos. **Brown-headed Cowbird**: 9 Jun (1 female) Roan Mtn at 4450 ft (RLK), unusual at high elevation. **Prothonotary Warbler**: 8 Jun (1) Chota Refuge, Monroe Co (Shelley Rutkin). **Cerulean Warbler**: 28 Jul (1) Pittman Center, Sevier Co (Keith Watson), a migrant. **Magnolia Warbler**: 6 Jun thru season (1-2 males) Roan Mtn (RLK); 16 Jun (1 male) Unaka Mtn (Glen Eller, PGR, Harry Lee Farthing, Jerry Bevins); 30 Jun (1-2 males) Mt Guyot, GSMNP (MGH); 15 Jul (1 male) Yellow Cr Gap, GSMNP (MGH). **Yellow-rumped Warbler**: 6 Jun thru season (1 male) Roan Mtn (RLK); 7 Jun (6 males) within 2 miles of the summit of Clingman's Dome along the Appalachian Trail, GSMNP (MGH); 30 Jun (1) Tricorner Knob, GSMNP (MGH); 15 Jul (1) Mt Guyot, GSMNP (MGH); 15 Jul (1) Yellow Cr Gap, GSMNP (MGH).

Locations: GSMNP - Great Smoky Mtns National Park; Roan Mtn - Carter Co; Unaka Mtn - Unicoi Co.

RICHARD P. LEWIS, Bristol, TN. mountainbirds@gmail.com

Observers

SA - Sharon Arnold	DDP - Dick D. Preston
EBo - Eric Bodker	PGR - Peter G. Range
BD - Bruce Dralle	DR - Daniel Redwine
GG - Graham Gerdeman	CR - Cyndi Routledge
MAG - Mark A. Greene	SR - Steve Routledge
JH - Joe Hall	DSh - Debbie Shannon
MGH - Michael G. Harvey	RSh - Roi Shannon
SHu - Susan Hubley	DSn - Dan Snell
RLK - Richard L. Knight	RS - Ruben Stoll
RoK - Roger Kroodsma	VS - Victor Stoll
EKL - Edmund K. LeGrand	MCT - Michael C. Todd
DMo - Daniel Moss	AT - Alan Troyer

SPECIES INDEX TO VOLUME 89

- A**
Accipiter sp. 22, 60
Anhinga 33, 37, 102, 104, 110, 111, 138, 139
Avocet, American 32, 36, 40, 45, 103, 107
- B**
Bittern,
 American 34, 37, 41, 55, 56, 59, 72, 78, 104, 108, 110, 111
 Least 54, 55, 59, 104, 108, 126, 127, 129, 139, 141
Blackbird,
 Brewer's 14, 18, 26, 75, 79, 115
 Red-winged 26, 64, 133
 Rusty 26
 Yellow-headed 72, 79, 80, 83, 106, 109
Bluebird, Eastern 26, 62, 132
Blue Jay 24, 61, 131
Bobolink 38, 42, 43, 63, 109, 113, 133
Bobwhite, Northern 11, 14, 20, 32, 57, 128, 139
Booby, Brown 49, 52
Brant 30, 39, 40
Bufflehead 20, 45, 57, 139
Bunting,
 Indigo 35, 39, 43, 46, 65, 75, 106, 110, 135
 Painted 54, 56, 65, 72, 76, 79, 106, 138
Buteo sp. 24, 130
- C**
Canvasback 20, 44, 74, 77, 80, 81
Caracara, Crested 30, 32, 34, 94, 95, 101
Cardinal, Northern 29, 44, 65, 135
Catbird, Gray 15, 16, 17, 26, 62, 75, 83, 85, 133
Chat, Yellow-breasted 31, 38, 63, 113, 133
Chickadee,
 Black-capped 24, 62
 Carolina 24, 62, 132
Chuck-will's-widow 58, 111
Coot, American 22, 58, 128, 138, 139, 141
Cormorant,
 Double-crested 22, 33, 41, 52, 59, 111, 129
 Neotropic 32, 33, 35, 37, 102, 103, 104, 106, 107, 137, 139
Cowbird,
 Bronzed 30, 32, 35, 101
 Brown-headed 26, 64, 133, 143
Crane,
 Sandhill 18, 22, 32, 36, 40, 45, 74, 77, 82, 107
 Whooping 16, 18, 22, 36, 40, 72, 74, 77, 82
Creeper, Brown 24, 62, 132
Crossbill,
 Red 38, 40, 42, 46, 85, 102, 103, 105, 106, 109, 115, 139, 140, 143
Crow,
 American 24, 61, 131
 Fish 24, 34, 38, 42, 61, 78, 83, 102, 105, 108, 110, 112, 131, 142
Cuckoo,
 Black-billed 45, 55, 58, 103, 106, 110, 111, 114, 127, 128, 141
 Yellow-billed 58, 128
- D**
Dickcissel 38, 39, 65, 105, 106, 109, 110, 113, 114, 140, 141, 142
Dove,
 Common Ground- 77
 Eurasian Collared- 22, 57, 114, 128
 Mourning 22, 57, 128, 142
 White-winged 106, 110, 111
Dowitcher sp. 111
 Long-billed 33, 36, 40, 41, 44, 45, 74, 104, 107, 111, 139
 Short-billed 36, 41, 104, 107, 111, 138
- Duck,
 American Black 20
 Black-bellied Whistling- 32, 57, 102, 103, 106, 128, 138
 Fulvous Whistling- 102, 103, 138
 Long-tailed 14, 20, 32, 36, 72, 76, 77, 80, 81, 85, 106
 Ring-necked 20, 57, 139, 141
 Ruddy 15, 20, 139
 Duck sp. 20
 Wood 20, 57, 106, 128
Dunlin 15, 22, 33, 36, 41, 74, 77, 82, 104, 107, 111, 114, 138
- E**
Eagle,
 Bald 22, 31, 59, 74, 94, 130
 Golden 18, 19, 24, 31, 37, 40, 42, 44, 45, 54, 60, 75, 76, 78, 108, 115
Egret,
 Cattle 37, 59, 104, 108, 112, 115, 141
 Great 22, 41, 45, 59, 74, 78, 82, 115, 129, 139, 141
 Snowy 34, 37, 59, 104, 108, 112
Eider, Common 30, 35, 36
Empidonax sp. 61, 131
- F**
Falcon,
 Peregrine 14, 15, 16, 24, 31, 34, 38, 42, 44, 45, 60, 75, 78, 82, 85, 105, 108, 112, 115, 131, 138, 140
 Prairie 72, 73, 75
Finch,
 House 26, 63, 133, 143
 Purple 26, 34, 38, 40, 54, 63, 79, 83, 85, 109, 115, 126, 133
Flicker, Northern 24, 60, 130

- Flycatcher,
 Acadian 38, 61, 105, 115, 131
 Alder 56, 61, 105, 108, 115,
 131, 142
 Great Crested 60, 108, 131
 Least 14, 24, 61, 115, 131
 Olive-sided 34, 38, 42, 54, 60,
 105, 108, 112, 115, 126, 127,
 131
 Scissor-tailed 30, 34, 35, 38,
 105, 108, 112, 138, 140, 141
 "Trail's" spp 126, 131
 Willow 34, 38, 61, 131, 138
 Yellow-bellied 34, 38, 42, 45,
 105, 108, 131
- G**
- Gadwall 20, 35, 57, 126, 128, 138,
 139, 141
- Gallinule,
 Common 40, 72, 80, 82, 102,
 103, 107, 110, 111
 Purple 110, 111, 126, 128
- Gnatcatcher, Blue-gray 16, 24,
 34, 42, 62, 72, 75, 79, 80, 83,
 105, 109, 113, 115, 132
- Godwit,
 Hudsonian 41
 Marbled 102, 103, 104, 106,
 107
- Goldeneye, Common 20, 74,
 77, 81, 85
- Goldfinch, American 26, 63, 133
- Goose 20
 Cackling 17, 18, 20, 35, 73,
 76, 80, 84
 Canada 20, 128
 Greater White-fronted 15,
 17, 18, 20, 35, 40, 44, 73,
 76, 80, 84, 106, 110, 141
 Ross's 15, 16, 17, 18, 20, 32,
 35, 40, 44, 54, 55, 57, 76,
 80, 84
 Ross's X Snow 80
 Snow 19, 20, 32, 35, 40, 73,
 76, 80, 84, 106, 110
- Goshawk, Northern 14, 15, 22,
 31, 40, 42, 74, 76, 78
- Grackle,
 Common 26, 64, 134
 Great-tailed 100
- Grebe,
 Eared 45, 85, 102, 103, 110,
 111, 114
 Horned 22, 40, 45, 77, 81, 85,
 103, 111, 114, 137, 139, 140,
 141, 142
 Pied-billed 20, 57, 128, 139, 141
 Red-necked 72, 73, 74, 77,
 106
 Western 72, 80, 81
- Grosbeak,
 Blue 43, 46, 65, 110, 113, 135
 Rose-breasted 43, 46, 65, 105,
 110, 126, 135, 142
- Grouse, Ruffed 20, 54, 57, 128
- Gull,
 Bonaparte's 22, 41, 45, 82, 111
 California 30, 32, 33, 77, 80, 82
 Franklin's 33, 36
 Glaucous 14, 18, 22, 33, 37, 74,
 78, 102, 107
 Great Black-backed 33, 35, 37,
 72, 76, 78
 Herring 22, 33, 41, 59, 104, 115,
 127, 129, 139
 Herring X Glaucous 78
 Iceland 31, 74, 102, 106, 107
 Laughing 33, 36, 49, 111, 115,
 139, 140, 141
 Lesser Black-backed 22, 33, 36,
 72, 74, 78, 82, 104, 107
 Little 18, 22, 74
 Mew 72, 80, 82
 Ring-billed 22, 59, 82, 129, 138,
 139, 141
 Sabine's 30, 32, 33, 35, 36
 Thayer's 31
- H**
- Harrier, Northern 18, 22, 31, 37,
 54, 59, 78, 84, 85, 130
- Hawk,
 Broad-winged 31, 60, 105,
 115, 130
 Cooper's 22, 31, 44, 60, 130
 Ferruginous 72, 73, 75
 Red-shouldered 22, 31, 60,
 84, 85, 130
 Red-tailed 24, 31, 34, 37, 44,
 60, 74, 108, 122, 130
 Rough-legged 37, 74, 78,
 108
 Sharp-shinned 22, 31, 44,
 60, 130
 Swainson's 102, 106, 108,
 122, 123
- Heron 22
 Black-crowned Night- 22,
 45, 59, 82, 112, 130
 Great Blue 22, 59, 129
 Great White 30, 40, 41
 Green 22, 41, 59, 72, 78, 112,
 129
 Little Blue 37, 41, 104, 108,
 104, 112, 127, 129, 141
 Tricolored 140, 141
 Yellow-crowned Night- 42,
 59, 108, 112, 115, 126,
 130, 141
- Hummingbird,
 Black-chinned 35, 36
 Broad-billed 96, 101
 Ruby-throated 36, 58, 96,
 107, 111, 128
 Rufous 22, 36, 40, 73, 74,
 81, 97
- I**
- Ibis,
 Glossy 102, 106, 108
 Plegadis 80
 White 127, 130, 138, 139,
 140, 141
 White-faced 37, 40, 42,
 54, 55, 59, 102, 103, 104,
 110, 112, 114, 115,

- J**
- Jaeger 49
Parasitic 30, 35, 36, 100
- Jay, Blue 24, 61, 131
- Junco, Dark-eyed 26, 26, 44, 63, 133
- K**
- Kestrel, American 24, 31, 44, 60, 131
- Killdeer 22, 58, 82, 120, 121, 129, 126
- Kingbird,
Eastern 60, 108, 131
Western 34, 102, 105, 106, 108, 138
- Kingfisher, Belted 24, 60, 130
- Kinglet,
Golden-crowned 24, 62, 132
Ruby-crowned 26, 44, 62, 132
- Kiskadee, Great 101
- Kite,
Mississippi 34, 37, 40, 42, 44, 45, 59, 104, 108, 138, 139
Swallow-tailed 30, 32, 34, 35, 37, 40, 42, 117, 118, 119
- Kittiwake, Black-legged 126, 129
- Knot, Red 30, 40, 41, 44, 45, 102, 103, 104
- L**
- Lark, Horned 24, 61, 112
- Limpkin 99, 100
- Longspur, Lapland 15, 18, 26, 75, 79
- Loon,
Common 22, 32, 33, 37, 41, 45, 59, 73, 74, 78, 104, 107, 139
Pacific 33, 37, 74, 80, 82, 107, 111
Red-throated 22, 33, 37, 41, 74, 78, 82, 104, 107, 111
- M**
- Mallard 20, 57, 77, 128
- Martin, Purple 38, 61, 75, 78, 112, 132
- Meadowlark,
Eastern 26, 63, 133
Western 18, 26, 35, 38, 75, 79, 105
- Merganser,
Common 1, 2, 3, 4, 5, 6, 7, 8, 9, 16, 20, 40, 54, 57, 74, 76, 77, 81, 85, 100, 102, 110, 114, 128, 137, 140, 141, 142
Hooded 5, 20, 45, 57, 138, 139, 141
Red-breasted 20, 45, 57, 74, 77, 85, 106, 114
- Merlin 24, 31, 34, 37, 42, 44, 45, 60, 75, 78, 82, 85, 108, 112, 131
- Mockingbird, Northern 26, 63, 133
- N**
- Nighthawk, Common 58, 72, 80, 81, 128
- Noddy, Brown 30, 35, 37, 51, 106, 111
- Nuthatch,
Brown-headed 24, 62, 78, 89, 90, 91, 92, 93, 113, 132
Red-breasted 24, 34, 38, 40, 42, 62, 75, 78, 83, 85, 108, 112, 132
White-breasted 24, 62, 132
- O**
- Oriole,
Baltimore 63, 109, 133
Orchard 63, 105, 113, 133
Hooded 97
- Osprey 15, 18, 22, 31, 37, 59, 74, 78, 82, 98, 112, 126, 130, 141
- Ovenbird 44, 46, 64, 72, 80, 83, 113, 134
- Owl,**
Barn 11, 14, 15, 18, 24, 45, 60, 141
Barred 24, 60, 127, 130
Eastern Screech- 24, 60, 130
Great Horned 24, 60, 130
Long-eared 46, 75
Northern Saw-whet 37, 44, 45, 115, 130, 142
Short-eared 15, 18, 24, 37, 44, 45, 72, 73, 75, 76, 78, 85, 84, 103, 105, 112, 115
Snowy 78
- P**
- Parula, Northern 35, 64, 105, 109, 134
- Pelican,
American White 22, 33, 37, 41, 52, 59, 74, 78, 82, 102, 104, 107, 111, 129, 138, 139, 141
Brown 40, 41, 72, 82, 137, 139, 141, 142
- Petrel, Black-capped 30, 35, 37, 49, 50, 51
- Phalarope,
Red 36, 40, 41
Red-necked 36, 44, 45
Wilson's 33, 41, 104
- Phoebe,
Eastern 24, 61, 131
Say's 30, 32, 34
- Pigeon, Rock 22, 57, 128
- Pintail, Northern 20, 35, 44, 77, 84
- Pipit, American 26, 75, 85
Plegadis sp. 37, 82, 104
- Plover,
American Golden- 32, 36, 40, 58, 103, 107
Black-bellied 32, 36, 40, 45, 103, 107
Piping 32, 138
Semipalmated 40, 58, 104, 107, 111, 114, 139, 141
Snowy 100

- R**
- Rail,
 Virginia 19, 22, 32, 36, 40,
 103, 107, 110, 111, 114, 128
 Yellow 30, 32, 35, 36
- Raven 110, 112, 115
 Common 24, 40, 42, 61, 83,
 85, 102, 112, 115, 131, 142
- Redhead 18, 20, 40, 77, 80,
 81, 84, 141
- Redpoll, Common 30, 40, 42,
 72, 73, 75
- Redstart, American 64, 109,
 113, 134, 140
- Robin, American 26, 62, 132
- S**
- Sanderling 32, 36, 41, 45, 104,
 138
- Sandpiper,
 Baird's 33, 36, 41, 104, 107
 Buff-breasted 33, 36, 41, 127,
 129
 Least 14, 22, 58, 74, 77, 82,
 129, 138
 Pectoral 33, 36, 41, 58, 104,
 111, 129, 138
 Semipalmated 104, 114, 129,
 138, 139
 Solitary 14, 15, 22, 58, 104,
 107, 111, 129, 139
 Spotted 15, 22, 33, 45, 58,
 77, 107, 129, 139
 Stilt 32, 58, 104, 107, 111, 138,
 139, 141
 Upland 32, 36, 104, 107, 138,
 139
 Western 58, 107, 129, 138
 White-rumped 33, 41, 58,
 104, 107, 110, 111, 114, 129,
 138, 139, 140, 141
- Sapsucker, Yellow-bellied 24,
 60, 130, 142
- Scaup,
 Greater 17, 20, 55, 57, 74,
 77, 80, 81, 84, 114
 Lesser 20, 74, 77, 81, 110, 139
- Scoter,
 Black 20, 32, 36, 44, 73, 74,
 77, 81, 103, 106, 110
 Surf 14, 20, 32, 36, 40, 74,
 77, 81, 84, 103, 106, 110
 White-winged 14, 20, 32, 44,
 74, 77, 81, 84, 106, 110
- Shoveler, Northern 20, 44, 57, 128,
 138
- Shrike, Loggerhead 18, 24, 38,
 54, 61, 115, 126, 140, 142
- Siskin, Pine 26, 34, 38, 40, 42,
 46, 63, 72, 75, 79, 80, 83, 85,
 109, 113, 115, 133, 143
- Snipe, Wilson's 22, 33, 41, 58, 129,
 139
- Sora 14, 17, 18, 22, 36, 40, 55,
 56, 58, 80, 82, 103, 107, 111,
 114, 126, 127, 128
- Sparrow,
 American Tree 26, 72, 75,
 76, 79, 80, 83
 Bachman's 109, 140
 Chipping 26, 63, 133, 143
 Clay-colored 35, 38, 40, 43
 Field 26, 63, 133
 Fox 26, 43
 Grasshopper 43, 46, 63, 113, 143
 Harris's 15, 26, 72, 76, 79, 80,
 83, 102, 106, 109, 110, 113
 Henslow's 34, 38, 63, 105,
 109, 140
 House 26, 63, 133
 Lark 34, 38, 80, 83, 102, 105,
 109, 113, 138, 140
 LeConte's 15, 26, 34, 38, 75,
 79, 105, 109
 Lincoln's 17, 26, 38, 43, 46,
 75, 79, 109, 113
 Nelson's 30, 32, 34, 38, 109
 Savannah 26, 63, 133, 142
 Song 26, 63, 133
 Swamp 26, 63, 115, 133
 Vesper 17, 26, 34, 38, 43, 46,
 54, 56, 63, 79, 83, 109, 113,
 115, 143
 White-crowned 26, 46, 63
 White-throated 26, 63, 133
- Sparrow sp. 26
- Spoonbill, Roseate 34, 137, 139,
 140, 141
- Starling, European 26, 63, 133
- Stilt,
 Black-necked 36, 58, 103,
 107, 129
 Sandpiper 32, 36, 41
- Stork, Wood 33
- Swallow 61
 Bank 38, 46, 61, 112
 Barn 14, 18, 24, 34, 38, 62,
 72, 75, 108, 112, 132
 Cliff 34, 62, 108, 112, 132
 Northern Rough-winged 34,
 61, 108, 112, 132
 Tree 34, 46, 61, 72, 75, 78, 80,
 83, 85, 132
- Swan sp. 20, 73, 81
 Mute 17, 20, 73, 76, 80
 Trumpeter 72, 73, 76, 80
 Tundra 14, 16, 20, 35, 44, 73,
 76, 77, 81, 84, 106
- Swift, Chimney 36, 58, 103,
 106, 111, 128
- T**
- Tanager,
 Scarlet 13, 14, 16, 18, 29, 65,
 113, 135
 Summer 65, 105, 109, 113,
 135
 Western 13, 72, 76, 79
- Teal,
 Blue-winged 20, 40, 44, 57,
 77, 81, 128, 138, 139, 141
 Cinnamon 102, 106
 Green-winged 20, 57, 138
- Tern,
 Black 33, 37, 41, 45, 51, 104,
 107, 139
 Caspian 33, 37, 41, 45, 59,
 107, 111, 115, 129, 138,
 139, 141
 Common 33, 37, 41, 45, 51,
 104, 107, 115, 129
 Forster's 15, 16, 22, 33, 37,
 41, 45, 59, 74, 78, 82, 107,
 111, 141

- Tern,
 Least 33, 37, 40, 41
 Royal 30, 32, 33
 Sooty 30, 35, 37, 40, 41, 49
 Thrasher, Brown 26, 62, 133
 Thrush 44
 Gray-cheeked 62, 105, 109, 132
 Hermit 26, 46, 115, 132, 142
 Swainson's 44, 62, 105, 113, 132
 Wood 62, 105, 109, 113, 132
 Titmouse, Tufted 24, 53, 62, 132
 Towhee
 Eastern 26, 63, 133
 Spotted 72, 73, 75
 Turkey, Wild 11, 20, 57, 128
 Turnstone, Ruddy 32, 36, 41, 104, 138
- V**
- Veery 62, 132, 142
 Vireo,
 Bell's 38, 54, 55, 61, 102, 105, 108, 138, 140
 Blue-headed 15, 17, 24, 34, 61, 72, 75, 80, 83, 85, 112, 115, 131
 Philadelphia 38, 42, 45, 61, 108, 112, 115, 131
 Red-eyed 61, 105, 112, 131
 Warbling 61, 105, 131, 142
 White-eyed 14, 17, 24, 61, 72, 80, 83, 105, 108, 112, 131
 Yellow-throated 61, 105, 108, 112, 115, 131
 Vulture,
 Black 22, 31, 59, 130
 Turkey 22, 59, 130
- W**
- Warbler
 Bay-breasted 43, 64, 113, 134
 Black-and-white 64, 72, 80, 83, 109, 113, 134
 Blackburnian 39, 43, 65, 109, 115, 134, 142
 Blackpoll 35, 43, 46, 65, 109, 113
 Black-throated Blue 43, 44, 65, 109, 113, 115, 134
 Black-throated Green 35, 39, 43, 65, 105, 109, 113, 116, 135
 Blue-winged 39, 46, 64, 109, 113, 134
 Brewster's 56, 109, 113, 127, 134
 Canada 39, 56, 65, 113, 135
 Cape May 39, 44, 46, 64, 105, 109, 113, 134
 Cerulean 12, 43, 64, 109, 113, 115, 134, 140, 142, 143
 Chestnut-sided 39, 43, 65, 105, 134, 142
 Connecticut 39
 Golden-winged 35, 39, 43, 46, 64, 105, 109, 113, 115, 134
 Hooded 64, 97, 105, 113, 115, 134, 140
 Kentucky 64, 93, 105, 109, 134
 Kirtland's 30, 44
 Magnolia 35, 39, 43, 44, 64, 105, 134, 143
 Mourning 105, 105, 109, 113, 134
 Nashville 64, 109, 113, 134, 142
 Orange-crowned 18, 26, 39, 43, 46, 64, 75, 79, 80, 83, 134
 Palm 26, 39, 43, 65, 79, 84, 135
 Pine 26, 65, 85, 135
 Prairie 43, 65, 113, 115, 135
 Prothonotary 46, 55, 64, 105, 109, 113, 115, 134, 143
 Swainson's 54, 64, 105, 109, 113, 115
 Tennessee 35, 39, 43, 44, 64, 105, 109, 134
 Wilson's 39, 43, 65, 109, 127, 135
 Worm-eating 55, 64, 109, 113, 134
 Yellow 43, 65, 100, 109, 113, 134, 140
 Yellow-rumped 26, 65, 135, 143
 Yellow-throated 46, 65, 79, 105, 109, 113, 115, 135
 Warbler sp. 135
- Waterthrush,
 Louisiana 64, 109, 113
 Northern 39, 64, 113, 134
 Waxwing, Cedar 26, 63, 133
 Whip-poor-will, Eastern 45, 58, 106, 114
 Wigeon,
 American 20, 35, 114
 Eurasian 14, 15, 20, 76, 77, 100
 Willet 104, 107, 111, 138, 141
 Woodcock, American 22, 58, 77, 127, 129
 Woodpecker,
 Downy 24, 60, 130
 Hairy 24, 60, 130
 Pileated 24, 60, 131
 Red-bellied 24, 60, 130
 Red-headed 15, 24, 45, 60, 126, 130, 142
 Wood-Pewee, Eastern 42, 61, 112, 131
 Wren,
 Bewick's 38
 Carolina 24, 62, 132
 House 18, 24, 34, 62, 78, 83, 85, 132
 Marsh 15, 17, 18, 24, 38, 42, 46, 55, 62, 75, 79, 80, 83, 109, 113, 115, 127, 132
 Sedge 15, 17, 24, 34, 38, 42, 78, 84, 85, 115, 127, 132
 Winter 24, 38, 62, 132
- Y**
- Yellowlegs,
 Greater 22, 58, 74, 77, 82, 104, 107, 129, 139, 141
 Lesser 33, 36, 41, 58, 74, 82, 104, 107, 129, 139
 Yellowlegs sp. 59
 Yellowthroat, Common 18, 26, 35, 39, 43, 46, 64, 75, 109, 113, 134

INSTRUCTIONS TO AUTHORS

The Migrant records observations and studies of birds in Tennessee and adjacent areas.

SUBMISSIONS: The manuscript should be submitted electronically to Bob Ford at editorthemigrant@gmail.com. Submission of hard copies is optional. If so desired the original and two copies of the manuscript should be sent to the: Editor: Bob Ford, 808 Hatchie, Brownsville, TN 38012. Manuscripts that have been published in other journals should not be submitted.

MATERIAL: The subject matter should relate to some phase of Tennessee ornithology. It should be original, factual, concise and scientifically accurate.

STYLE: Both articles and short notes are solicited; recent issues of *The Migrant* should be used as a guide in the preparation of manuscripts. Where more detail is needed, reference should be made to *Scientific Style and Format*, eighth edition, by the Council of Science Editors, councilscienceeditors.org.

COPY: Manuscripts should be double-spaced with adequate margins for editorial notations and emailed in Word.docx. Tables and figures should be prepared in a separate file with appropriate headings; see *Scientific Style and Format* for examples of appropriate form for tables. Photographs intended for reproduction should be at least 300 dpi or sharp with good contrast on glossy white paper. Weights, measurements, and distances should be in metric units. Dates should be in “continental” form (e.g., 16 March 1997). Use the 24-hour clock (e.g., 0500 or 1900).

NOMENCLATURE: The scientific name of a species should be given after the first use of the full common name in the text. The scientific name should be italicized and in parentheses. Names should follow the *A. O. U. Check-list of North American Birds* (seventh edition, 1998, or supplements).

TITLE: The title should be concise, specific and descriptive.

ABSTRACT: Manuscripts of five or more pages should include an abstract. The abstract should be less than 5% of the length of the manuscript. It should include a brief explanation of why the research was done, the major results, and why the results are important.

LITERATURE CITED: List all literature citations in a “Literature Cited” section at the end of the text. Text citations should include the author and year.

IDENTIFICATION: Manuscripts including reports of rare or unusual species or of species at atypical times should include: date and time, light and weather conditions, exact location, habitat, optical equipment, distance, behavior of bird, comparison with other similar species, characteristic markings, experience of observer, other observers verifying the identification and reference works consulted.

REPRINTS: Reprints are available to authors on request. Billing to authors will be through the TOS treasurer. Request for reprints must be made well in advance of printing.

SEASON REPORTS: Observations that are to be considered for publication in “The Season” should be mailed to the appropriate Regional Compiler. Consult a recent issue of *The Migrant* for the name and address of the compiler.

CONTENTS

STATUS OF THE SWALLOW-TAILED KITE IN TENNESSEE	
Richard L. Knight.....	117
LATE NESTING KILLDEER IN HAMILTON COUNTY	
David Aborn.....	120
ROUND TABLE NOTE	
FIRST SWAINSON'S HAWK RECORD IN MIDDLE TENNESSEE	
Daniel Moss	122
MINUTES OF THE FALL TOS BOARD OF DIRECTORS MEETING	
Cyndi Routledge.....	124
2018 TENNESSEE FALL BIRD COUNTS	
Ron Hoff	126
THE SUMMER SEASON: 1 JUNE - 31 JULY 2018	
Richard L. Knight.....	137
WESTERN COASTAL PLAIN REGION	
Dick D. Preston.....	138
HIGHLAND RIM AND BASIN REGION	
Stephen C. Zipperer	139
CUMBERLAND PLATEAU/RIDGE AND VALLEY REGION	
Richard L. Knight.....	140
EASTERN MOUNTAIN REGION	
Richard P. Lewis.....	142
SPECIES INDEX TO VOLUME 89	144