Field Records July 1984



by George W. Gove, Robert H. Stymeist, Lee E. Taylor

The warm, wet, and cloudy month of July 1984 had an average temperature of 74.7 degrees. That was 1.2 degrees above normal, but still 3.3 degrees cooler than the blistering July of 1983. Ice cream cones were melting, and the beer was flowing on July 15 when the temperature reached 98 degrees, breaking a 101-year-old record for that date by 2 degrees. Seven days had the mercury soaring to 90 degrees or higher, two more days than average, but only half the number of days last year. The sun shone 63 percent of the time, which is the monthly average, but the nights were relatively comfortable, although the evening of July 23 was a sticky 79 degrees, the highest overnight low of the month. The lowest temperature was a mild 58 degrees on July 9, 10, 27, and 28. Precipitation totaled 4.43 inches, 1.75 inches more than normal; the most in any one day was 1.19 inches on July 18. No severe local storms were noted near Boston, but several tornadoes spiraled across other parts of Massachusetts, New Hampshire, and Connecticut.

LOONS THROUGH HERONS

A breeding-plumaged Red-throated Loon was observed at Stage Harbor in Chatham on the fifteenth. At Wareham 28 Common Loons were found in Little Harbor, nearly half of which were in summer plumage.

Twenty-five Cory's, 130 Greater, and just one Sooty shearwater were noted at Stellwagen Bank on the fifth. A single Greater Shearwater was observed sitting on the beach at Monomoy, being continually harassed by terns. Two immature Great Cormorants were noted near the target ship off Eastham, and Double-crested Cormorants were found in good numbers at East Boston, Squantum, and on Plum Island.

Fifty-three Great Blue Herons were counted at the Suasco Dam area in Westboro, a location where they nest. Tricolored Herons were found in just three locations, Plum Island, South Monomoy, and Nantucket. Likewise Little Blue Herons were few and far between with reports of three on South Monomoy and one or two from Essex County. At the spillway on the Charles River in Watertown, counts of Black-crowned Night-Herons were made several times during the month. On July 6, 97 were counted, including 11 immatures, and on July 14, 73 birds were recorded including just 7 immatures. This area has been a traditional gathering spot for night-herons; on July 1, 1976, 201 were tallied there.

R.H.S.

| SPECIES/DATE | LOCATION | NUMBER | OBSERVERS JULY 1984 |
|-----------------|-----------------------|---------------------|-------------------------|
| Red-throated Lo | on: | | |
| 15 | Chatham(Stage I.) | 1 br. pl. (ph.) | V.Cordes |
| Common Loon: | | | |
| 14,28 | Wareham(Little I.) | 20+, 28 | L.+C.Robinson |
| Pied-billed Gre | be: | | |
| 11 | S.Monomoy | 1 | B.Nikula# |
| Cory's Shearwat | er: | | |
| 5 | Stellwagen | 25 | R.Heil |
| Greater Shearwa | | | |
| 4,5 | N.Monomoy, Stellwagen | 1 sitting on beach, | 130 R.Heil |
| Sooty Shearwate | | | |
| 5 | Stellwagen | 1 | R.Heil |
| Great Cormorant | | 92 V3 = | |
| 10 | Eastham(target ship) | 2 imm. | A.Morgan |
| Double-crested | Cormorant: | | WE NO TO 1984 |
| thr. | E.Boston(Belle Isle) | max. 250 (7/22) | S.Zendeh# |
| 28,29 | P.I.area, Squantum | 350+, 300 | G.d'Entremont#,S.Zendeh |
| American Bitter | | | |
| 7,8 | P.I.,E.Boston | 1, 1 | R.Titus, S.Zendeh |
| 14,22 | Lancaster, W. Newbury | 2, 1 | S.Carroll#,R.Titus |
| | | | |

| SPECIES/DATE | LOCATION | NUMBER | OBSERVERS JULY 1984 |
|------------------|---------------------------|-------------------|-------------------------------|
| Least Bittern: | | | |
| thr. | P.I. | max. 3 | V.O. |
| 8,22 | Marshfield, GMNWR | 3-4, 2 imm. | W.Petersen#,M.Lynch# |
| Great Blue Heron | : | , - 1mm. | w.recerseng, n. Lynchy |
| 10 | Westboro(Suasco) | 53 | M.Lynch, S. Carroll |
| Great Egret: | | | M. Lynch, S. Calloli |
| 22;29 | IRWS; P.I., Westport | 3; 2, 20 | P Titue I Borry P Charlet |
| Snowy Egret: | | , -, 20 | R.Titus, J.Berry, R.Stymeist# |
| thr. | E.Boston(Belle Isle) | 59 (7/8) may 70 | (7/22 S. Zendeh |
| 2.7 | P.I., Squantum | 45, 40 | BBC,R.Abrams |
| Little Blue Hero | | , , , , | DBC, R. ADI aus |
| 1,6 | P.I., Rowley | 1, 1 | I.Giriunas, R. Forster |
| 11:29 | S. Monomoy; P.I., Ipswich | 3; 1 ad., 1 ad. | D.Holt#;G.Gove |
| Tricolored Heron | | ,, 1 001 | D.HOLE , G. GOVE |
| 1-2,11 | P.I., S. Monomoy | 1, 1 | I.Giriunas#,D.Holt |
| 18 | Nantucket | 1 | N.Jenks-Jay |
| Cattle Egret: | | | n.ochko ody |
| 4,29,30 | Ipswich | 3, 8, 3 | J.Berry, G.Gove, J.Berry |
| Green-backed Her | on: | | |
| thr.,14 | P.I.,Lancaster | max. 8, 11 | v.o.,M.Lynch# |
| Black-crowned Ni | ght-Heron: | | , |
| thr. | Watertown(Square) max. | 97+ incl. 11 imm. | (7/6) B.Phillips |
| 13,22 | Eastham(Hem, Landing) | 44, 68 | B.Nikula |
| 20,25 | Charlestown, Squantum | 19, 10 | T.Ahearn, G.d'Entremont |
| Yellow-crowned N | ight-Heron: | | , |
| 8,22 | N. Scituate, Eastham | 1 imm., 3 imm. | R.Titus, B.Nikula |
| 28,28-29 | N. Monomoy, P.I. | 1 imm., 1 ad. | CCBC, v.o. |
| Glossy Ibis: | | | 74.57.79.24.41.41.41 |
| thr. | P.I., E. Boston max. 9 | (7/21), max. 9 (7 | /22) E.Nielsen#,S.Zendeh |
| 4 | Natick(Broadmoor) | 1 | E.Taylor |
| 3,14 | N. Monomoy | 6, 8 | D.Holt#+v.o. |

WATERFOWL THROUGH NORTHERN BOBWHITE

Forty-six Wood Ducks were tallied at Great Meadows on July 22, and a female Pintail had eight young in tow on July 15. Unusual July waterfowl were Greater Scaup in Wareham, a pair of Bufflehead at Wachusett Reservoir in Clinton, and the interesting report of three young Hooded Mergansers in the Bolton Flat area of Lancaster.

On July 3, an immature Bald Eagle was seen feeding on carrion on the river flats between Woodbridge Island and the Sportsman's Lodge, Newburyport, occasionally getting up and flying around only to be harassed by gulls and terns, then settling back down to feed. Another immature Bald Eagle was observed flying very low over the Worcester Art Museum on July 27. In Newburyport, two young Cooper's Hawks were out of the nest on July 11; this nest, found in June, was the first known nesting of this species in many years for Essex County.

A <u>Mississippi</u> <u>Kite</u> was well observed on July 8 in Wareham. This sub-adult bird was the eighth record for the state, the most recent being a bird seen in Chatham and Truro on May 6 and 7, 1983.

R.H.S.

| SPECIES/DATE | LOCATION | NUMBER | OBSERVERS | JULY 1984 |
|-----------------|--------------------|----------------|------------------|-----------|
| Mute Swan: | | | | |
| 14 | P.I. | 1 | G.d'Entremont# | |
| Wood Duck: | | | | |
| 22 | GMNWR | 46 | BBC | |
| Northern Pintai | 1: | | | |
| 15 | P.I. | 1 f. + 8 yg. | G.Gove | |
| Blue-winged Tea | 1: | | | |
| 29 | P.I. | 30 + incl. yg. | J.Berry | |
| Gadwall: | | | | |
| 21,28 | P.I. | 7, 7 | BBC | |
| American Wigeon | : | | | |
| 11,29 | P.I. | 3, pr. | W.Petersen#,J.Be | rry |
| Greater Scaup: | | | | |
| 14 | Wareham(Little H.) | 1 m. | L.+ C.Robinson | |
| Oldsquaw: | | | | |
| 2 | Chatham | 3 | B.Nikula | |

| SPECIES/DATE | LOCATION | NUMBER | OBSERVERS JULY 1984 |
|-----------------|--|--|-----------------------------|
| Black Scoter: | | | |
| 14,21 | Wareham | 7 m., 5 m., 2 f. | L.+ C.Robinson |
| White-winged Sc | oter: | | |
| 14,21 | Wareham | 25, 58 | L.+ C.Robinson |
| Bufflehead: | | | |
| 1 | Clinton(Wachusett Reser | voir) pr. | R.Titus |
| Hooded Merganse | r: | | |
| 14 | Lancaster (Bolton Flats) | 3 yg. | M.Lynch, S. Carroll |
| Red-breasted Me | rganser: | | |
| 21,23 | Wareham, S. Dartmouth | 1 f., 1 f. | L.+ C.Robinson, R. Laubach |
| Turkey Vulture: | | | |
| 2,9 | Milton, Hopkinton | 2, 1 | W.Regan, P.Anderson |
| 10,20 | Dedham, Wayland | 1, 1 | J.Marshall, H. Parker |
| Osprey: | | | |
| 2,11 | Wareham, Lancaster | 3, 1 | L+C.Robinson,M.Lynch# |
| 21,29 | Newbypt.; M.V., Westport | 1, 7, 54 | BBC; A. Ellis, R. Stymeist# |
| Mississippi Kit | | | |
| 8 | Wareham | 1 sub-ad. | G.LeBaron |
| Bald Eagle: | | | |
| 3 | Newbypt. | 1 imm. | B.Sorrie |
| 27 | Worcester | 1 imm. | M. Lynch# |
| Cooper's Hawk: | | | |
| 11 | Newbypt. Pr. at r | nest + 2 yg. out of | nest. W.Petersen |
| Northern Goshav | | | |
| 1 | Lincoln | 1 ad. | R.Forster |
| Red-shouldered | Hawk: | | |
| 2.14 | E.Middleboro, N. Carver | 1, 1 imm. | K.Anderson |
| American Kestr | AND THE RESIDENCE OF THE PROPERTY OF THE PROPE | and the second s | |
| 21 | Plymouth Beach 2 ad. | + 3 m. + 2 f. yg. | G.Gove# |
| 29 | Lincoln(Hanscom) | 6+ | J.Carter |
| Northern Bobwh | | | |
| 3,29 | Canton, Bedford (Hanscom) | 1, 2 | G.d'Entremont, J. Carter |

RAILS THROUGH WOODPECKERS

On a BBC walk at Great Meadows, 10 Virginia Rails and 3 Soras were noted. Reports of Lesser Golden-Plover are scarce in summer; nevertheless, two were reported this month—one in breeding plumage. The highest counts in 12 years of records of American Oyster—catchers (40) and Willets (95) were recorded this month from North Monomoy, where a very high count of Least Sandpipers was also made. Also, excellent numbers of Whimbrels were counted there with 148 and 195 seen going to roost on July 19 and 30, respectively.

Among a flock of 500 Red Knots seen on Plymouth Beach were 10 to 12 color-banded birds. These were some of the knots that had been banded at Scituate (Third Cliff) in the summer of 1983. Three, and possibly four, <u>Curlew Sandpipers</u> were noted this month. The possibility of four birds exists because the bird seen at Squantum on July 23-29 may or may not have been the same bird that had been reported to Massachusetts Audubon Society on the twenty-seventh as having been seen at Thompson's Island for a week previous.

The Common Black-headed Gulls, which had failed in their nesting attempt in June, were still present on North Monomoy, where a Lesser Black-backed Gull in second summer plumage was also seen. An albino Great Black-backed Gull, at first mistaken for a Glaucous, caused excitement at Plymouth. A Gull-billed Tern was noted at Plum Island and a Caspian and 11 to 13 Royal Terns were reported, one of the latter being a banded bird. Last month it was said that Sandwich Terns are of irregular occurrence; this month 3 were seen at three different locations. Arctic Terns were present in good numbers. This is the southern limit of their breeding range but most of these birds were of the portlandica form; that is, first year, non-breeding birds. The largest and most productive Least Tern colony in the state, at Nantucket, had 560 pairs of adults and 310 immature birds. The pair of Black Skimmers nesting on North Monomoy was seen with three young; we hope the young successfully fledged and the skimmers will return to breed here next year.

No woodpeckers of any species were reported this month! This does not mean, however, that there are no woodpeckers in Massachusetts.

| | | | O. W. G. |
|-----------------------------------|-------------------------------|--|--|
| LOCATION | NUMBER | OBSERVERS | JULY 1984 |
| P.I. | 1 heard | BBC | |
| Lancaster, P.I. GMNWR, Chatham | 5, 1 f. + 5 yg. 10, 1 dead | M.Lynch#,BBC BBC,J.Aylward | |
| | P.I. Lancaster,P.I. | P.I. 1 heard Lancaster, P.I. 5, 1 f. + 5 yg. | P.I. 1 heard BBC Lancaster, P.I. 5, 1 f. + 5 yg. M.Lynch#, BBC |

| SPECIES/DATE | LOCATION | NUMBER | OBSERVERS JULY 1984 |
|--------------------------------|---|--------------------------------|--|
| Sora: | | | |
| 22 | GMNWR | 3 | BBC |
| Common Moorhen: | | | |
| thr. | GMNWR | 1-2 | V.O. |
| 11,29 Black-bellied Pl | S.Monomoy, P.I. | 1, 2 ad. + 6 yg. | B.Nikula#,J.Berry |
| thr. | N.Monomoy | max. 400 | B.Nikula# |
| 8,31 | Scituate, Plymouth | 12, 7 | W.Petersen, K. Anderson# |
| Lesser Golden-Pl | | 1200 Hall 120 | 1000000 120 II |
| 4,21 | N.Monomoy, Plymouth | 1 br. pl., 1 | G.Gove# |
| Semipalmated Plo thr. | N.Monomoy | max. 300 (7/28) | B.Nikula# |
| 8,28 | Scituate, P.I. | 2, 10 | W.Petersen,BBC |
| Piping Plover: | | | |
| 8,19 | Scituate | 8, 4 ad. + 6 yg. | |
| 26 Killdeer: | Monomoy | max. 15 | B.Nikula# |
| 1 | Halifax, Bedford | 50, 10 | W.Petersen, J. Carter |
| 10,21 | Sharon, P.I. | 16, 15 | R.Titus, BBC |
| American Oystero | | | |
| 14 | Nantucket, N. Monomoy | 7, 20 | A.Ellis,BBC |
| 21,26 Greater Yellowle | Chatham, N. Monomoy | 1, 40 | W.Reagan, B. Nikula# |
| thr. | | 32 (7/29) may 20 | (7/28) S.Zendeh, BBC |
| 14 | N. Monomoy | 100 | BBC S. Zeilden, BBC |
| Lesser Yellowleg | | | |
| thr. | E.Boston, P.I. max. | 40 (7/14), max. 21 | (7/21) S.Zendeh, v.o. |
| 19,20 22 | N.Monomoy,Halifax Newbypt | 130, 11 223 | B.Nikula#,K.Anderson |
| Solitary Sandpip | | 223 | R.Titus |
| 14-16 | P.I. | 1 | v.o. |
| 20,22 | Halifax, GMNWR | 3, 1 | K.Anderson, BBC |
| Willet: | | | |
| 2-28 10,28 | P.I. Barnstable, Squantum | max. 6-8 | v.o. |
| thr. | N.Monomoy | nest. pr., 8 max. 95 (7/28) | G.Wilson,W.Reagan B.Nikula# |
| Spotted Sandpipe | | mm. 33 (//20) | b.Nikulay |
| 14,15 | Lancaster, Belmont | 1 ad. + 1 yg., 2 | M.Lynch#,L.Taylor |
| 16,22 | P.I.,GMNWR | 4, 3 | BBC |
| 17,27 Upland Sandpiper | Nantucket, E. Boston | 2, 3 | N. Jenks-Jay, S. Zendeh |
| 1,4 | Halifax, E. Middleboro | 5, 2 | W.Petersen, K.Anderson |
| 15,29 | Newbypt, Bedford | 3, 19 | G.Gove, J. Carter |
| Whimbrel: | | total cere | |
| 12,19,30 | N. Monomoy | 40, 148, 195 | B.Nikula# |
| 15;21,31 1-2 birds fi | Nantucket; Plymouth rom each of 6 locations | 10; 8, 5 | N. Jenks-Jay; G. Gove, W. Smith |
| Hudsonian Godwit | | | |
| thr. | N. Monomoy | max. 63 (7/28) | v.o. |
| 7-29 | Newbypt-P.I. | max 8 (7/28) | v.o. |
| 8-29 | E.Boston | max. 11 (7/22) | S.Zendeh |
| Ruddy Turnstone: 12,25 | Nantucket | 2, 7 | N T1- T |
| 21,30 | Plymouth, Wareham | 20, 40 | N.Jenks-Jay G.Gove#,L+C.Robinson |
| Red Knot: | 150 (50) | | |
| thr. | N. Monomoy | max. 1400 (7/28) | |
| 19,29 21 | N.Scituate Plymouth | 150, 1100 500 | G.d'Entremont#, R.Titus |
| Sanderling: | Fiymoden | 300 | G.Gove# |
| thr. | N.Monomoy | max. 3000 (7/24) | B.Nikula# |
| 22,29 | E.Boston, Nantucket | 150, 46 | S.Zendeh, N. Jenks-Jay |
| Semipalmated Sand | | 0000 | The state of the s |
| thr. 22,28 | N.Monomoy E.Boston, P.I. | max. 2000 (7/24, 2 | |
| 29 | Scituate | 500, 75 400 | S. Zendeh, BBC |
| Least Sandpiper: | (C) 4 (C) | | R.Titus |
| thr. | N.Monomoy | max. 1400 (7/19) | B.Nikula# |
| 20,22 | Halifax, Newbypt | 67, 265 | K.Anderson, R. Titus |
| 29 Postoval Sandaina | Scituate | 150 | R.Titus |
| Pectoral Sandpipe 2,7,21,22 | r: P.I. | 1 1 1 1 | |
| 4 | N. Monomoy | 1, 1, 1, 1 | v.o. |
| 22,29 | E.Boston | 3, 2 | G.Gove S.Zendeh |
| | | (d) | w recorded to |

| SPECIES/DATE | LOCATION | NUMBER | OBSERVERS JULY 198 |
|----------------------------|--|-------------------------|---|
| Dunlin: | | 1 1 | |
| 15 Curlow Sandainer | P.I. | 1 br. pl. | G.Gove |
| Curlew Sandpiper: 14,22 | Nantucket, N. Monomoy | 1 br. pl., 1 | A E1114 - Y m1 - # |
| 22-29 | Squantum | 1 br. pl. | A.Ellis, L.Taylor# |
| Stilt Sandpiper: | 55.* SACOSSTANA | 26 MANUAL SANDA | |
| 6-29 | P.I. | max. 34 (7/28) | v.o. |
| Short-billed Dowi | tcher: | | |
| thr. | N.Monomoy | max. 2300 (7/14) | v.o. |
| thr. | P.I., Newbypt | max. 375 (7/22) | v.o. |
| thr. | E.Boston | max. 120 (7/8) | S.Zendeh# |
| 7,8 | Squantum, Scituate | 200, 25 | R.Abrams, W. Petersen |
| Long-billed Dowit | P.I. | 1 | U Patawan# |
| Common Snipe: | | • | W.Petersen# |
| 11 | Lancaster | 2 | S.Carroll# |
| American Woodcock | c: | | |
| 8;21,23 | Lincoln; Newbury | 1; 1 | J.Carter; BBC |
| Wilson's Phalarop | | | |
| 2,26 | P.I., N. Monomoy | 2, 1 | BBC,D.Holt# |
| Parasitic Jaeger: | | | |
| 4,6 | Marion | 5, 1 | P.Hallowell |
| Laughing Gull: | P.I. | 1 ad + 1 4mm 1 | hr. pl . C. C |
| 15,28 18,25 | Nantucket, Squantum | 1 ad. + 1 imm., 1 45, 5 | br. pl. G.Gove N.Jenks-Jay,G.d'Entremont |
| Little Gull: | rane deliver ; oquan cum | 75, 5 | N. Jenks-Jay, G. G Eneremone |
| 7,8 | P.I., Newbypt | 1 imm., 1 | R.Titus,BBC |
| Common Black-head | ded Gull: | | |
| thr. | N.Monomoy | 2-3 ad. | D.Holt# |
| 2,22 | E.Boston | 1 | S.Zendeh |
| Bonaparte's Gull | | 22 22 | 1995 A SO S 19 |
| 8 | P.I., Revere | 60, 90 | BBC,S.Zendeh |
| Ring-billed Gull | | 1 6 | |
| 23,30 Lesser Black-bacl | Westport | 1, 6 | R.+ C.Laubach |
| 22 | N.Monomoy | 1 (2 S) | C.Floyd# |
| | ed Gull (albino): | - (/ | 0.1.10,00 |
| 8 | Plymouth | 1 | W.Smith |
| Gull-billed Tern | : | | |
| 1 | P.I. | 1 | J.Soucy, L. Jodrey |
| Caspian Tern: | | | |
| 1 | P.I. | 1 | R.Williams |
| Royal Tern: 1-22 | N.Monomoy | max. 3 (7/4) | v.o. |
| 3,15 | Nantucket | 2, 3 | E.Andrews, N. Jenks-Jay |
| 8,10 | Revere, Plymouth | 2, 1 b. | S.Zendeh, W.Smith |
| 14 | P.I. | 2 | G.d'Entremont# |
| Sandwich Tern: | | | |
| 6,11 | P.I., S. Monomoy | 1, 1 (ph.) | R.Heil, B.Nikula# |
| 29 | Nantucket | 1 | N. Jenks-Jay |
| Roseate Tern: | | | |
| 4,15 | P.I. | 2, 1 br. pl. | G.d'Entremont, G. Gove |
| 8 | Nantucket, Scituate | 1, 4 | N.Jenks-Jay, W.Petersen |
| Common Tern: | N | /6 126 200 | N Jonks Jon |
| 8,29,30 | Nantucket | 46, 136, 200 | N.Jenks-Jay |
| Arctic Tern: | N. Monomoy | max. 80 (7/14) | B.Nikula# |
| thr. | Orleans | max. 25 (7/21) | B.Nikula# |
| 8 | Nantucket | 3 ad. + 2 imm. | N.Jenks-Jay |
| Forster's Tern: | | | |
| 23 | N. Monomoy | 1 | R.Humphrey# |
| Least Tern: | | | |
| thr. | P.I. | max. 67 (7/15) | v.o. |
| thr. | Nantucket | 1430 | N.Jenks-Jay |
| Black Tern: | 111111111111111111111111111111111111111 | 1 11 1 1 | C. Count |
| 4 | N. Monomoy | 1 br. pl. + 1 | G.Gove# N.Jenks-Jay |
| 8,18,25,29 | Nantucket | 2, 1, 1, 1 | n.Jenka-Jay |
| Black Skimmer: thr. | N.Monomoy 1 pr. with | 3 chicks, max. 6 | D.Holt# |
| thr. | Orleans | pr. | P.Trul1# |
| 8,25 | P.I., Naushon I. | 1, 2 | v.o.,J.Hatch |
| Black-billed Cuc | 그리는 전화보다 가장 아이를 바다 되었다면 하는 것이 없는 아이들이 되었다면 하는데 | | and the second second |
| 1,25 | Ipswich, S. Monomoy | 1, 1 | J.Berry, B. Nikula# |
| | ur locations. | | |

| SPECIES/DATE | LOCATION | NUMBER | OBSERVERS JULY 1984 |
|----------------|-----------------------|--------------|------------------------|
| Yellow-billed | Cuckoo: | | |
| thr. | E.Middleboro | 1-2 | K.Anderson |
| 22 | Newbypt | 2 | R.Titus |
| 6 from 6 | locations | | |
| Eastern Screec | h-Owl: | | |
| 29 | M.V. | 2 gray-phase | A.Ellis# |
| Great Horned O | wl: | | |
| 7 | P.I. | 1 | J.Berry# |
| Barred Owl: | | | |
| 1 | Lakeville, Medfield | pr., 4-5 | W.Petersen#, B.Cassie# |
| Short-eared Ow | 1: | | |
| thr. | N. Monomoy | 3 | D.Holt# |
| 25 | S.Monomoy | 5 | B.Nikula# |
| Northern Saw-w | thet Owl: | | |
| 7 | Wellesley | 1 | fide N.King |
| Common Nightha | wk: | | |
| 18 | Stoughton | 1 | R.Titus |
| Whip-poor-will | : | | |
| 2,4 | Newbury, Natick | 2, 3 | BBC |
| 14,23 | E.Middleboro, Newbury | 1, 2 | K.Anderson,BBC |
| Chimney Swift: | | | |
| 4 | Natick | 15 | BBC |
| Ruby-throated | Hummingbird: | | |
| 23 | E.Middleboro | 1 | P.Anderson |
| Belted Kingfis | her: | | |
| 22,23 | GMNWR, Randolph | 2, 1 | BBC,G.d'Entremont |

FLYCATCHERS THROUGH PURPLE FINCH

An individual Acadian Flycatcher seen in Wareham constituted an unusual summer location for this species, which is much more regular as a breeder in central and western Massachusetts. Another surprise from the southeastern part of the state was the nesting pair of Willow Flycatchers at Whitman; this was only the third breeding record in Plymouth County known to Wayne Petersen. Cliff Swallows started out the season auspiciously at the maintenance buildings on Plum Island, with a large number of nests. By mid-July, however, only two adults were present and the nests were all destroyed or abandoned. The apparent cause was competing House Sparrows. A Sedge Wren was heard and carefully observed during the first week of the month in Nonquitt, near New Bedford. At month's end an adult Swainson's Thrush was banded at Rockport.

July observers at coastal sites turned up some nice rarities. For three days beginning July 6, an immature Scissor-tailed Flycatcher was seen at Plum Island. Another report of what was very likely the same individual was received from Chatham a week later. Interestingly, of the two long tailed flycatcher species, this North American one has occurred less often in recent years than the South American Fork-tailed. Interest in the Jackdaw near Siasconset on Nantucket heightened when apparent plumage differences were detected. Indeed, on July 9 two Jackdaws were seen together at that site. This is not quite so astounding an occurrence in the light of other recent Jackdaw activity in North America. An article on this situation will be published in a later issue of this magazine.

| SPECIES/DATE | LOCATION | NUMBER | OBSERVERS JULY 1984 |
|-----------------|--------------------|---------------|--------------------------------------|
| Olive-sided Fly | catcher: | | |
| 1 | Medfield | 1 | W.Reagan |
| Eastern Wood-Pe | wee: | | |
| thr.,9 | E.Middleboro, P.I. | 1, 1 | K.Anderson,BBC |
| Acadian Flycatc | her: | | |
| 1 | Wareham | 1 | L.+ C.Robinson |
| Alder Flycatche | r: | | |
| 14.15 | Lancaster,Lincoln | 1, 1 | S.Carroll#, C.Floyd# |
| Willow Flycatch | er: | 1000 | Secretary tests - es secretary tests |
| 1 | Whitman, Lincoln | pr., 2 | W.Petersen, J. Carter |
| 14,22 | Lancaster, GMNWR | 8, 4 | M. Lynch#, BBC |
| Eastern Phoebe: | | | |
| 6 | Concord | 3 | J.Carter |
| Great Crested F | lycatcher: | | |
| 14 | Sharon | 3 | R.Titus |
| Eastern Kingbir | d: | | |
| 2,14 | Sharon, P.I. | 16, 10 | R.Titus, BBC |
| Scissor-tailed | | | |
| 6-8,14-15 | P.I., Chatham | 1 imm., 1 ad. | R.Heil + v.o.,L.Pivacek# |

| SPECIES/DATE | LOCATION | NUMBER | OBSERVERS JULY 1984 |
|------------------------------------|---------------------------------------|-------------------|---|
| Horned Lark: | Bedford | 3 | J.Carter |
| Purple Martin: | bearora | 3 | 5. Garter |
| 1-18 | Marshfield, Middleboro | 16 pr., 43 pr. | R.Turner |
| 1-18 | Halifax, Rehoboth | 12 pr., 17 pr. | R.Turner |
| 21-28 | | 30 max. | |
| | P.I. | 30 max. | v.o. |
| Northern Rough-wi 4,28 | Wayland, Stoughton | 7, 1 | R.Forster, R.Titus |
| Bank Swallow: | | | |
| 2-21,12 | P.I.,Lincoln | 6 max., 100+ | BBC, J. Carter |
| Cliff Swallow: | | | |
| 2-14,8 Blue Jay: | P.I., Marshfield | 2 max., 1 | v.o.,W.Petersen# |
| 23-25 Fish Crow: | Chatham | several flocks mi | gr. D.Holt |
| 7-22 | Canton | 2 max. | v.o. |
| Jackdaw: | W | 2 | N. Tooloo Tood |
| 9-31 | Nantucket | 2 | N.Jenks-Jay# |
| Red-breasted Nuth | | | 2 5 100 100 100 100 100 100 100 100 100 1 |
| 9,11 | Concord, Newbypt | 2, 1 | J.Carter, W.Petersen# |
| 25,26 | Framingham, E. Middleboro | 1, 1 | R.Forster, K.Anderson |
| Brown Creeper: | | | |
| 3-14 Sedge Wren: | Concord | pr. + 2 yg. | C.Floyd# |
| 6-7 | Nonquitt | 1 | A.Ellis |
| Marsh Wren: 2-21,22 | P.I.,GMNWR | 12 max., 17 | v.o.,BBC |
| Golden-crowned Ki | inglet: | | |
| from June-1 Eastern Bluebird: | Lakeville | 2 pr. | W.Petersen |
| 7 Veery: | Lexington | 1 | R.Titus |
| 15 | Bolton | 7. | R.Titus# |
| Swainson's Thrush 30 | Rockport | 1 ad. b. | R.Norris |
| Hermit Thrush: | Sharon | 2 | R.Titus |
| Gray Catbird: 22 | Newbypt | 37 | R.Titus |
| Solitary Vireo: | 2.70 | 1 | |
| 11 | Sharon | 1 | R.Titus |
| Yellow-throated V from June-1,1 | Halifax,Bridgewater | nesting pr., pr. | W.Petersen# |
| Warbling Vireo: 22 | GMNWR | 2 | BBC |
| Red-eyed Vireo: 14 | Sharon | 23 | R.Titus |
| Blue-winged Warb | | | |
| 2 Nashville Warble | r: | 1 m. | J.Carter |
| 1 Northern Parula: | Berlin | 1 | R.Titus |
| 11 Chestnut-sided Wa | W.Newbury | 1 m. | W.Petersen |
| 15 | Waltham | 1 m. | L.Taylor |
| Black-throated Gr 1,30 | reen Warbler: Lakeville,Middleboro | 1, 1 m. | W.Petersen#,K.Anderson |
| Pine Warbler: | Lancaster, Medfield | 2, 1 | R.Titus, W.Reagan |
| 3,14 Prairie Warbler: | E.Middleboro,Concord | 2 m., pr. | K.Anderson, J. Carter |
| 1,3 Black-and-white N | N.Carver,Concord | 1 m., 2 m. | K.Anderson,J.Carter |
| 14,21 | Sharon, Newbury | 4, 2 | R.Titus, BBC |
| Worm-eating Warb from May-14 | Sharon | 1 | R.Titus |
| Northern Waterth | rush: Medfield,Lakeville | 1, 5 | W.Reagan, W.Petersen# |
| Canada Warbler: | Lakeville | 2 pr. | W.Petersen# |
| Indigo Bunting: | Concord, Halifax | 5, 4 m. | J.Carter, K.Anderson |
| 3,20 | Concord, natitax | J, 4 m. | J. Garter, K. Anderson |

| SPECIES/DATE | LOCATION | NUMBER | OBSERVERS | JULY 1984 |
|-----------------|----------------------------------|------------------|-------------------------------------|-----------|
| Field Sparrow: | | | | |
| 3,20 | Concord, Halifax | 5, 1 m. | J. Carter, K. Ande | rson |
| Grasshopper Spa | irrow: | | | |
| 9,15 | Lincoln, M.V. | 3 ad. + 2 yg., 1 | J. Carter, M. Vaug | han |
| Sharp-tailed Sp | arrow: | | PERSONAL CONTRACTOR OF CASE OF CASE | |
| thr. | E.Boston, P.I. | 8-10, 4 max. | S.Zendeh, v.o. | |
| Seaside Sparrow | : | | | |
| 1, thr. | P.I., E. Boston | 2, 1-2 | I.Giriunas, S.Ze | ndeh |
| Swamp Sparrow: | Antan Trend to Mark State (1997) | | | |
| 10,14 | Needham, Lancaster | 8, 11 | J.Marshall, S.Ca | rroll# |
| White-throated | Sparrow: | C26 2701 | | |
| 11 | W.Newbury | pr. | W.Petersen# | |
| Bobolink: | 30.40.30.30.45.00 . | 7/2000 | | |
| 4,21 | Sherborn, Rowley | 3, 35 | BBC | |
| Eastern Meadowl | lark: | | | |
| 4,9 | Sherborn, P.I. | 4, 2 nesting | BBC | |
| 13,20 | Lincoln, Halifax | 2 nesting, 6 | J. Carter, K. Ande | rson |
| Orchard Oriole: | | | | |
| 4,8 | E.Orleans, Scituate | 2, 1 m. | A.Williams, W.Pe | tersen |
| Purple Finch: | | 45 50 MM | | |
| 28 | P.I. | 10 | BBC | |
| | | | | |



CHRISTMAS BIRD COUNT

A Christmas Bird Count is a popular census of birdlife taken each year by a number of birdwatchers between December 10 and January 2. This traditional activity, sponsored by the National Audubon Society and the U.S. Fish and Wildlife Service, dates back to 1899 and has grown from a limited affair involving only a few people to an event in which tens of thousands of birders in Canada, United States, Mexico, Central and South America participate. In the 79th Christmas Bird Count (CBC) of 1978, 31,140 people joined in 1269 separate counts to report a total of 103,403,790 individual birds of 600 species. In the CBC of 1982, the number of species reported in any one count ranged from 3 to 306 with the highest count in the U.S. being 222.

Although the participants vary in expertise from beginner to seasoned expert, the magnitude of the undertaking provides data of some scientific significance. For this reason, very specific rules are set up to guide those who take part. Each CBC is a day-long census of all birds within a circular area 15 miles in diameter, and compilers try to "field" as many competent birders as they can to cover the roughly 177 square miles of a count circle, pairing up the less experienced with the better qualified so that all sightings can be confirmed. The competition is keen to produce the highest number of species, and this undoubtedly insures the thoroughness with which the count areas are covered. And with the great number of birders in the field, there are bound to be a number of "rare bird" sightings - something that enables Massachusetts birders to participate enthusiastically despite the rigors of a cold winter's day. The day's activity usually ends with a celebratory suppertime gathering during which the final tally or "countdown" is made.

This year's count period is from Saturday, December 15, 1984 through January 1, 1985. <u>Bird Observer</u> encourages anyone wishing more information or desiring to participate, whatever your level of birding accomplishment, to send a stamped and self-addressed envelope to Robert H. Stymeist, 98 Boylston Street, Watertown, MA 02172.

