

was at Clearwater (Pinellas Co.) in February 1880 (Howell 1932), but recently the species has been reported on Christmas Bird Counts north to Sanford (Seminole Co.) and Titusville (Brevard Co.). Just outside Florida there are sight records of wintering Northern Waterthrushes at Savannah, Georgia, 26 December 1931 (Burleigh 1958: 550) and Fort Morgan, Alabama, 12 February 1973 (Imhof 1976: 360).

All previously collected specimens cited above (Florida, South Carolina, Texas, Pennsylvania) have been referred to *S. n. notabilis*, the mid-continental race with a more grayish back and whiter underparts than the nominate race. Our specimen matches those of *S. n. notabilis* in the Tall Timbers collection, but Mengel (1965: 423) casts doubt on the validity of these two races.

Manuel Pescador, of Florida A. & M. University, and William Heard, of Florida State University, kindly examined the stomach contents of the bird I collected, and Robert L. Crawford prepared the study skin.

LITERATURE CITED

- AMERICAN ORNITHOLOGISTS' UNION. 1957. Check-list of North American birds. Baltimore, Lord Baltimore Press, Inc.
- BURLEIGH, T. D. 1958. Georgia birds. Norman, Univ. Oklahoma Press.
- HOWELL, A. H. 1932. Florida bird life. Tallahassee, Florida Dept. Game and Fresh Water Fish.
- IMHOF, T. A. 1976. Alabama birds (2nd ed.). Univ., Ala., Univ. Ala. Press.
- MENDEL, R. M. 1965. The birds of Kentucky. Ornithol. Monog. No. 3, Amer. Ornitho. Union.
- STEVENSON, H. M. 1949. Seasonal notes in the Tallahassee region. Fla. Nat. 22: 62-63.
- HENRY M. STEVENSON, Tall Timbers Research Station, Rt. 1, Box 160, Tallahassee, Florida 32312.

A Cardinal at the Dry Tortugas, Florida.—Cardinals (*Cardinalis cardinalis*) are widespread in North America (Gruson 1976, Checklist of the World's birds, London, William Collins Sons and Co., p. 155) and are considered nonmigratory (Bent 1968, U.S. Natl. Mus. Bull. 237, p. 15). Howell (1932, Florida bird life, Tallahassee, Fla. Dept. Game Fresh Water Fish, p. 437) gives its southern range as extending through the Florida Keys to the Marquesas Keys but indicates that the species is rare in Key West and states, "A Cardinal struck Sombrero Key Light, September 10, 1884—an unusual occurrence for a nonmigratory bird (Merriam 1885, p. 61)". Sprunt (1954, Florida bird life, New York, Coward-McCann Inc. p. 450) states, "It has since been recorded even from Tortugas: Gabrielson saw one there, March 29, 1948." This is the only published record of a Cardinal at the Dry Tortugas (W. B. Robertson, Jr., pers. comm.).

On 6 April 1978 at 0700, in the parade ground of Fort Jefferson, Garden Key, Dry Tortugas, Florida, I observed a female Cardinal working through the trees. The bird was also observed on the 6th and 8th by members of the tern-banding party (W. B. Robertson, Jr., et al.). The bird was not reported again and was not seen in June by the tern-banding party. I thank William B. Robertson, Jr., for his comments on this note.—THEODORE H. BELOW, 3697 North Road, Naples, Florida 33942.

Pine Siskins in the Florida Keys.—The Pine Siskin (*Carduelis pinus*) winters irregularly to north Florida, and rarely to south Florida (Sprunt 1954, Florida bird life, New York, Coward-McCann, Inc.). The last irruption of siskins reaching peninsular Florida occurred during the winter of 1969-70. In that season Stevenson (1970, Amer. Birds 24:497) noted large numbers seen near St. Petersburg (Pinellas Co.) and West Palm Beach (Palm Beach Co.), and listed several other reports from peninsular Florida. Southernmost reports were two sightings from Everglades Na-