

FLORIDA BIRDS IN THE PERIODICAL LITERATURE

FRED E. LOHRER

Archbold Biological Station, Route 2, Box 180, Lake Placid, Florida 33852

- AKEY, B. L., J. K. NAYAR, AND D. J. FORRESTER. 1981. Avian pox in Florida Wild Turkeys: *Culex nigripalpus* and *Wyeomiu vanduzeei* as experimental vectors. *J. Wild. Dis.* 17: 597-599.
- ALDRICH, J. W. 1981. Geographic variation in White-winged Doves with reference to possible sources of new Florida population. *Proc. Biol. Soc. Wash.* 94: 641-651.—With so much overlap and lack of agreement in size and color characters between White-wing populations in the Gulf of Mexico-West Indies region, all these populations are considered as belonging to one subspecific taxon *Zenaida a. asiatica*. Southern Florida specimens most closely resemble samples from Hispaniola and Cuba and the author suggests that the Florida population may have originated in either of these two islands and reached Florida in avicultural trade through Tampico. Natural range expansion from Cuba is also possible.
- ATKINSON, R. D., T. S. BASKETT, AND K. C. SADLER. 1982. Population dynamics of Mourning Doves banded in Missouri. *U.S. Fish Wild. Serv., Spec. Sci. Rept.-Wild.* No. 250: 1-20.—Includes Florida recoveries.
- BARBOUR, D. B., AND A. R. DEGANGE. 1982. Communal roosting in wintering Hooded Mergansers. *J. Field Ornithol.* 53: 279-280.—Pasco Co.
- BARBOUR, D. B., AND A. R. DEGANGE. 1982. Reciprocal allopreening in the Brown-headed Nuthatch. *Auk* 99: 171.—Marion Co.
- BASHAM, B. 1982. Plantation Report: the 1982 ABA Convention. *Birding* 14: 129-135.—Report of the 10-17 May 1982 American Birding Association annual meeting in southern Florida.
- BESKE, A. E. 1982. Local migratory movements of radio-tagged juvenile harriers. *Raptor Res.* 16: 39-53.—A nestling Northern Harrier banded in Buena Vista Marsh, Central Wisconsin, was recorded in February of its first winter in southeastern Florida (probably Palm Beach Co.).
- BLACK, B. B., AND L. D. HARRIS. 1981. Winter foraging patterns of Gulf coast Black Skimmers. *Colonial Waterbirds* 4: 187-193.—Near Cedar Key, 1979-1980.
- BLACK, B. B., AND M. W. COLLOPY. 1982. Nocturnal activity of Great Blue Herons in a north Florida salt marsh. *J. Field Ornithol.* 53: 403-406.—At Cedar Key tidal flats.
- BUHRMAN, C. B. 1982. The Florida that birders need. *Birding* 14: 137-139.—This brief article lists the major bird species that Florida visitors seek; the pitfalls and pests of Florida birding; and 5 current research projects that birders can contribute sightings to.
- CAPRIO, A., AND D. L. TAYLOR. 1982. Breeding bird census. No. 87. Cutover subtropical slash pine forest. *Amer. Birds* 36: 74.—Everglades National Park, Dade Co.
- CAPRIO, A., D. L. TAYLOR, M. BRITTEN, AND R. ROCHEFORT. 1982. Breeding bird census. No. 88. Virgin subtropical slash pine forest—February burn. *Amer. Birds* 36: 74.—Big Cypress National Preserve, (BCNP) Collier Co.

- CAPRIO, A., D. L. TAYLOR, M. BRITTEN, AND R. ROCHEFORT. 1982. Breeding bird census. No. 89. Virgin subtropical slash pine forest—December burn. *Amer. Birds* 36: 74-75.—BCNP, Collier Co.
- CAPRIO, A., D. L. TAYLOR, M. BRITTEN, AND R. ROCHEFORT. 1982. Breeding bird census. No. 90. Virgin subtropical slash pine forest—Summer burn. *Amer. Birds* 36: 75.—BCNP, Collier Co.
- CAPRIO, A., D. L. TAYLOR, M. BRITTEN, AND R. ROCHEFORT. 1982. Breeding Bird census. No. 91. Virgin subtropical slash pine forest—unburned control. *Amer. Birds* 36: 75.—BCNP, Collier Co.
- CLAPP, R. B., M. K. KLIMKIEWICZ, AND J. H. KENNARD. 1982. Longevity records of North American birds: Gaviidae through Alcidae. *J. Field Ornithol.* 53: 81-124.—A Laysan Albatross holds the record for the oldest N. A. bird, 37+ years (or 39-02), but a Dry Tortugas Sooty Tern is close at 32 years. Other Florida records are Brown Pelican, 19-08 (years, months); Double-crested Cormorant, 17-08; White Ibis, 16-04; Black Skimmer, 20-00.
- CLAPP, R. B., R. C. BANKS, D. MORGAN-JONES, AND W. A. HOFFMAN. 1982. Marine birds of the southeastern United States and Gulf of Mexico. Part I. Gaviiformes through Pelecaniformes. U.S. Fish Wild. Serv., Off. Biol. Serv. FWS/OBS-82/01, 637 pp.
- CLAPP, R. B., D. MORGAN-JONES, AND R. C. BANKS. 1982. Marine birds of the southeastern United States and Gulf of Mexico. Part II. Anseriformes. U.S. Fish Wild. Serv., Off. Biol. Serv. FWS/OBS-82/20, 491 pp.
- CLARK, F. S. 1982. Florida, Kissimmee, Sandhill Crane, Limpkin. *Birding* 14: 128(u-v).—A birding-site, tear-out sheet to a site on Reedy Creek and on Lake Tohopekaliga, both fairly close to Orlando.
- CONTI, J. A., AND D. J. FORRESTER. 1981. Interrelationships of parasites of White-winged Doves and Mourning Doves in Florida. *J. Wild. Mgmt.* 17: 529-536.—Mourning Doves from Alachua and Dade cos. compared with White-wings from Dade Co.
- DOLBEER, R. A. 1982. Migration patterns for age and sex classes of blackbirds and starlings. *J. Field Ornithol.* 53: 28-46.—One or more Red-winged Blackbirds banded in Detroit area was recovered in northern Florida.
- DUNCAN, C. D. 1982. Western Gull in Alabama and northwestern Florida. *Amer. Birds* 36: 899-902.—A one-legged gull first sighted 24 October 1977 at Mobile Bay and identified as a Lesser Black-backed Gull has been subsequently seen in 1978 and in 1981 in Fort Pickens, Florida, and reidentified as a Western Gull.
- DUNKS, J. H., R. E. TOMLINSON, H. M. REEVES, D. D. DOLTON, C. E. BRAUN, AND T. P. ZAPATKA. 1982. Migration, harvest, and population dynamics of Mourning Doves banded in the Central Management Unit, 1967-77. U.S. Fish Wild. Serv., Spec. Sci. Rept. Wild. No 249: 1-128.—The CMU includes 14 states from the Rockies to the Mississippi River. Several tables include state of origin for birds recovered in Florida.
- ELOWSON-HALEY, A. M. 1982. Osprey spreads wings after fishing. *Wilson Bull.* 94: 366-368.—As observed in Everglades National Park, Dade Co.
- ENGSTROM, T. 1982. Breeding bird census. No. 31. Mature beech-magnolia forest. *Amer. Birds* 36: 61.—Tall Timbers Res. Sta., Leon Co.
- ERWIN, R. M. 1981. Censusing wading bird colonies: An update on the "flight-line" count method. *Colonial Waterbirds* 4: 91-95.—Based primarily on ten colonies of Cattle Egrets, Louisiana Herons, Snowy Egrets, and/or Little Blue Herons studied in south-central Florida during Spring 1980.

- EVANS, D. L. 1982. Status reports on twelve raptors. U.S. Fish Wild. Serv., Spec. Sci. Rept.-Wild. 238: 1-68.—Distribution, ecology, management and status; compiled largely from the Literature through mid-1977 with some update through 1979. Nine of the 12 species occur regularly in Florida; Bald Eagle, Burrowing Owl, Caracara, Cooper's Hawk, Marsh Hawk, Merlin, Osprey, Peregrine Falcon, Sharp-shinned Hawk. However, only the Caracara and the Burrowing Owl accounts contain much information about Florida populations.
- FICKETT, S. B. JR., AND S. NESBITT. 1982. Bald Eagle surveys in Florida. ENFO 82-3: 4-5.
- FISHER, R. P. 1982. An albinistic Brown Pelican. Fla. Nat. 55(5): 7-10.—Includes 2 cover photos. At New Smyrna Beach. Same bird as in *FFN* 7: 6.
- FITZPATRICK, J. W., AND G. E. WOOLFENDEN. 1981. Demography is a cornerstone of Sociobiology. *Auk* 98: 406-407.—A commentary based on Scrub Jay studies conducted at Archbold Biol. Sta., Highlands Co.
- FORRESTER, D. J. 1980. Haematozoa and mallophaga from the White Ibis, *Eudocimus albus* L. in Florida. *J. Parasitol.* 66: 58.
- FORRESTER, D. J., AND P. P. HUMPHREY. 1981. Susceptibility of the knot (*Calidris canutus*) to *Plasmodium hermani*. *J. Parasitol.* 67: 747-748.—Study based on knots from the St. Petersburg area.
- FORRESTER, D. J., P. R. HUMPHREY, S. R. TELFORD, JR., AND L. E. WILLIAMS, JR. 1980. Effects of blood-induced infections of *Plasmodium hermani* on domestic and wild Turkey poults. *J. Wild. Dis.* 16: 237-244.—Wild Turkeys from Glades Co.
- FRITTS, T. H., AND R. P. REYNOLDS. 1981. Pilot studies of the marine mammals, birds, and turtles in the OCS areas of the Gulf of Mexico. U.S. Fish. Wild. Serv., Office Biol. Ser. FWS/OBS-81/36, 139 pp.—Results of aerial surveys in 2 areas off Florida during August and November 1979.
- GEE, G. F., J. W. CARPENTER, AND G. L. HENSLER. 1981. Species differences in hematological values of captive cranes, geese, raptors, and quail. *J. Wild. Mgmt.* 45: 463-483.—Includes values for Florida Sandhill Cranes.
- GOOD, J. M. 1982. Winter bird population study. No. 9. Slash pine. *Amer. Birds* 36: 31. Everglades Natl. Park, Dade Co.
- GOOD, J. M. 1982. Winter bird population study. No. 51. Canal bank and associated spoil bank. *Amer. Birds* 36: 41.—2 km N of Homestead, Dade Co.
- GOOD, J. M. 1982. Winter bird population study. No. 70. Avocado grove. *Amer. Birds* 36: 45. 1.6 km N of Homestead, Dade Co.
- GREINER, E. C., D. J. BLACK, AND W. O. IVERSON. 1981. *Plasmodium* in a Bald Eagle (*Haliaeetus leucocephalus*) in Florida. *J. Wild. Dis.* 17: 555-558.—Adult eagle found injured along north-central Florida road.
- HAILMAN, J. P., AND J. R. REED. 1982. Head wind promotes skimming in Laughing Gulls. *Wilson Bull.* 94: 223-225.—At Flamingo, Everglades National Park, Monroe Co.
- HARDY, J. W., AND M. H. CLENCH. 1982. First United States specimen of the White-collared Swift. *Amer. Birds* 36: 139-141.—Found dead, clinging to a screen door of a condominium on Perdido Key, 15 km W of Pensacola, Escambia Co., on 25 January 1981. Specimen to FSM.
- HARRINGTON, B. A., AND L. E. LEDDY. 1982. Sightings of knots banded and color-marked in Massachusetts in August 1980. *J. Field Ornithol.* 53: 55-57.—During January 1981 the authors looked for marked Red Knots among 4 500 in the Sarasota area but found no color-marked individuals.

- HARRINGTON, B. A., D. C. TWICHELL, AND L. E. LEDDY. 1982. Knots in Florida—A migration mystery. *Fla. Nat.* 55(5): 4-5.—Portions of this paper delivered at 1981 fall FOS meeting.
- HARRINGTON, B. A. 1982. Untying the enigma of the Red Knot. *The Living Bird Quarterly* 1(2): 4-7.—Includes mention of Florida's Gulf coast knots.
- HENNEMANN, III, WILLARD W. Energetics and spread-winged behavior of Anhingas in Florida. *Condor* 84: 91-96.—Studies of four hand-reared captives conducted at Gainesville and presumably collected somewhere in Florida.
- HOLLER, N. R., AND E. W. SCHAFER, JR. 1982. Potential secondary hazards of avitrol baits to Sharp-shinned Hawks and American Kestrels. *J. Wild. Mgmt.* 46: 457-462.—Sharp-shins collected near Gainesville, Alachua Co., Nov. and Dec. 1979.
- KALE, H. W., II, AND R. W. LOFTIN. 1982. A longevity record for the Black Skimmer. *N. A. Bird Bander* 7: 54.—Banded in Georgia by Kale, recovered near Mayport by Loftin, age 20+ years. See Clapp et al. above.
- KENNER, K. C., AND W. R. MARION. 1981. Wood Duck production on a northern Florida phosphate mine. *J. Wild. Mgmt.* 45: 1037-1042.—Hamilton Co.
- KILHAM, L. 1982. Florida Red-shouldered Hawk robs American Crows. *Wilson Bull.* 94: 566-567.—Highlands Co.
- KILHAM, L. 1982. Cleaning/feeding symbioses of Common Crows with cattle and feral hogs. *J. Field Ornithol.* 53: 275-276.—Highlands Co.
- KLASS, E. E., H. M. OHLENDORF, AND E. CROMARTIE. 1980. Organochlorine residues and shell thickness in eggs of the Clapper Rail, Common Gallinule, Purple Gallinule, and Limpkin (Class Aves), eastern and southern United States, 1972-1974. *Pesticides Monitoring J.* 14: 90-94.—Gallinule eggs (*Porphyryla*, n=10; *Gallinula*, n=15) collected from Brevard, Glades, Lee, and Marion co., and Limpkin eggs (n=14) collected from Lake, Okeechobee, and Glades co. No evidence of eggshell thinning was found for any species studied, but the gallinule eggs contained residues of DDE and PCBs.
- KROENKE, S. R. 1982. Eastern Bluebirds in Jefferson County, Florida. *Sialia* 4: 15-16.—Nesting data for 10 boxes in 1981.
- KUSHLAN, J. A., AND L. C. MCEWAN. 1982. Nesting phenology of the Double-crested Cormorant. *Wilson Bull.* 94: 201-206.—In southern Florida.
- KUSHLAN, J. A., O. L. BASS, JR., AND L. C. MCEWAN. 1982. Wintering waterfowl in the Everglades estuaries. *Amer. Birds* 36: 815-819.
- LARSON, S. E. 1982. Winter population trends in the Cattle Egret. *Amer. Birds* 36: 354-357.—A considerable portion of the discussion considers Florida.
- LAYNE, J. N. 1982. Status of sibling aggression in Florida Sandhill Cranes. *J. Field Ornithol.* 53: 272-274.—In 10 south-central Florida counties.
- LAYNE, J. N. 1982. Analysis of Florida-related banding data for the American Kestrel. *N. A. Bird Bander* 7: 94-99.
- LAYNE, J. N. 1982. The caracara in Florida. *ENFO* 82-3: 10-12.
- LEBERMAN, R. C. 1982. Powdermill Nature Reserve-1980 recoveries. *N. A. Bird Bander* 7: 53.—Two birds banded at Powdermill NR (of Carnegie Museum of Natural History) in western Pennsylvania were recovered in Florida: An American Robin found dead 17 January 1980, St. Petersburg, age 2 yrs., 7 mo; A Yellow-rumped Warbler found dead 15 January 1980, St. Cloud, age less than 1 year.
- LEENHOUTS, W. P. 1982. Breeding bird census. No. 186. Cordgrass salt marsh I. *Amer. Birds* 36: 99.—St. Johns River valley, Brevard Co.

- LEENHOUTS, W. P. 1982. Breeding bird census. No. 187. Cordgrass salt marsh II Amer. Birds 36: 99-100.—St. Johns River valley, Brevard Co.
- LEENHOUTS, W. P., AND J. L. BAKER. 1982. Vegetational dynamics in Dusky Seaside Sparrow habitat on Merritt Island National Wildlife Refuge. Wild. Soc. Bull. 10: 127-132.—Brevard Co.
- LOFLIN, R. K. 1982. Ani male apparently killed by other anis while attempting to parasitize nest. Auk 99: 787-788.—At Virginia Key, Dade Co.
- LOVVERN, J. R., AND C. M. KIRKPATRICK. 1982. Fall census of Eastern Greater Sandhill Cranes. Pp. 259-265 in Lewis, J. C. (ed.), Proc. 1981 Crane Workshop, Natl. Aud. Soc., Tavernier, FL.—The results of the 1979 and 1980 census include some Florida data.
- MAEHR, D. S. 1982. The adaptable Least Tern. Fla. Nat. 53(3): 7-10.—Popular account of the discovery and history of Least Terns nesting at a phosphate mine in Madison Co.
- MAGER, D. 1982. Notes on eagle nesting in Florida. ENFO 82-3: 6-7.
- MARION, W. R., AND T. E. O'MEARA. 1982. Limpkins. P 75 in Davis, D. E. (ed.), CRC Handbook of census methods for terrestrial vertebrates. CRC Press, Boca Raton, FL.—Techniques perfected in Florida.
- MARION, W. R., T. E. O'MEARA, AND L. D. HARRIS. 1981. Characteristics of the Mourning Dove harvest in Florida. J. Wild. Mgmt. 45: 1062-1066.—Describes the source (29 states East of SD and Ontario) and movements of Mourning Doves shot in Florida during 1931-1978 based on 1625 band recoveries. 43 percent of recoveries were birds banded in Florida. Of the remainder, the following 5 states, in descending order, account for half of the recoveries: Illinois; Missouri; Wisconsin; Georgia; Alabama.
- MARION, W. R., T. E. O'MEARA, AND D. S. MAEHR. 1981. Use of playback recordings in sampling elusive or secretive birds. Stud. Avian Biol. 6: 81-85.—King and Virginia rails and Soras in phosphate mines in Hamilton Co.; Limpkins on Lake Ocklawaha, Marion and Putnam co.; Plain Chachalacas in Texas.
- MCDONALD, M. V. 1982. Breeding bird census. No. 188. Gulf coast salt marsh. Amer. Birds 36: 100.—6 km NW of Cedar Key, Levy Co.
- MONTALBANO, F., III. 1982. Summer use of two central Florida phosphate settling ponds by Florida Ducks. Proc. Ann. Conf. S.E. Assoc. Fish Wild. Agencies 34: 584-590.—Polk Co.
- MURPHY, W. L. 1982. The Ash-throated Flycatcher in the East: an overview. Amer. Birds 36: 241-247.—Florida information includes an account of a Virginia Key, Dade Co., sighting (20 January 1980), and a listing of the three previous Florida records.
- NESBITT, S. A., P. E. COWAN, P. W. RANKIN, N. P. THOMPSON, AND L. E. WILLIAMS, JR. 1981. Chlorinated hydrocarbon residues in Florida Brown Pelicans. Colonial Waterbirds 4: 77-84.—Study based on 56 birds obtained in Florida, 1971-1972.
- NESBITT, S. A. 1982. The past, present, and future of the Whooping Crane in Florida. Pp. 151-154 in Lewis, J. C. (ed.), Proc. 1981 Crane Workshop, Natl. Aud. Soc., Tavernier, FL.
- NESBITT, S. A., B. A. HARRIS, R. W. REPENNING, AND C. B. BROWNSMITH. 1982. Notes on Red-cockaded Woodpecker study techniques. Wild. Soc. Bull. 10: 160-163.—Techniques developed on Florida birds.
- ODGEN, J. C., H. W. KALE, II, AND S. A. NESBITT. 1980. The influence of annual variation in rainfall and water levels on nesting by Florida populations of

- wading birds. Trans. Linnaean Soc. N.Y. 9: 115-126.—Based on state-wide aerial surveys conducted during April-May and June-July, 1976, 1977, 1978.
- O'MEARA, T. E. 1981. A field test of two density estimators for transect data. Stud. Avian Biol. 6: 193-196.—Breeding-bird censuses conducted in slash pine plantations in Alachua and Bradford co.
- POOLE, A. 1981. The effects of human disturbance on Osprey reproductive success. Colonial Waterbirds 4: 20-27.—In Florida Bay portions of Everglades National Park and in NE U.S.
- POOLE, A. 1982. Brood reduction in temperate and sub-tropical Ospreys. Oecologia 53: 111-119.—Florida Bay and Long Island, N.Y.
- POST, W., AND F. B. ANTONIO. 1981. Breeding and rearing of Seaside Sparrows *Ammodramus maritimus* in captivity. Int. Zoo. Yearbook 21: 123-128.—6 hand-reared nestlings and 19 wild-caught adult *A. m. peninsulae* from Cedar Keys and 3 adult male *A. m. nigrescens* were bred.
- RECHER, H. F., AND J. A. RECHER. 1980. Why are there different kinds of herons? Trans. Linnaean Soc. N. Y. 9: 135-158.—Based in part on observations of feeding herons in Everglades National Park and the Florida Keys.
- RICKERT, J. E. 1982. Florida Ornithological Society. Birding 14: 128(s).—Tear-out sheet. Revised club supplement for the book, A guide to North American bird clubs.
- ROBERTSON, W. B., JR. 1982. Raptors on the National Park Service areas of southern Florida. ENFO 82-3: 7-10.
- RUSCH, D. H. 1982. Great Horned Owl. Pp. 87-88 in Davis, D. E. (ed.), CRC Handbook of census methods for terrestrial vertebrates, CRC Press, Boca Raton.—Includes a table of mean hatching date vs latitude, with Florida the earliest at the third week of January.
- RYDER, R. A. 1981. Movements and mortality of White Pelicans fledged in Colorado. Colonial Waterbirds 4: 72-76.—Of 67 recoveries from elsewhere than Colorado, 2 were from Florida. There were 37 recoveries from coastal Texas, Mexico, and Central America. Where do Florida's White Pelicans originate?
- SCHNEIDER, D. 1982. Predation by Ruddy Turnstones (*Arenaria interpres*) on a polymorphic clam (*Donax variabilis*) at Sanibel Island, Florida. Bull. Marine Sci. 341-344.—In March 1980, the turnstones were capturing clams in the swash zone, using vision rather than probing, carrying them beyond the reach of the waves, and pecking them open. The coloration of the predated clams did not differ significantly from the coloration of clams sampled from the beach.
- SCOTT, M. D., AND J. A. POWELL. 1982. Commensal feeding of Little Blue Herons with manatees. Wilson Bull. 94: 215-216.—St. Johns River, Volusia Co.
- SERAFIN, J. A. 1982. The influence of diet composition upon growth and development of Sandhill Cranes. Condor 84: 427-434.—Results of experiments conducted on captive *Grus canadensis pratensis* and *G. c. tabida* at Patuxent Wildlife Research Center indicate that the non-migratory Florida Sandhills grow slower than migratory Greater Sandhills regardless of type of ration and did not develop the leg or wing abnormalities that Greater were prone to when raised on a high protein diet.
- SHAPIRO, A. E., F. MONTALBANO, III, AND D. MAGER. 1982. Implications of construction of a flood control project upon a Bald Eagle nesting activity. Wilson Bull. 94: 55-63.—Kissimmee Basin in Osceola, Polk, Okeechobee,

Highlands and Glades co.

- SYKES, P. W., JR. 1982. North Carolina, South Carolina, Georgia, Florida. Amer. Birds 36: 384-387.—Summary of 82nd Audubon CBC; includes summaries of diurnal raptor counts for the region and parulids reported for 1979-80, 1980-81, and 1982-82 CBCs in Florida.
- SYKES, P. W., JR. 1982. Everglade Kite. Pp. 43-44 in Davis, D. E. (ed.), CRC Handbook of census methods for terrestrial vertebrates. CRC Press, Boca Raton.—Two census methods used are airboat transects and evening counts at roosts.
- SMALLWOOD, J. A., WOODREY, N. J. SMALLWOOD, AND M. A. KETTLER. 1982. Foraging by Cattle Egrets and American Kestrels at a fire's edge. J. Field Ornithol. 53: 171-172.—At Clewiston and Immokalee.
- STEVENSON, H. M. 1982. Bachman's Warbler . . . rarest North American bird? Fla. Wildlife 35(6): 36-39.—A history of our knowledge about this enigmatic bird.
- STEWART, P. A. 1982. Migration of Blue Jays in eastern North America. N. A. Bird Bander 7: 107-112.—Includes Florida data.
- THRELFALL, W. 1982. Endoparasites of the Double-crested Cormorant (*Phalacrocorax auritus*) in Florida. Proc. Helminthol. Soc. Wash. 49: 103-108.
- THRELFALL, W. 1982. Ectoparasites (Mallophaga, Acarina) from the Double-crested Cormorant (*Phalacrocorax auritus*) in Florida. Proc. Entomol. Soc. Wash. 84: 369-375.—Based on examination of 77 cormorants from several locations in Florida (1973-1977).
- WALKINSHAW, L. H. 1981. The Sandhill Cranes. Pp. 151-162 in Lewis, J. C., and H. Masatomi (eds.), Crane research around the World, Baraboo, WI, International Crane Foundation.—Includes the Florida Sandhill; habitat, breeding biology, growth of chicks, and population status.
- WALKINSHAW, L. H. 1982. Nesting of the Florida Sandhill Crane in central Florida. Pp. 53-62 in Lewis, J. C. (ed.), Proc. 1981 Crane Workshop, Natl. Aud. Soc., Tavernier, FL.
- WEBBER, L. A. 1981. A census of birds and fish in areas treated with mosquito adulticides versus similar untreated areas. J. Fla. Anti-Mosquito Assoc. 52: 51-55.—As studied in Collier Co. from December 1979 to January 1981.
- WHITE, F. H., D. J. FORRESTER, AND L. E. WILLIAMS, JR. 1981. Isolation of *Salmonella* from Wild Turkeys in Florida. J. Wildl. Mgmt. 17: 327-330.—411 birds were collected from 17 sites in Florida.
- WILSON, J., AND R. W. LOFTIN. 1982. Winter bird population study. No. 50. Barrier beach and saltwater estuary. Amer. Birds. 35: 41.—N. side of St. Johns River inlet, Duval Co.
- WINGATE, D. B. 1982. Successful reintroduction of the Yellow-crowned Night-Heron as a nesting resident on Bermuda. Colonial Waterbirds 5: 104-115.—Nestlings were obtained from Alafia Banks heronry, Tampa Bay, 1976-1978.