

EVIDENCE OF KRILL IN THE DIET OF BALEARIC SHEARWATERS *PUFFINUS MAURETANICUS*

MAITE LOUZAO^{1,2}, DAVID GARCÍA^{3,4}, BENEHARO RODRÍGUEZ^{3,5} & PERE ABELLÓ⁶

¹Centro Oceanográfico de Xixón, Instituto Español de Oceanografía (IEO),
Camín de l'Arbeyal s/n, E-33212 Gijón/Xixón, Spain (maite.louzao@gmail.com)

²AZTI, Calle Herrera Kaia, Portualdea z/g, 20110 Pasaia, Spain

³SEO/BirdLife, Carrer Múrcia 2-8, local 13 interior 08026 Barcelona, Spain

⁴Iniciativa de Recerca de la Biodiversitat de les Illes (IRBI), C/ Son Borràs, 14, E-07340 Alaró, Balearic Islands, Spain

⁵Grupo de Ornitología e Historia Natural de las Islas Canarias (GOHNIC),
La Malecita s/n, E-38480 Buenavista del Norte, Tenerife, Spain

⁶Institut de Ciències del Mar (CSIC), Passeig Marítim de la Barceloneta, 37-49, E-08003 Barcelona, Spain

Received 11 August 2014, accepted 20 November 2014

Seabird diets are generally composed of a few major taxa such as pelagic fish, squid and crustaceans (Shealer 2002), with small pelagic fish being the main foraging resource in temperate regions (Furness & Monaghan 1987). This is also true for the critically endangered Balearic Shearwater *Puffinus mauretanicus*. Its diet includes small pelagic fish and demersal fish obtained from trawling discards (Gutiérrez & Figuerola 1995, Arcos & Oro 2002, Louzao *et al.* 2006b, Käkälä *et al.* 2010). Direct observations at sea determined that this species also feeds on plankton, although the species of plankton have not been identified (Arcos *et al.* 2000, Arcos & Oro 2002). In this study, we report observations that, during breeding season, the diet of the endemic Balearic Shearwater includes macrozooplankton, specifically the krill *Nyctiphanes couchii*.

METHODS

We monitored the breeding population of Balearic Shearwaters located in the south of Sa Conillera Island (southwest of Ibiza Island in the Mediterranean Sea) during the chick-rearing period (May) of 2013. Sa Conillera Island is part of the protected area *Reserves Naturals des Vedrà, es Vedranell i els Illots de Ponent*, where approximately 400 breeding pairs of Balearic Shearwater nest (Arcos 2011). The chick-rearing period starts in late April and can last until early July (Ruiz & Martí 2004, Louzao *et al.* 2006b). During this period, adults visit the colony to feed their young during the night. In May 2013, we collected a Balearic Shearwater

regurgitation, which contained crustaceans only (Fig. 1), and we preserved the sample in 70% ethanol. Body length measurements of the prey were taken using a Nikon AZ100 microscope (10× magnification) fitted with a Nikon DS-Fi1 camera and NIS-Elements software.

RESULTS AND DISCUSSION

The entire regurgitation was composed of the euphausiid *Nyctiphanes couchii*. A subsample of this material was deposited in the Biological Reference Collections (CBR) of the Institut de Ciències del Mar (CSIC), Barcelona (reference code ICMR000001).

We measured both the total and cephalothorax length of 100 individuals in the sample (Fig. 2). The mean total length was 14.36 mm ± 1.73 mm (SD), ranging from 11.26 mm to 18.27 mm, and the mean cephalothorax length was 4.09 mm ± 0.55 mm (SD), ranging from 3.27 mm to 5.76 mm. Both measurements were positively related ($y = 0.268 + 0.265 * x$, $F_{1,83} = 215.6$, $P > 0.001$, R^2 -adjusted = 0.718). Based on the characteristics of the related *N. australis* (Ross & Quetin 2000), all individuals were adults.

This is the first evidence that Balearic Shearwaters feed on krill during the breeding season (chick-rearing), a period when the species forages in productive waters of the continental shelf of the western Mediterranean (Louzao *et al.* 2006a, 2012; Arcos *et al.* 2012). According to previous studies, the diet of the chicks consisted mainly of anchovies, a high-energy source (Navarro *et al.* 2009).

In the Mediterranean, euphausiids are most abundant in waters of the continental slope deeper than 200 m, and at night they perform diel vertical migrations that reach the sea surface (Wiebe & D'Abramo 1972). During this period, this species could be accessible to Balearic Shearwaters, which are able to dive to 26 m depth (Aguilar *et al.* 2003). Among krill species, *Meganyctiphanes norvegica* is the most abundant species found in the deeper waters over Mediterranean continental slopes, while *N. couchii* is generally restricted to the continental shelf (Lindley 1982). The habitat of Balearic Shearwaters matches the distribution of *N. couchii* (Louzao *et al.* 2006a, 2012), and the birds have been observed feeding on unidentified zooplankton (Arcos *et al.* 2000, Arcos & Oro 2002).



Fig. 1. Regurgitation from a Balearic Shearwater obtained in May 2013, composed solely of *Nyctiphanes couchii* (Photo by M. Louzao).

Further research is needed to accurately assess the importance of euphausiids in the diet of the Balearic Shearwater. However, given that this species is critically endangered, i.e. very few birds remain, it does not lend itself to intensive research. Thus, opportunistic observations, such as the present study, are very important in learning more about the natural history of this species.

ACKNOWLEDGEMENTS

Thanks to Joan Cartes, Félix Pérez and Juan Bueno, who provided comments about the biology, identification and measurements of krill; to Iñaki Huskin, who provided krill images; to José Manuel Arcos and to the Editor, who provided valuable comments on the manuscript. We are especially grateful to *Protecció d'Espècies* and *Espais de Natura Balear* (Balearic Government) for their support in providing permits and logistics. Special thanks also to Virginia Picorelli, Helena Ribas and Jose Vicente Ripoll as well as to Esteban Cardona and his family. Thanks to Karine Delord, Henri Weimerskirch, Amélie Boué, Thierry Micol, Asunción Ruiz, Sandra Benbeniste and Isabel Bermejo for providing logistical support. This study was funded by CONPELHAB, a Marie-Curie Career Integration Grant (PCIG09-GA-2011-293774), the LIFE+ INDEMARES fund (LIFE07NAT/E/000732) and the Ibiza Preservation Fund. M.L. was funded by a *Juan de la Cierva* postdoctoral fellowship (JCI-2010-07639) of the Ministerio de Ciencia e Innovación and a *Ramón y Cajal* postdoctoral fellowship (RYC-2012-09897). This paper is contribution number 698 from AZTI-Marine Research.

REFERENCES

AGUILAR, J.S., BENVENUTI, S., ANTONIA, L.D., MCMINN-GRIVÉ, M. & MAYOL-SERRA, J. 2003. Preliminary results on the foraging ecology of Balearic shearwaters (*Puffinus mauretanicus*)

- from bird-borne data loggers. *Scientia Marina* 67: 129–134.
- ARCOS, J.M. 2011. International species action plan for the Balearic Shearwater, *Puffinus mauretanicus*. Cambridge, UK: SEO/BirdLife & BirdLife International.
- ARCOS, J.M., BÉCARES, J., VILLERO, D., BROTONS, L., RODRÍGUEZ, B. & RUIZ, A. 2012. Assessing the location and stability of foraging hotspots for pelagic seabirds: an approach to identify marine Important Bird Areas (IBAs) in Spain. *Biological Conservation* 156: 30–42.
- ARCOS, J.M., MASSUTÍ, E., ABELLÓ, P., ORO, D. 2000. Brief report: Fish associated with floating drifting objects as a feeding resource for Balearic Shearwaters *Puffinus mauretanicus* during the breeding season. *Ornis Fennica* 77: 177–182.
- ARCOS, J.M. & ORO, D. 2002. Significance of fisheries discards for a threatened Mediterranean seabird, the Balearic shearwater *Puffinus mauretanicus*. *Marine Ecology Progress Series* 239: 209–220.
- FURNESS, R.W. & MONAGHAN, P. 1987. Seabird ecology. Glasgow, UK: Blackie.
- GUTIÉRREZ, R. & FIGUEROLA, J. 1995. Wintering distribution of the Balearic shearwater (*Puffinus yelkouan mauretanicus*, Lowe 1921) off the Northeastern coast of Spain. *Ardeola* 42: 161–166.
- KÄKELÄ, R., KÄKELÄ, A., MARTÍNEZ-ABRAÍN, A., SARZO, B., LOUZAO, M., GERIQUE, C., VILLUENDAS, E., STRANDBERG, U., FURNESS, R.W. & ORO, D. 2010. Fatty acid signature analysis confirms foraging resources of a globally endangered Mediterranean seabird species: calibration test and application to the wild. *Marine Ecology Progress Series* 398: 245–258.
- LINDLEY, J.A. 1982. Population dynamics and production of euphausiids. *Marine Biology* 66: 37–46.
- LOUZAO, M., DELORD, K., GARCÍA, D., BOUÉ, A. & WEIMERSKIRCH, H. 2012. Protecting persistent dynamic oceanographic features: Transboundary conservation efforts are needed for the critically endangered Balearic shearwater. *PLoS*




Fig. 2. Histogram of *Nectiphanes couchii* measurements: total length and cephalothorax length (mm) were taken from the Balearic Shearwater regurgitation in Fig. 1.

- One* 7: e35728.
- LOUZAO, M., HYRENBACH, K.D., ARCOS, J.M., ABELLÓ, P., GIL DE SOLA, L. & ORO, D. 2006a. Oceanographic habitat of an endangered Mediterranean Procellariiform: implications for the design of marine protected areas. *Ecological Applications* 16: 1683–1695.
- LOUZAO, M., IGUAL, J.M., MCMINN, M., AGUILAR, J.S., TRIAY, R. & ORO, D. 2006b. Small pelagic fish, trawling discards and breeding performance of the critically endangered Balearic shearwater: improving conservation diagnosis. *Marine Ecology Progress Series* 318: 247–254.
- NAVARRO, J., LOUZAO, M., IGUAL, J.M., ORO, D., DELGADO, A., ARCOS, J.M., GENOVART, M., HOBSON, K.A. & FORERO, M.G. 2009. Seasonal changes in the diet of a critically endangered seabird and the importance of trawling discards. *Marine Biology* 156: 2571–2578.
- ROSS, R. & QUETIN, L. 2000. Reproduction in euphausiacea. In: Everson, I. (Ed.) *Krill biology, ecology and fisheries*. Oxford, UK: Blackwell Science. pp. 150–181.
- RUIZ, A. & MARTÍ, R. 2004. *La pardela balear*. Madrid, Spain: SEO/BirdLife-Conselleria de Medi Ambient del Govern de les Illes Balears.
- SHEALER, D.A. 2002. Foraging behavior and food of seabirds. In: Schreiber EA, Burger J (Eds.) *Biology of marine birds*. Boca Raton, FL: CRC Press. pp. 137–177.
- WIEBE, P.H. & D'ABRAMO, L. 1972. Distribution of euphausiid assemblages in the Mediterranean Sea. *Marine Biology* 15: 139–149.
-