

THEMIGRANI

A QUARTERLY JOURNAL DEVOTED TO TENNESSES BIRDS

Published by THE TENNESSEE ORNITHOLOGICAL SOCIETY

Joms' Is.

> **SEPTEMBER, 1986** VOL. 57, NO. 3

THE MIGRANT

A QUARTERLY JOURNAL OF ORNITHOLOGY

FIRST PUBLISHED, JUNE 1930

PUBLISHED BY

THE TENNESSEE ORNITHOLOGICAL SOCIETY

Founded at Nashville, Tenn., 7 October 1915
A non-profit, educational, scientific, and conservation organization.

EDITORIAL STAFF

Editor—Charles P. Nicholson, Box 402, Norris, TN 37828 State Count Compiler—Richard L. Knight, 804 North Hills Dr., Johnson City, TN

Editorial Advisors-T. David Pitts, Morris D. Williams

OFFICERS FOR 1985-1987

President—James A. Ferguson, 3090 Joslyn St., Memphis, TN 38128 Vice Presidents:

East Tenn.—Lillian H. Dubke, 8139 Roy Lane, Ooltewah, TN 37363 Middle Tenn.—Thomas Guschke, 113 Taggart Ave., Nashville, TN 37205 West Tenn.—Dollyann Daily, 701 Stevens, Memphis, TN 38116

Directors-at-Large:

East Tenn.—Dee Eiklor, Rt. 11, Box 178, Gray, TN 37615
Middle Tenn.—Ruth McMillan, Rt. 2, Box 141, Lascassas, TN 37085
West Tenn.—Don Manning, Rt. 1, Box 303, McKenzie, TN 38201
Curator—James T. Tanner, Rt. 28, Box 155, Knoxville, TN 37920
Secretary—Carolyn Bullock, 4746 Cole Rd., Memphis, TN 38117
Treasurer—George R. Payne, 6643 Monmouth Dr., Memphis, TN 38119

All TOS members receive *The Migrant* and the *TOS Newsletter*, *The Tennessee Warbler*. The *Newsletter* carries information about meetings, forays, and club activities.

Annual dues, \$6.00; Sustaining, \$10.00; Life, \$150.00; Student (under 18 years), \$3.00; Family, \$7.00; (chapters may collect additional fees to cover local expenses). Dues, contributions, and bequests are deductible from Federal income and estate taxes. Back numbers may be had from Dr. James T. Tanner, Rt. 28, Box 155, Knoxville, Tenn. 37920. Correspond with the Treasurer for subscriptions, memberships, and changes of address.

Published quarterly (March, June, September, and December). Printed by Hubert E. Hodge Printing, Inc., 1800-1802 Grand Avenue, Knoxville, Tennessee 37916, U.S.A. Postage paid and mailed at Norris, Tennessee 37828, U.S.A.

THE MIGRANT

Published by the Tennessee Ornithological Society, to Record and Encourage the Study of Birds in Tennessee. Issued in March, June, September, and December.

VOL. 57

SEPTEMBER, 1986

NO. 3

THE 1986 SPRING FIELD DAYS Richard L. Knight

The 1986 Spring Field Days yielded 220 species from 12 counts across Tennessee. The species total has been exceeded only once (225 species in 1983) and equaled once (1984). The most notable of several unusual species were Anhinga, Greater White-fronted Goose, American Avocet, Baird's Sandpiper, Laughing Gull, Least Tern, and Scissor-tailed Flycatcher. Many rarities were reported with little or no documentation, a trend that needs to be reversed.

INFORMATION ON THE COUNTS

MEMPHIS—27 April; 0730 to 2000; clear; temp. 62-85°F; Ben B. Coffey, Jr. (compiler—672 N. Belvedere, Memphis 38107).

LAWRENCEBURG—3-4 May; Sat. - 0500 to 2000, clear, temp 42-69°F; Sun. - 0530 to 2000, clear, temp. 35-70°F; 12 observers in 7 parties—Andy Augustin, Michael Beuerlein, Helen & Lloyd M. Clayton, Daphnie Griggs, Ann & Lee Mann, Chester & Dot McConnell, Damien J. Simbeck (compiler—Rt. 2, Box 3A, Loretto 38469), Don Simbeck, Morris Williams, Wylie Willis.

CLARKSVILLE—3-4 May; clear; av. temp. 60°F; 10 observers—Joe Allen, George Beringer, Mike Dinsmore, Debbie Gillis, John Habermel, Annie Heilman, Nancy Hughes, Marilyn Lucy, Linda Mattes, John Quick (compiler—Rt. 1, Box 114B, Cunningham 37052).

COLUMBIA—2-3 May; Fri. afternoon & all day Sat.; clear; temp. 40-70°F; 9 observers in 5 parties—Kenneth Anderson, Daniel Gray, William Jernigan, Anne & Bedford Lochridge, Cleo & George Mayfield (compiler—999 Sunnyside Dr., Columbia 38401), Mark Mayfield, Dianne Pressnell.

NASHVILLE—3-4 May; about 42 observers—Jerry Drewry (compiler). In addition, one *Empidonax* sp. was reported. Black-headed Grosbeak deleted; no details.

MURFREESBORO—3-4 May; Sat. - temp. 45-80°F, windy; Sun. - temp. 34-80°F, no wind; 16 observers—Lou A. Erwin, C.E. Greever, Edith Haynes, Anne L. Hettish (compiler—1018 Lawndale Dr., Murfreesboro 37130), Bob Hettish, John Hooper, Richard B. & Wilma J. Hunter, M. David McCarroll, Larry McFarlin, Ruth V. McMillan, Gene & Ruth Odom, John A. Patten, Alberta Spence, Terry J. Witt. In addition, four *Empidonax* sp. were reported.

1986 SPRING FIELD DAYS

| | Метр | Lawr | Clar | Colu | Nash | Murf | Leba | High | Chat | Knox | Gree | Eliz |
|---|----------|----------|---------------|----------|---------|-------|---------|---------|----------|------|----------|------|
| Common Loon | - | 20 | - | - | 2 | - | 120 | 21 | 1 | 222 | 22 | 1022 |
| Pied-billed Grebe | 2 | | 1 | 2 | 1 | 1 | - | - | | | 20 | 6 |
| Horned Grebe | | | | | | 122 | | | | | | 3 |
| Double-cr. Cormorant | 27 10 | | | 77 | 4 | - | 16 | | | 10 | | |
| Anhinga | 10 | | | - | | 77 | 7.0 | | | | - | |
| American Bittern | | | - | | 1 | ~ | | - | | | 1 | - |
| Least Bittern | - | | | | 1 | | | - | 7.7 | - | | - |
| Great Blue Heron | 14 | 33 | | 1 | 14 | - | 5 | 3 | 99 | 4 | 5 | 3 |
| Great Egret Snowy Egret | -1 | | | | 4 | | | | | 2 | | - |
| 2007) 28.00 | | | | | | | | | | | | |
| Little Blue Heron | 2 | - | | | 700 | .1 | 77 | - | 1 | 3.77 | | |
| Cattle Egret | 8 | 6 | | 4 | 23 | 25 | 5 | 1 | 2 | 10 | 6 | 7 |
| Black-cr. Night-Heron Yellow-cr. Night-Heron | | | ** | 1 | 87 6 | 1 | | | 2 | 1 | - | 1 |
| Gr. White-fronted Goose | | | | 1 | | | | | -5 | | | - |
| | | | | | | | | | | | | |
| Canada Goose | 1 | 12 | 4 | 60 | 623 | 41 | 67 | 44 | 29 | 26 | 20 | 25 |
| Wood Duck Mallard | 16 | 4 | 1 | 14 | 168 | 19 | 1 | 2 | 20 35 | 40 | 32 16 | 23 |
| Maliard Blue-winged Teal | 9 | 2 | | 10 | 24 | | 17 | 1 | 19 | 16 | 4 | 13 |
| Northern Shoveler | | | - | | | | | - | 1 | | -2 | 1 |
| | | | | | | | | | | | | |
| Gadwall | 2.5 | | $\overline{}$ | | - | 3.5 | 7 | - | 5 | | 77 | -55 |
| American Wigeon Redhead | ** | | | 2 | | | | | 4 | | | |
| Ring-necked Duck | | | ** | 1 | | ** | ** | | 1 | - | ** | |
| Lesser Scaup | 2 | | - | | | | | | 4 | | | 25 |
| | | | | | | | | | | | | |
| Bufflehead | - | | | | - | | | | | ~~ | | 6 |
| Red-breasted Merganser Ruddy Duck | 9 | 35 | 1 | 7.7 | 723 | 7.7 | 7.5 | | 7.7 | | - | 2.8 |
| Black Vulture | 1 | 2 | 2 | 8 | 83 | 32 | 6.1 | 4 | 3 | | 2 | |
| Turkey Vulture | 2 | 35 | 28 | 23 | 124 | 30 | 30 | 16 | 10 | 15 | 20 | 35 |
| | | | | | | | | | | | | |
| Osprey | | | | | 3 | | 1 | | 6 | 8 | 2 | 5 |
| Mississippi Kite Northern Harrier | 6 | | | 1 | | == | | | 7 | 1 | | |
| Sharp-shinned Hawk | | 1 | | 2 | 3 | | | | | 1 | 1 | 1 |
| Copper's Hawk | 1 | 1 | - | 1 | 4 | 22 | 1 | | | | 1 | 223 |
| | | | | | | | | | | | | |
| Red-shouldered Hawk | 4 | 4 | 4 | 1 | 9 | | 1 | | 6 | | 7 | 19 |
| Broad-winged Hawk Red-tailed Hawk | 8 | 3 | 1 | 9 | 37 | 6 | 2 14 | 7.0 | 11 | 6 | 10 | 5 |
| American Kestrel | 1 | 2 | 3 | 9 | 41 | 59 | 5 | 1 | ï | 4 | 4 | 5 |
| Ruffed Grouse | | | | | | | - | | | 2 | | 5 |
| | | | | | | | | | | | | |
| Wild Turkëy Northern Bobwhite | 2 26 | 14 25 | 14 | 16 34 | 160 | 33 | 13 | 5 | 33 | 41 | 32 | 12 |
| King Rail | 26 | 23 | 1.0 | 34 | 100 | -33 | 13 | | 3.3 | 41 | -12 | 12 |
| Virginia Rail | | | | 1 | | | | | | | | |
| Sora | | 1 | | 7 | 5 | 1 | 5.77 | 2.7 | | 1. | | 1 |
| N N SECT | 1 | | | 4 | 4 | | | 2 | 82 | | 2 | Ė |
| American Coot Black-bellied Plover | 1 | - | | - | * | | 22 | 4 | 84 | | | - |
| Lesser Golden-Plover | 1 | | 3.06.00 | | | | - | - | | | | |
| Semipalmated Plover | 32 | - 3 | | | 1.3 | | | | 4 | | | |
| Killdeer | 27 | 10 | 6 | 7 | 86 | 52 | 10 | 8 | 27 | 59 | 31 | 39 |
| American Avocet | | | | | 4 | | | | | | - | - |
| Greater Yellowlegs | 25 | 1 | 1225 | | 10 | 2 | 1 | der tes | 18 | 11 | 5 | 2 |
| Lesser Yellowlegs | 78 | 5 | - | 10 | 4.1 | 2 | 2 | | 90 | 23 | 2 | 10 |
| Solitary Sandpiper | 54 | 18 | ++ | 7 | 44 | 13 | 13 | - 2 | 18 | 31 | 46 | 20 |
| Spotted Sandpiper | 19 | 4 | 4 | 1 | 25 | 8 | 2 | | 12 | 4 | 5 | 14 |
| Ruddy Turnstone | 5.23 | 62 | -22 | | 1 | | | *** | | - | | |
| Semipalmated Sandpiper | 10 | 4 | | 1 | 1 | | | | 8 | ** | | ** |
| Western Sandpiper | 1 | 1 | | | 1 | | | | | - | | |
| Least Sandpiper | 114 | 15 | | 1 | 165 | -55 | 55 | 1 | 104 | 1 | | 7.7 |
| White-rumped Sandpiper | 4 | | | | 1 | | | ** | | ** | | |
| Baird's Sandpiper | . 1 | - | 122 | - | - 25 | | 22 | 124 | | | | 4.5 |
| Pectoral Sandpiper | 189 | 2 | | 2 | 10 | 155 | | | 24 | 1 | | 9 |
| Dun1in | 1 | - | *** | | 2 | | | | | | | |
| Dowitcher sp. | | 3 | × 4 | | 1 | 94.34 | | | - | 4 | | 77 |
| Common Snipe | 6 | 1 | N-41 | 1 | 11 | | | | | 4 | 3 | 7 |

| | Memp | Lawr | Clar | Colu | Nash | Murf | Leba | High | Čhat | Knox | Gree | Eliz |
|---|----------|------|----------|----------|------------|-----------|-----------|-----------|-----------|------------|------------|------------|
| American Woodcock | | | | 2 | 9 | | | | _ | 179990 | - | 3 . |
| Wilson's Phalarope | 2 | | | | | | | | | | -2 | |
| Laughing Gull | | | | | | | 7.7 | | 1 | | 77 | |
| Bonaparte's Gull Ring-billed Gull | 3 | 85 | 77 | | 1 32 | | 15 | - | 103 | 32 | | 52 |
| Herring Gull | | en. | | | 6 | | 2 | 11 | | | | |
| Caspian Tern | | | | | 7 | ** | | 3.70 | | . 77 | | 77.0 |
| Forster's Tern Least Tern | 33 | | | | 9 | | 2 | | 4 | | | 10 |
| Rock Dove | 1 | 15 | 17 | 29 | 147 | 30 | 43 | 14 | 64 | . 74 | 46 | 52 |
| Mourning Dove | 105 | 88 | 56 | 55 | 270 | 90 | 90 | 23 | 93 | 246 | 88 | 156 |
| Black-billed Cuckoo Yellow-billed Cuckoo | 100 | 1 | 177 | 1 | 1 | | 1 | = | | | | |
| Common Barn-Dwl | 2 | | 4 | | 5 | 3 | 1 | | | 1 | 1 | |
| Eastern Screech-Owl | | 1 | ** | 1 | 5 | | 10 | | 7 | 4 | 1 | 2 |
| Great Horned Owl | 1 | 45 | | 3 | 5 | | 6 | 1 | 3 | 4 | 2 | 2 |
| Barred Owl | 3 | 7 | 4 | 7 | 15 | 72 | 18 | 2 | 6 | 6 | - | |
| Common Nighthawk Chuck-wills-widow | | 8 | 1 | 5 | 13 | 15 | 21 17 | 1 | 7 2 | 8 12 | 1 | 2 |
| Whip-poor-will | -22 | 3 | 2 | 6 | 41 | 5 | 7 | 3 | 2 | 4 | 3 | 6 |
| Chimney Swift | 45 | 26 | 62 | 65 | 510 | 1000 | 79 | 30 | 119 | 209 | 150 | 283 |
| Ruby-thr. Hummingbird | 20 | 4 | 6 | 4 | 24 | 9 | 1 | 2 | 8 | 16 | 1 | 7 |
| Belted Kingfisher Red-headed Woodpecker | 1 21 | 5 2 | 3 | 6 12 | 15 | 10 | 8 | 2 | 17 | 10 | 1 | 11 |
| Red-hellied Woodpecker | 63 | 22 | 20 | 28 | 98 | 21 | 13 | 8 | 19 | 73 | 23 | 22 |
| | | | | | 5.701 | 2 | ** | | 300 | 190 | 2.3 | 5.5 |
| Yellow-bel. Sapsucker Downy Woodpecker | 17 | 11 | 1 8 | 26 | 76 | 12 | 6 | 1 | 18 | 19 | 19 | 26 |
| Hairy Woodpecker | 3 | 5 | 1 | 7 | 13 | 3 | 2 | 1 | 4 | 2 | 3 | 3 |
| Northern Flicker | 11 | 2 | 4 | 7 | 27 | 10 | 5 | 7 | 19 | 32 | 26 | 35 |
| Pileated Woodpecker | 18 | 6 | 6 | 17 | 36 | 4 | 2 | 2 | 15 | 21 | 21 | 13 |
| Eastern Wood-Pewee | 15 | 6 | .6 | 17 | 43 | 5 | 4 | 1 | 15 | 1 | 4 | |
| Acadian Flycatcher | 15 | 11 | 4 | 2 | 22 | | | | 9 | | | |
| Least Flycatcher | 2 | 13 | 7 | 15 | 1 66 | 5 | 22 | | | 1 | | |
| Eastern Phoebe Great Cr. Flycatcher | 33 | 14 | 6 | 20 | 54 | 13 | 9 | 4 | 12 12 | 30 21 | 12 20 | 45 5 |
| Eastern Kingbird | 21 | 17 | 13 | 19 | 126 | 49 | 55 | 19 | 40 | 20 | 9 | 9 |
| Scissor-t. Flycatcher | 22 | | | | | ++ | 1. | - | | 277 | 77 | 75 |
| Horned Lark Purple Martin | 15 33 | 17 | 45 | 30 | 108 | 42 | 32 | 28 | 124 | 119 | 3 41 | 158 |
| Tree Swallow | 5 | 12 | 1 | 8 | 98 | 57 | 4 | 20 | 111 | 6 | 16 | 12 |
| N. Rough-winged Swallow | 9 | 24 | 36 | 36 | 294 | 10 | 49 | 6 | 90 | 46 | 19 | 106 |
| Bank Swallow | 3 | 7.70 | 2 | 10 | 44 | 10 | 77 | | 14 | | | - |
| Cliff Swallow | 91 | 50 | 250 | 2 | 3154 | | | 2 | 118 | | | 453 |
| Barn Swallow Blue Jay | 29 | 43 | 15 22 | 46 39 | 455 208 | 173 59 | 114 85 | 104 33 | 151 82 | 153 447 | 177 286 | 157 181 |
| American Crow | 31 | 38 | 78 | 41 | 297 | 650 | 90 | 27 | 170 | 203 | 138 | 184 |
| Fish Crow | 8 | | | | - | | | | | | | - |
| Common Raven | 73 | 14 | 20 | | 177 | | 27 | 2 | | | | 6 |
| Carolina Chickadee Tufted Titmouse | 79 | 25 | 35 | 28 32 | 156 | 33 23 | 18 | 7 | 76 54 | 101 | 47 77 | 91 |
| Red-breasted Nuthatch | 1 | | | | 1 | 3 | | - | | 5 | | |
| White-breasted Nuthatch | 3 | 4 | 4 | 7 | 34 | | | 3 | 5 | 15 | 2 | 7 |
| Brown-headed Nuthatch | 55 | 7.75 | 170 | 770 | 377 | 37.5 | | 7.7 | 1 | - | | === |
| Brown Creeper Carolina Wren | 45 | 18 | 20 | 22 | 108 | 10 | 9 | 5 | 1 51 | 115 | 59 | 69 |
| Bewick's Wren | | _ | | 2 | 1. | 3 | 4 | | | | | |
| House Wren | | *** | 40 | 2 | 22 | 13 | 14 | | 1 | 6 | 18 | 9 |
| Sedge Wren | - | | 17.7 | 1 | | | - | 1 | | 1 | - | |
| Marsh Wren | | - | | | 7 | | | -22 | | | 1 | 2 |
| Golden-crowned Kinglet | 75 | 22.5 | | | | 700 | 6.75 | | | | | |
| Ruby-crowned Kinglet | 1.12 | 21 | 1 25 | 2 | 2 | | 16 | 8 | 105 | 31 | 6 | 7 |
| Blue-gr. Gnatcatcher Eastern Bluebird | 112 | 103 | 25 | 29 41 | 191 212 | 99 | 201 | 26 | 105 | 127 97 | 42 86 | 35 52 |
| Veery | 3 | 2 | | 4 | 10 | 1 | 241 | 20 | 1 | 2 | | 32 |
| Gray-cheeked Thrush | 2 | 1 | | | 10 | | | 1 | 3 | - | ** | ** |
| Swainson's Thrush | 53 | - | 5 | 3. | 107 | 17 | - | 1 | 4 | 4 | | 16.00 |
| Hermit Thrush Wood Thrush | 57 | 6 | 1 15 | 16 | 58 | 1 12 | 6 | 1 | 32 | 75 | 7 | 20 |
| American Robin | 42 | 53 | 120 | 72 | 561 | 1200 | 109 | 76 | 152 | 547 | 201 | 389 |
| Gray Catbird | 4 | 8 | 8 | 12 | 69 | 52 | 15 | i | 27 | 11 | 4 | 3 |

| | Memp | Lawr | Clar | Colu | Nash | Murf | Leba | High | Chat | Knox | Gree | Eliz |
|---|---------|------|------|---------|------------|-----------|---------|------|----------|------|-----------|----------|
| Northern Mockingbird | 18 | 33 | 47 | 31 | 142 | 75 | 77 | 40 | 60 | 125 | 64 | 81 |
| Brown Thrasher Water Pipit | 12 | 26 | 20 | 40 | 113 | 36 | 68 | 9 | 36 | 54 | 61 | 26 90 |
| Cedar Waxwing | 6 | 3 | 15 | 11 | 60 | 52 | 5 | | 22 | 50 | 21 45 | 13 |
| Loggerhead Shrike | 4 | 15 | 1 | 11 | 67 | 7 | 12 | | 6 | 2 | 2 | 1 |
| European Starling | 183 | 125 | 169 | 61 | 936 109 | 6000 | 178 | 78 | 275 | 1513 | 406 | 700 |
| White-eyed Vireo Solitary Vireo | 92 | 10 | 19 | 35 | 109 | 10 | 6 2 | 3 | 17 | 41 | 13 | 20 18 |
| Yellow-throated Vireo | 14 | 6 | 1 | 6 | 12 | 1 | 4 | 1 | 12 | 9 | 2 | 6 |
| Warbling Vireo | 10 | | 7 | | 34 | | 1 | | 1 | 222 | 2.2 | 1 |
| Philadelphia Vireo Red-eyed Vireo | 69 | 32 | 15 | 25 | 1114 | 29 | 7 | 9 | 150 | 170 | 1 9 | 36 |
| Blue-winged Warbler | 2 | 3 | 2 | 3 | 31 | 27 | <u></u> | 1 | 1 | 4 | 1 | 20 |
| Golden-winged Warbler | 1 | | | | 2 | | 1 | 1 | | | | *** |
| Tennessee Warbler | 107 | 4 | 38 | 42 | 167 | 26 | 6 | 4 | 44 | 22 | | ** |
| Orange-crowned Warbler Nashville Warbler | 7 | | 5 | | 17 | 2 | 2 | | 2 | 111 | 1 | - |
| Northern Parula | 97 | 12 | 2 | 10 | 25 | | | | 2 | 1 | | 12 |
| Yellow Warbler | 1 | 2 | 12 | 22 | 68 | 7 | 13 | | 21 | 14 | 5 | 34 |
| Chestnut-sided Warbler | - | 2 | 3 | 4 | 12 | 5 | 5 | - | 12 | 4 | 75 | 10 |
| Magnolia Warbler Cape May Warbler | 4 | 3 | - | 2 | 6 2 | 7 5 | 2 | - | 8 11 | 1 4 | 1 | 1 |
| Black-thr. Blue Warbler | | | | | | | 1 | - | | | | 8 |
| Yellow-rumped Warbler | 17 | 2 | 14 | 13 | 129 | 27 | 16 | 13 | 41 | 216 | 28 | 92 |
| Black-thr. Green Warbler | 2 | | - | 10 | 22 | 1 | 6 | 1 | 8 | 27 | | 26 |
| Blackburnian Warbler | - | | | 4 | 2 | 1 | 2 | *** | 4 | | | 1 |
| Yellow-throated Warbler Pine Warbler | 8 | 7 2 | 2 | 5 | 34 | 6 | 3 | 1 | 25 38 | 6 | | 5 |
| Prairie Warbler | 93 | 2 | 328 | 12 | 47 | 19 | 5 | 6 | 28 | 2 | 7 | ï |
| Palm Warbler | 77 | 2 | | 12 | 47 | 19 | 5 | 6 | 28 | 2 | 7 | 1 |
| Bay-breasted Warbler | 1 | 1 | 1 | 5 | 14 | 16 | | | 8 | | | 84.0 |
| Blackpoll Warbler Cerulean Warbler | 21 | 4 | | 7 | 15 | 3 | | | 2 | 1 5 | | 1 |
| Black & white Warbler | | 6 | 4 | 10 | 23 | 3 | 4 | | 10 | 19 | 1 | 31 |
| American Redstart | 6 | 2 | 1 | 11 | 9 | 1 | 1 | 32 | 41 | 1 | 1 | 2 |
| Prothonotary Warbler | 42 | 3 | 5 | 15 5 | 26 13 | 1 | .7 | 144 | 12 | 11 | 2 | 1 5 |
| Worm-eating Warbler Swainson's Warbler | 10 | | 17 | | 1.3 | | 1 | | | 12 | | |
| Ovenbird | 5 | | | 8 | 10 | 3 | | 100 | 2 | 45 | 3 | 66 |
| Northern Waterthrush | 1 | 2 | 1 | 1 | 16 | 5 | 6 | | 2 | 1 | | |
| Louisiana Waterthrush | 15 | 12 | 3 | 9 | 30 | 2 | 2 | | . 5 | | ~~ | 7 |
| Kentucky Warbler Connecticut Warbler | 54 | 9 | 8 | 21 | 49 | 2 | 4 | 1 | 17 | 2 | 1 | 1 |
| Mourning Warbler | | - | | | 1 | | 1 | | | | | |
| Common Yello⊌throat | 27 | 14 | 19 | 30 | 251 | 35 | 30 | 11 | 57 | 101 | 45 | 18 |
| Hooded Warbler | 11 | 4 | 2 | 15 | 13 | | 227 | 122 | 16 | 17 | 3 | 42 |
| Wilson's Warbler Canada Warbler | ** | 1 | | | 3 | | | | 1 | 1 | | |
| Yellow-breasted Chat | 7 | 6 | 10 | 28 | 76 | 9 | 13 | 4 | 27 | 19 | 4 | 1 |
| Summer Tanager | 63 | 14 | 9 | 25 | 64 | 6 | 12 | 5 | 9 | 8 | 4 | 1 |
| Scarlet Tanager | 11 | 8 | 7 | 22 | 60 | 17 | 2 | 4 | 19 | 37 | 8 | 11 |
| Northern Cardinal Rose-breasted Grosbeak | 151 | 99 | 101 | 83 | 580 | 300 21 | 123 | 36 | 170 | 408 | 161 10 | 182 |
| Blue Grosbeak | 2 | 6 | 6 | 4 | 15 | 4 | 3 | | 7 | 5 | 11 | 2 |
| Indigo Bunting | 93 | 11 | 54 | 76 | 303 | 47 | 67 | 8 | 106 | 58 | 11 | 4 |
| Painted Bunting | 1 | 55 | | =5 | - | 3 | 77 | 22 | 33 | 12 | 33 | 30 |
| Dickcissel Rufous-sided Towhee | 6 16 | 31 | 2- | 43 | 163 | 47 | 23 | 5 | 88 | 124 | 69 | 78 |
| Chipping Sparrow | 10 | 11 | 6 | 9 | 76 | . 6 | 8 | 3 | 31 | 39 | 32 | 30 |
| Field Sparrow | 1 | 11 | 8 | 22 | 151 | 58 | 63 | 8 | 95 | 173 | 88 | 32 |
| Lark Sparrow | 21 | 2 | - | | 41 | - | 2 | 3 | 5 | | 4 | 10 |
| Savannah Sparrow Grasshopper Sparrow | | 1 | | | 3 | | 1 | | 2 | 6 | 10 | 3 |
| Fox Sparrow | 122 | 1 | 2777 | | | | | | | | | |
| Song Sparrow | 3.5 | 1 | 5 | 1 | 37 | 14 | 5 | .9 | 68 | 179 | 134 | 309 |
| Lincoln's Sparrow Swamp Sparrow | 3 | 2 | *** | 1- | 11 | 1 | | ** | 12 | 10 | | 11 |
| White-throated Sparrow | 57 | 29 | 6 | 16 | 93 | 31 | 29 | 21 | 22 | 249 | 125 | 48 |
| White-crowned Sparrow | 11 | 1 | 6 | 19 | 22 | 14 | 1 | 2E0 | 02 | 3 | 14 | 11 |
| Dark-eyed Junco | | 1 | | | - | | | | | | 2 | 11 |

| | Метр | Lawr | Clar | Colu | Nash | Murf | Leba | High | Chat | Knox | Gree | Eliz | |
|----------------------|------|------|------|------|------|-------|------|------|------|------|------|------|--|
| Bobolink | | 109 | 13 | 100 | 137 | 50 | | | 6 | 6 | - | 2.2 | |
| Red-winged Blackbird | 440 | 116 | 98 | 170 | 812 | 2500 | 75 | 71 | 251 | 241 | 315 | 127 | |
| Eastern Meadowlark | 57 | 105 | 46 | 63 | 429 | 600 | 96 | 121 | 170 | 205 | 255 | 182 | |
| Rusty Blackbird | | 1 | | | 1 | | 2 | | | | | | |
| Common Grackle | 115 | 78 | 89 | 220 | 683 | 10000 | 221 | 105 | 197 | 463 | 446 | 550 | |
| Brown-headed Cowbird | 171 | 43 | 49 | 84 | 332 | 100 | 86 | 8 | 115 | 131 | 67 | 67 | |
| Orchard Oriole | 29 | 11 | 2 | 23 | 146 | 9 | 20 | 7 | 33 | 20 | 5 | 5 | |
| Northern Oriole | 26 | 3 | 2 | 2 | 33 | 6 | 2 | | 8 | 3 | 2 | 2 | |
| Purple Finch | | | - | 1 | 5 | 1 | 2 | 440 | | 22 | 11 | 25 | |
| House Finch | 1 | | 4 | | 4 | 3 | 1 | 6 | 17 | 9 | 9 | 25 | |
| Pine Siskin | 4 | 45 | | 31 | 37 | 5 | 122 | 18 | 180 | 1 | 1 | 22 | |
| American Goldfinch | 20 | 78 | 59 | 97 | 553 | 69 | 50 | 94 | 179 | 438 | 225 | 525 | |
| Evening Grosbeak | | 13 | | 1.1 | 45 | 2 | - | 12 | 45 | 14 | 3 | 87 | |
| House Sparrow | 1 | 182 | 82 | 52 | 213 | 2500 | 42 | 43 | 60 | 86 | 55 | 110 | |
| Total Species | 144 | 130 | 105 | 144 | 176 | 121 | 127 | 87 | 147 | 136 | 115 | 132 | |

LEBANON—3-4 May; fair & windy; temp. 42-70°F; 10 observers—Ceil & Fred Detlefsen, Robert Hosier, Ruth V. McMillan (compiler—Rt. 2, Box 141, Lascassas 37085), Kay Norris, Ray Pope, Billy & Ramona See, Grace Speck, Melissa Tucker.

HIGHLAND RIM—26 April; Philip Medley (compiler—Rt. 2, Box 173B, Hillsboro 37342).

CHATTANOOGA—3-4 May; Sat. - 0500 to 2100, temp. 49-70°F, windy; Sun. - 0500 to 2200, temp. 40-79°F; 21 observers—Edith Bromley, Angela Davis, Jack & Linda Doyle, Ken & Lil Dubke, Gertrude Fleming, Paul Harris, Albert Jenkins, Ross Kinderman, Jonnie Sue Lyons, Barbara & Mike McMahon, Johnny Parks (compiler—1610 East 17th St., Chattanooga 37404), David & Tom Patton, Janet Phillips, Mary Ann Potter, Peter Robinson, Jim Rowell, Robin Rudd.

KNOXVILLE—27 April; 0500 to 2000; clear; temp. 51-91°F; 27 observers in 14 parties—Jane Beintema, Frank Bills, Sharon Bostick, Jim & Betty Reid Campbell, Howard Chitwood, Bob Collier, Barry Fleming, Brice Gilmer, Paul Hartigan (compiler—824 Thomas Circle, Seymour 37865), Audrey & Ron Hoff, Joe Howell, Susan Hoyle, Hal Hubbs, Burt Jones, Tony Koella, Beth Lacy, Holly Overton, J.B. Owen, Paul Pardue, Truett Patterson, Mark Reeves, Ken & Mable Sanders, Boyd Sharp, Jim Tanner. In addition, two *Empidonax* sp. were reported.

GREENEVILLE—27 April; clear; temp. reaching 91°F; 17 observers—Ben, Orland, & Phine Britton, Tio Carlos, Herbert Cragin, King & Margaret Gaut, Lenard Heckert, Richard & Willie Ruth Nevius, Daniel J. Nieves (compiler—112 Magnolia Dr., Greeneville 37743), Terry R. Nieves, Linda & Roy Northrop, Doug Ratledge, Royal Spees, Virginia Williams. Greater White-fronted Goose (reported incubating eggs) and Ring-necked Pheasant (no established population) were deleted.

ELIZABETHTON—26 April; 0330 to 1930; clear; temp. 50-88°F; 16 observers in 7 parties—Glen Eller (compiler—Rt. 9, Box 1340, Elizabethton 37643).

ROUND TABLE NOTES

SECOND TENNESSEE RECORD OF THE GROOVE-VILLED ANI—On 17 October, 1985, while birding at the Tigrett Wildlife Management Area about 8 km east of Dyersburg, Dyer County, Tennessee, I was attracted by the continuous calling and tail-twirling behavior of a small glossy blackbird in a fence row. Using 12x binoculars, I observed the bird at a distance of 20m for 3 minutes. It had a long rounded tail which it twirled in a circle, first in one direction, then the other way. After the bird flew to a small tree, I noted that its bill was high, curved and distinctly grooved. After about 2 minutes, it flew short distances in "flap and glide" flights, while continuing to twirl its tail and constantly repeat a short, sharp whistle. The bird flew out of sight, but I tracked it by its call and again studied it for several minutes. My total observation time was about 10 minutes, and I eventually lost the bird in thick brush. After consulting the 4th edition of Peterson's "A field guide to the birds", I identified it as a Groove-billed Ani (Crotophaga sulcirostris).

I alerted other birders who searched for the ani. To my knowledge, the only other individual to observe it was Don Manning, who saw it about 400 m west of where I observed it. The only previous Tennessee record of the Groove-billed Ani is that of one shot by a hunter 16 km north of Dyersburg on 29 November 1968. This bird was secured and identified by Kenneth Leggett (Migrant 40:7-9, 1969), and a post-mortem was performed by Albert Ganier (ibid. 40:9-11, 1969). This bird was also in Dyer County. Outside Tennessee, this tropical species has been reported along the Gulf Coast and as far north as Ontario (Amer. Birds 38(2):197, 1984). Most inland records have been in the fall.

WILLIAM G. CRISWELL, Route 4, Box 206, Dyersburg, TN 38024. Accepted 10 March, 1986.

NASHVILLE, TENNESSEE TELEVISION TOWER CASUALTIES, 1984—Avian casualties were collected during the fall of 1984 at the WSMV (formerly WSM) and WKRN (formerly WNGE) television towers at Nashville, Davidson County, Tennessee. This was the twenty-fifth year of this project, and the most recently published report, covering the years 1976-1983, was in the September 1984 *Migrant* (55:53-57). This note summarizes the 1984 results.

WSMV Tower Casualties—The WSMV tower was monitored each day from 1 September through 1 November, except for 2 days when no collection was made. Two hundred fifteen birds of 34 species were found (Table 1). The highest count of casualties was on 27 September when 66 birds were collected, and the next highest count was on 1 October with 24 birds. No unusual weather conditions were recorded by the Weather Bureau on 26-27 September, although there was some ground fog present at the tower on the morning of 27 September. The most numerous species collected was the Tennessee Warbler, with 43 individuals, followed by the Red-eyed Vireo, 24, Bay-breasted Warbler, 23, and Ovenbird, 19. No new species were added to the overall list, and there were no new early or late dates set. The Red-headed Woodpecker found on 23 September was only the second collected during 25 years, and the previous one was on 29 September 1963.

TABLE 1. Television tower casualties, fall 1984, Nashville, Tennessee.

| Species | Season WSMV | Totals | Dates of Occurrence and Number Collected (in parentheses) |
|--|----------------|--------|--|
| Sora | 0 | 1 | Sep 27(1)* |
| American Coot | 0 | 1 | Sep 28(1) |
| Cuckoo, Blkbil. | 0 | 1 | Sep 27(1) |
| Yellow-billed | 2 | 1 | Sep 17(1), 19(1), 24(1) |
| Belted Kingfisher | 0 | 1 | Sep 27(1) |
| Red-h. Woodpecker | 1 | 0 | Sep 23(1) |
| Marsh Wren | 2 | 0 | Sep 23(1), 30(1) |
| Kinglet, Golcr. | 0 | 1 | Oct 21(1) |
| Ruby-crowned | 2 | 0 | Sep 27(1), Nov 1(1) |
| Thrush, Swainson's | 2 | 1 | Sep 5(1), 27(1), Oct 6(1) |
| Wood | 5 | 1 | Sep 27(1)(1), Oct 7(1), 11(2), 12(1) |
| Gray Catbird | 6 | 2 | Sep 27(1)(1), 30(1), Oct 5(1), 6(1), 11(1), 14(1), |
| Gray Catoliu | 0 | - | 20(1) |
| European Starling | 1 | 0 | Oct 4(1) |
| | 3 | 0 | Sep 27(1), Oct 20(1), 25(1) |
| Vireo, Yelthr. | 5 | 0 | Sep 27(3), Oct 1(1), 26(1) |
| Philadelphia | 24 | 0 | Sep 27(6), 29(1), 30(2), Oct 3(2), 4(1), 5(1), |
| Red-eyed | 2.4 | U | 6(2), 7(7), 11(1), 12(1) |
| and the second s | 2.74 | 7 | Sep 6(1), 24(1), 27(16)(2), 28(1), 29(1), 30(4), |
| Warbler, Tennessee | 43 | 100 | |
| 7 44 5 TO 1 - 1 TO 1 TO 1 TO 1 TO 1 TO 1 TO 1 | | | 0et 1(4), 5(5), 6(6)(1), 7(2)(1), 20(3)(2) |
| Orange-crowned | 1 | 0 | Oct 26(1) |
| Nashville | 6 | 0 | Oct 5(1), 6(4), 14(1) |
| N. Parula | 1 | 0 | Sep 27(1) |
| Chestnut-s. | 3 | 1 | Sep 27(2)(1), Oct 5(1) |
| Magnolia | 10 | 3 | Sep 26(1), 27(6), 29(2), Oct 5(1), 6(1), 9(1), 20(1 |
| Yellow-rumped | 3 | 0 | Oct 18(1), 20(2) |
| Black-th, Green | 5 | 0. | Sep 30(1), Oct 6(2), 11(2) |
| Blackburnian | 3 | 0 | Sep 27(2), Oct 6(1) |
| Palm | 4 | 1 | Sep 27(1)(1), 28(1), Oct 1(2) |
| Bay-breasted | 23 | 4 | Sep 11(1), Oct 1(2), 5(4), 6(5), 7(4)(1), 9(1), |
| | | | 20(6)(2), 25(1) |
| Black-and-white | 7 | 0 | Sep 27(4), Oct 1(2), 11(1) |
| Amer. Redstart | 6 | 1 | Sep 27(5), 30(1)(<u>1</u>) |
| Worm-eating | 1 | 0 | Oct 5(1) |
| Ovenbird | 19 | 5 | Sep $24(1)$, $25(\underline{1})$, $26(2)$, $27(7)(\underline{1})$, $28(\underline{3})$, $30(1)$, |
| | | | Oct 1(2), 7(1), 11(1), 20(4) |
| Common Yellowthroat | 10 | 2 | Sep 27(4)(1), 30(3)(1), Oct 11(2), 12(1) |
| Canada Warbler | 0 | 1 | Sep 4(1) |
| Yellow-br. Chat | 3 | 0 | Sep 11(2), 28(1) |
| Scarlet Tanager | 2 | 0 | Sep 27(1), Oct 23(1) |
| Northern Cardinal | 1 | 0 | Sep 27(1) |
| Rose-br. Grosbeak | 1 | 3 | Sep 27(1)(2), 30(1) |
| Indigo Bunting | 4 | 0 | Sep 27(1), Oct 10(1), 11(1), 18(1) |
| Sparrow, Song | 1 | 0 | Oct 6(1) |
| Swamp | 1 | 0 | Oct 9(1) |
| Unidentified | 4 | 0 | |
| Total | 215 | 38 | |
| Number Species | 34 | 19 | |

^{*}Underlined number is from WKRN tower; others from WSMV.

The total collection of 215 at WSMV may be compared with 106 in 1983, 107 in 1982, and 84 in 1981.

WKRN Tower Casualties—Monitoring at the WKRN tower was irregular in comparison with WSMV, and predation by house cats was a continuing problem. Collectors found numerous piles of feathers, many of which were unidentifiable. Because of the irregular collection and cat predation, WKRN results (Table 1) are not comparable with those of previous years or with WSMV results. Thirty-eight individuals of 19 species were collected on 14 days. The most numerous species were the Tennessee Warbler, with 7 collected, and the Ovenbird, with 5 collected. The Belted Kingfisher collected on 27 September was only the second collected, and the previous one was on 28 September 1968.

This long-term project would not have been possible without the interest and active participation of a group of Tennessee Ornithological Society members. Collectors this season were Sandra Bivens, Louise Jackson, Portia MacMillan, Henry Parmer, Ann Tarbell, David Vogt and Miriam Weinstein. We also thank the maintenance staff of WSMV for their assistance. Bird specimens in good condition were donated to the Cumberland Museum and Science Center or to the Warner Park Nature Center.


KATHERINE A. GOODPASTURE, 3407 Hopkins Lane, Nashville, TN 37215. Accepted 15 March, 1986.

COMMON REDPOLL AT SEWANEE, TENNESSEE—During a period of moderately heavy snowfall at about 1300 on 20 December 1985, I observed a single female Common Redpoll (Carduelis flammea) perched on a feeding platform about 5 m from my living room window in Sewanee, Franklin County, Tennessee. At this distance, I could clearly discern the bird's dark chin spot, which separated it from the Pine Siskins (C. pinus) with which the redpoll was feeding. With the use of 7x binoculars I was able to note the reddish cap as well as the following differentiating field marks. In contrast to the siskins, this individual appeared to be less boldly streaked, with lighter brown on the back and rump. The breast and sides were an off-white with some dark brown streaking especially noticeable on the sides. Both my wife Cheri and I observed the bird for approximately 15 minutes during which time I obtained several recognizable photographs taken through the window with a 300 mm telephoto lens.

Although this bird was rather unpredictable and generally would visit my feeder for only 5 to 15 minutes at a time, it stayed in the area until 14 March 1986. One other Sewanee birder, Jerry Ingles, was able to observe the bird in early January. During the numerous occasions I had to observe the bird feeding, I was rather surprised to find that this individual never extracted sunflower seeds by cracking their husks, but instead it scavenged for bits of seed dropped by the other finches.

JAMES R. PETERS, SPO 1157, University of the South, Sewanee, TN 37375. Accepted 25 August 1986.

THE SEASON



SPRING: 1 MARCH - 30 MAY 1986

This was, again, a very dry season, with less than half the normal amount of rainfall in middle and east Tennessee. The drought was apparently not as severe in the extreme western part of the state. The regional compilers differed in their assessments of temperatures, which were described as cool in the west, above normal in middle Tennessee, and generally above normal in the Ridge and Valley Region.

The warbler migration in middle Tennessee was described as dull, and the passerine migration in the Ridge and Valley Region was described as unspectacular. Judging from the reports from the other areas, these assessments probably apply statewide. Among the unusual passerines reported were a Harris' Sparrow and a LeConte's Sparrow in west Tennessee in March, and a Sharp-tailed Sparrow and Black-headed Grosbeak in middle Tennessee in May. The nonpasserine migration seemed to be somewhat more exciting. There was the usual abundance of shorebird reports, and several reports of rare species. These included the highest number of Short-eared Owls reported in the state in several years at Duck River Unit in March, a Purple Gallinule in Stewart County, and Least Terns at Cross Creeks refuge. A Great Black-backed Gull was reported at Paris Landing in March. This is probably the sixth report from the state, although the lack of substantiating details supporting some previous reports makes evaluating them difficult. Finally, a bird tentatively identified as a Clapper Rail was found in Elizabethton in April.

As most readers know, this is the first year of the Tennessee Breeding Bird Atlas project, and many began atlas fieldwork this season. Among the exciting results received to date are Bachman's Sparrows in Giles County for the second year, several Brown Creepers in west Tennessee river bottoms, and the first nest record of the Rose-breasted Grosbeak in the Cumberland Mountains of Tennessee.

Readers will notice that the name of the Central Plateau and Basin Region is changed to the Highland Rim and Basin Region. The new name is closer to the commonly used names for the physiographic regions making up middle Tennessee, i.e. the Highland Rim and Central Basin. An updated map will appear in the next issue.—CHARLES P. NICHOLSON.

Abbreviations used in the following reports include: ers - earliest report of season; lrs - latest report of season; max - maximum number during season; m.ob. - many observers.

WESTERN COASTAL PLAIN REGION—This season can best be characterized by low rainfall and cool temperatures. With a decrease in the flow of southerly winds, the number of species and individual migrants was lower than in preceding years. Sightings of infrequently recorded species were made by field observers new to west Tennessee. These conscientious, consistent observations greatly enhanced the coverage within the region. Although the number of species was low for 1986, several early dates were recorded.

Loon-Crane: Common Loon: 16 Mar (3) PLS (DWB); 23 Mar (21), 4 May (3) Pace Point, HNC (JCR). Horned Grebe: 8 Mar - 10 Apr (24-1) PLS (JCR). Double-crested Cormorant: 29 Mar (4) PLS (JCR); 12 Apr (110) Bass Landing, DeSoto Co., MS/SBC line (BBC, LCC); 13 Apr (76) same area (MTOS). Anhinga: 27 Apr (kettle of 10 riding thermals) SFP (LCC et al.); 15 May (2) Waller Pond, Anderson-Tully WMA, LDC (Dave Vandergriff, Paul B. Hamel, RPF). Green-backed Heron: 19 Mar (1) Wapanocca NWR. ARK (HBD, Sue C. Ferguson, Selma L. Lewis). Tundra Swan: 15 Mar (1) Gooch WMA, Obion Co. (JCR). Canvasback: 16 Mar (31) BRF (DWB, JCR). Ruddy Duck: 9 Mar (2000), 23 Apr (650) Robco Lake, SBC (BBC, LCC). Mississippi Kite: 19 May (2 ad, 1 imm) Nixon Cr. at Forked Deer, HYC (RPF, PIK); 31 May (1) HWR (Susan N. McWhirter, N.P. McWhirter et al.). Cooper's Hawk: 12 Apr (1) PEF (JWR); 27 Apr (1) North Lake, SBC (LCC); 4 May (1) BRF (JCR). Golden Eagle: 1 Mar (1 imm) BSU (JCR). Sora Rail: 13 Apr (1) ESL (JRW); 27 Apr (2) PEF (MGW); 28 Apr (1) ESL (Diane P. Bean, DAD, JRW, MGW); 30 May (1) ESL (CHB, DAD, MGW). American Coot: 16 Mar (1600+) PLS (DWB); 13 Apr (1190) PLS (JCR). SANDHILL CRANE: 13, 18 May (2 ad) Fort Pillow State Park, LDC (RPF, Robert L. Brown, Dick Whittington).

Plover—Gull: Black-bellied Plover: 27 Apr (1) ESL (WRP). Lesser Golden-Plover: 15 Mar (206) Lake and Dyer Cos. (JCR). Semipalmated Plover: 27 Apr (32), 11 May (31) ESL (JRW). Lesser Yellowlegs: 27 Apr (78) ESL (MTOS). Solitary Sandpiper: 27 Apr (54) ESL (MTOS); 28 Apr (102) ESL (CHB, DAD, MGW); 3 May (114), 4 May (92), 11 May (38), 18 May (6) ESL (JRW). Upland Sandpiper: 12 Apr (8) PEF (JRW). Semipalmated Sandpiper: 16 May (27) PEF (JRW); 18 May (365) ESL (JRW). Western Sandpiper: 27 Apr (1), 18 May (1) ESL (JRW). White-rumped Sandpiper: 27 Apr (4), 11 May (13), 18 May (57) ESL (JRW). Baird's Sandpiper: 27 Apr (1), 1 May (1) ESL (JRW). Dunlin: 27 Apr - 11 May (1), 18 May (2) ESL (JRW). Short-billed Dowitcher: 13 Apr (1), 18 May (1) ESL (JRW); 9 May (10) LKC (JCR). Wilson's Phalarope: 27 Apr (2) ESL (WRP). Laughing Gull: 9 May (1) Island 13, LKC (JCR). Bonaparte's Gull: 2 Mar (3) PLS (DWB). GREAT BLACK-BACKED GULL: 2-3 Mar (1) PLS (DWB); 8 Mar (1) PLS (ECC, MHI).

Swallow-Finch: Cliff Swallow: 30 Mar (2) PLS (JCR). Barn Swallow: 23 Mar (4) Robco Lake, SBC, 23 Mar (8) MEM, 6 May (72) MEM (BBC, LCC). Fish Crow: 23 Mar (2) MEM (BBC); max 11 Apr (10) OVP (BBC). Brown Creeper: 1 May (1) N of Rossville, Fayette Co.; 5 May (1) Forked Deer River, Madison Co.; 14 May (1) Reelfoot Lake, LKC; 23 May (1) Obion River, HNC (RPF). Water Pipit: 9 May (1) LKC (JCR). Solitary Vireo: 28 Mar (1) Big Hill Pond, McNairy Co. (MGW). Swainson's Warbler: 14 Apr (1) SFP (LCC, HBD); Apr-May (0) OVP (BBC); 31 May (14) along Hatchie River, HYC (BBC). Vesper Sparrow: 23 Mar (1) BRF; 13 Apr (1) BSU (JCR).

LeConte's Sparrow: 29 Mar (1) BRF (JCR). Lincoln's Sparrow: 11 Mar (2) MEM (JRW); 16 Mar (1) MEM (MGW); 16 Mar (1) BRF, 9 May (1) LKC (JCR). Harris' Sparrow: 16 Mar (1) BRF (DWB, JCR); 21 Mar (1) BRF (MHI). House Finch: small numbers through 30 Mar, uncommon except at one location in SE MEM, where they are nesting (BBC).

Locations: BRF - Britton Ford, Henry Co.: BSU - Big Sandy Unit, Tennessee Nat. Wildl. Refuge, Henry Co.; ESL - Ensley Sewage Lagoons, Shelby Co.; HNC - Henry Co.; LDC - Lauderdale Co.; LKC - Lake Co.; MEM Memphis; OVP - Overton Park, Shelby Co.; PEF - Shelby Co. Penal Farm; PLS - Paris Landing State Park, Henry Co.; SBC - Shelby Co.; SFP - Shelby Forest State Park, Shelby Co.

MARTHA G. WALDRON, 1626 Yorkshire Dr., Memphis, TN 38119.

HIGHLAND RIM AND BASIN REGION (formerly Central Plateau and Basin Region)-Migration this spring caused little excitement among field observers until mid-May, when a flurry of unusual sightings occurred. Prior to that, generally warm, dry weather with southerly winds produced excellent conditions for migration (as in the spring of 1985), thus reducing numbers of birds encountered on the ground. Rainfall in March (2.29 in) was well below normal in Nashville, while that for April (0.52 in) was the lowest accumulation ever recorded for that month; May precipitation (3.36 in) was an inch deficient with most of it falling toward the end of the month. Temperatures were above normal during the period though some cooler days occurred (e.g., on 23 April a record low of 31°F was recorded in Nashville, followed on 25 April by a record-tying high of 87°F; at least 2 other record highs were recorded in that month). As a result, leaf emergence was ahead of schedule for many species of plants.

Fair numbers of unusual species were reported during the season, with a few occurring in March and quite a few in May. Early reports included Lapland Longspur and Brewer's Blackbird in Stewart and Lawrence Cos.. respectively, and up to 7 Short-eared Owls in Humphreys Co. Some of the highlights of the May migration included Peregrines, Purple Gallinule, and Sharp-tailed Sparrow in Stewart Co., Bachman's Sparrows in Giles and Stewart Cos., and Black-headed Grosbeak in Davidson Co. About a normal number of early and late dates of arrival and departure were reported during the season, with early swallows being perhaps the most notable among these.

The warbler migration through the region was a dull one by any standards; few reports came in of great numbers of warblers in the woods. though a few high daily species totals were reported. To bring some perspective to the migration of these species through the region, Ruth V. McMillan agreed to undertake an analysis of spring count warbler data from middle Tennessee. The results of her study (Table 1) reveal some interesting trends in the relative abundance of the various warbler species in the region over the past 26 years. Note particularly that 4 transient warblers—Tennessee, Magnolia, Bay-breasted, and Blackpoll, all probably benefiting from the outbreak of spruce budworms and gypsy moths in the northern coniferous forests—have become increasingly abundant over the past 26 years, while no warbler breeding in the region has increased. On the other hand, several warblers which breed in the region are declining in abundance on the spring counts, with the Cerulean and Redstart being the most drastically affected. Regional observers interested in analyzing spring count data of other groups of birds for future spring reports from the region are encouraged to contact the compiler.

The volume of records submitted for inclusion in this report showed a significant increase over the volume submitted in any of the past three spring migration seasons, with many sightings reported on the "Unusual Sighting" cards available from TWRA. However, some areas of the region remain poorly represented in the following report, even though significant numbers of field observers reside in or near them. Most notable among these areas are those covered by the Highland Rim, Columbia, and Nashville Chapters of the TOS. Observers in these areas are in a position to contribute many records which will enrich our understanding of the region's birdlife and that of birds students in the decades and centuries to come. So, to the many silent fieldworkers who reside among us, let me issue a request to hear from you at the end of future seasons.

I thank the many observers who did submit records for this report, which is more the result of their work than my own. I also thank Ruth McMillan for the considerable effort which she expended in analyzing the spring warbler data presented herein. Finally, I am indebted to John C. Robinson for prompt and careful review of earlier versions of this report.

Loon-Night-Heron: Common Loon: 3 May (2) OHL (LFK), Irs. Piedbilled Grebe: 19 May (1) GRP (DTC, JPC), lrs. American Bittern: 2-6 Apr, 7 May (up to 4) CCNWR (JCR, DWB); 12 Apr - 18 May (up to 4) BEC (JCR, DWB); 27 Apr - 5 May (1) MC (DFV, CGD, CKC, et al.); only reports. Least Bittern: 3 May (1) marsh SE of ACI on Cumberland R. (RJM), ers (in possible breeding habitat); 4-30 May (2) BEC (JCR, CKC, et al.), also in suitable breeding habitat. Great Egret: through EOP (up to 3) CCNWR (JCR); 4 May (2) CCP Ponds (BHS, SJS), few WMC records; 4, 30 May (2) GSP (DTC, JPC); only reports. Snowy Egret: 4-16 May (1) MC (CGD, CKC, RJM, m.ob.); 8-12 May (1) CCNWR (JCR); 18 May (2) BEC (JCR); high number of reports. Little Blue Heron: 12 Apr (3 ad) BEC (JCR, DWB); 6 May (2 ad) BEC (JCR, DWB); 8-18 May (2 ad) CCNWR (JCR, CKC), late if not possible breeders; 22-23 May (1 second yr.) CCNWR (JCR); only reports. Cattle Egret: 3 May (5) near BEC (DWB), ers; 5 May (3) CCNWR (JCR); 26 May (1) RUC (TJW, DMM); 27 May (1) CCNWR (JCR); only reports. Green-backed Heron: 7 Apr (1) CCNWR (EJW), ers. Yellowcrowned Night-Heron: 5 Apr (5) WSB (DWB), ers; 21 Apr (1) Cheatham WMA, CHC (EJW).

Goose—Duck: Greater White-fronted Goose: 9 Mar (9) CCNWR (JCR), lrs; only report. Green-winged Teal: 10 Apr (2) CCNWR (JCR), lrs. American Black Duck: 25 May (1) GSP (DTC, JPC), lrs. Blue-winged Teal: 25 Feb (1) CCNWR (JCR), ers; 25 May (1) LWP (DTC, JPC), lrs. Northern Shoveler: 26 Apr (1) MC (SJS, CGD), lrs. Gadwall: 17 Apr (6) GSP (DTC, JPC), lrs. American Wigeon: 16 Apr (1) FRR (SJS, JCR), lrs. Canvasback: 6 Apr (1) Kentucky L., SWC (JCR), lrs. Redhead: 30 Mar (1) MOP (DCC), lrs. Ring-necked Duck: 13 Mar (317) CCNWR (JCR), max; 26 Apr (1) BEC (JCR), lrs. Lesser Scaup: 25 Mar (172) CCNWR (JCR), max; through EOP (1) MOP (WNJ et al.), lrs. Bufflehead: 12 Apr (3) CCNWR (JCR), lrs. Hooded Merganser: 6 May (1) CCNWR (JCR), lrs. Common Merganser: 1 Mar (17) DRU (JCR), lrs; only report. Red-breased Merganser: 6 May (1) CCNWR (JCR), lrs. Ruddy Duck: 5 Apr (2) FRR (SJS), lrs.

TABLE 1. Rank¹ of warblers on spring counts in the Highland Rim and Basin Region, 1961-1986.

| | RANK | DURING Y | EARS |
|--------------------------|---------|----------|------|
| SPECIES | 1961-70 | | |
| Blue-winged Warbler | 12 | 9 | 12 |
| Golden-winged Warbler | 29 | 30 | 30 |
| Tennessee Warbler* | 10 | 4 | 2 |
| Orange-crowned Warbler | 32 | 34 | 32 |
| Nashville Warbler | 22 | 18 | 23 |
| Northern Parula | 20 | 19 | 18 |
| Yellow Warbler | 6 | 7 | 5 |
| Chestnut-sided Warbler | 23 | 26 | 19 |
| Magnolia Warbler* | 27 | 27 | 17 |
| Cape May Warbler | 28 | 25 | 29 |
| Black-thr. Blue Warbler | 31 | 35 | 34 |
| Yellow-rumped Warbler | 1 | 2 | 3 |
| Black-thr. Green Warbler | 19 | 22 | 13 |
| Blackburnian Warbler | 26 | 29 | 26 |
| Yellow-throated Warbler | 14 | 17 | 15 |
| Pine Warbler | 30 | 28 | 28 |
| Prairie Warbler | 9 | 5 | 8 |
| Palm Warbler | 4 | 6 | 4 |
| Bay-breasted Warbler* | 25 | 24 | 14 |
| Blackpoll Warbler | 15 | 11 | 6 |
| Cerulean Warbler*** | 8 | 15 | 21 |
| Black-and-white Warbler | 13 | 13 | 11 |
| American Redstart** | 16 | 21 | 24 |
| Prothonotary Warbler | 7 | 10 | 9 |
| Worm-eating Warbler | 21 | 20 | 20 |
| Swainson's Warbler | 35 | 33 | 35 |
| Ovenbird** | 18 | 14 | 25 |
| Northern Waterthrush | 24 | 23 | 27 |
| Louisiana Waterthrush** | 11 | 12 | 16 |
| Kentucky Warbler** | 5 | 8 | 10 |
| Connecticut Warbler | 35 | 36 | 37 |
| Mourning Warbler | 37 | 37 | 36 |
| Common Yellowthroat | 2 | 1 | 1 |
| Hooded Warbler** | 17 | 16 | 22 |
| Wilson's Warbler | 32 | 32 | 33 |
| Canada Warbler | 34 | 31 | 31 |
| Yellow-breasted Chat** | 3 | 3 | 7 |

¹Rank determined by adding totals for each warbler on all spring counts in middle Tennessee for the period and by ranking the results (with 1 being the most abundant species and 37 the least abundant); * - species increasing in abundance; ** - species declining moderately; *** - species declining drastically.

Osprey-Falcon: Osprey: 31 Mar (1) Long Hunter S.P., DVC (RVM), ers; 24 May (3) S OVC (DJS, SJS); 25 May (1) Celina, CYC (DJS, SJS); last 2 sightings are of very late birds if not of possible breeders. Bald Eagle: 23 Mar (5 imm) DRU (JCR), last report of wintering birds. Northern Harrier: 9 Mar (24 seen simultaneously) DRU (JCR), very high number; 4 May (1) BEC (JCR); 17 May (1) Whitleyville, JAC (SDB); 21 May (1 imm) Putnam Co. (Joseph W. Wahl) lrs; very late. Sharp-shinned Hawk: 18 reports (6 Mar 25 May), double the 1985 data (mainly due to 8 reports from CCNWR). Cooper's Hawk: 16 reports (4 Mar - 26 May), also double the 1985 data (with only 1 report from CCNWR); many of the May birds were probable breeders (see Summer 86 report). Broad-winged Hawk: 5 Apr (1) FTD (JCR), ers. Golden Eagle: 5 Apr (1 imm) Model Fire Tower, LBL (DWB), only rep. Merlin: 27 Mar (1) LBL (DWB); 26 Apr (1) FTN (CGD et al.); only reports. Peregrine Falcon: 13 Mar (1 ad male) CCNWR (JCR); 5 May (1 ad male) CCNWR (JCR); 15 May (1 ad male) CCNWR (JCR); only reports; as JCR noted. CCNWR was definitely "THE place to be for migrating Peregrines this spring."

Turkey—Crane: Wild Turkey: 24 Mar (6) Lebanon S.F., WLC (RVM); 28 Mar (3) WSB (DWB); 6 Apr (14) Shacklett, CHC (CKC); only reports. King Rail: 5 May (1) marsh southeast of ACI on Cumberland R., CHC (RJM), only report. Virginia Rail: 29 Mar (1) MOP (BHS, DCC), ers; 6 May (1) BEC (JCR, DWB), lrs; only reports. Sora: 9 Mar (1), 12 Apr (1), 12, 17 May (1) CCNWR (JCR), first sighting ers; 29 Mar (4) MOP (BHS, DCC); 12-27 Apr (up to 6) CCSP (JCR), max; Apr-May (1) Anne Hettish Bird Sanc., RUC (ALH, DMM, TJW, Eula Kellor); 4-18 May (2) BEC (JCR). PURPLE GALLINULE: 18 May - EOP (1 ad) BEC (JCR, DWB, CKC), very rare in region. Common Moorhen: 31 May (1) CCNWR (DWB), only report. American Coot: 30 May (1) BEC (DWB, JCR), lrs. Sandhill Crane: 2 Mar (30), 7 Mar (22), 8 Mar (160), 9 Mar (75), 10 Mar (25) PIC (Robbie C. Hassler), only reports; according to this observer, the crane migration was "a bit above normal."

Golden-Plover-Snipe: Lesser Golden-Plover: 8 Mar (2) CCNWR (JCR, ECC, MHI, DWB), ers; 14 Mar (60) CCNWR (JCR); 15 Mar (105) DRU (CKC, MHI); 15-23 Mar (up to 192) LOR (DJS), max; 13 Apr (1) CCSP (JCR), Irs. Black-bellied Plover: 15 Mar (1) LOR (DJS), ers; 30 Mar (1) ROP (DJS): only reports (very early). Semipalmated Plover: 27 Apr (3) SWC (JCR, DWB), ers; 19 May (24) LWP (DTC, JPC), max; 25 May (11) GSP (DTC, JPC), Irs. American Avocet: 4 May (4) Cage's Bend, SUC (DRJ, DTC, JPC), 7th NA spring record. Greater Yellowlegs: 9 Mar (3) CCNWR/DRU (JCR), ers; 14 May (5) CCNWR (JCR), lrs. Lesser Yellowlegs: 8 Mar (1) CCNWR (DWB), ers; 20 May (3) CCNWR (JCR), lrs. Willet: 27 Apr (1) MC (DFV), only report. Spotted Sandpiper: 10 Apr (1) CCP ponds (SJS), ers; 31 May (1) CCSP (JCR), lrs. Upland Sandpiper: 27 Apr (1) SWC (JCR), only report. Ruddy Turnstone: 3-4 May (1) LWP (DRJ, DTC, JPC), only report; 12th NA record and first since 1979. Sanderling: 11 May (1) LWP (DTC, JPC), only report. Semipalmated Sandpiper: 27 Apr (1) CCSP (JCR), ers; 31 May (3) CCNWR (DWB), lrs. Western Sandpiper: 11 May (10) GSP (DTC, JPC), ers and max; 25 May (6) LWP (DTC, JPC), lrs. Least Sandpiper: 14 Mar (1) CCNWR (DWB), ers; 4 May (70) LWP (DRJ, DTC, JPC), max; 25 May (23) LWP and GSP (DTC, JPC), lrs. Whiterumped Sandpiper: 3 May (1) near ACI (CGD et al.), ers; 19 May (16) LWP (DTC, JPC), max; 29 May (1) Hillsboro Pond, CFC (James R. Peters); 31 May (2) CCNWR (DWB) lrs. Baird's Sandpiper: 19, 30 May (4/4) LWP/GSP (DTC, JPC), only reports. Pectoral Sandpiper: 8 Mar (2) CCNWR (DWB), ers; 19 Mar (72) CCNWR (JCR), max; 19 May (1) LWP (DTC, JPC), lrs. Dunlin: 3 May (2) SUC (DRJ, DTC, JPC); 11-12, 25 May (2/1) LWP (DTC, JPC, CKC); only reports. Stilt Sandpiper: 12 May (1) LWP (CKC, MHI), only report. Dowitcher sp.: 11 May (22/20) GRP/GSP (DTC, JPC), max; 18 May (2) MC (DFV). Short-billed Dowitcher: 16 May (4) CCNWR (JCR), only report; this bird was noted to have "pale cinnamon color on breast and belly with whitish lower belly and undertail coverts," characteristics reasonably definitive of this species in alternate plumage (Birding, 15:151-166, 1983). Common Snipe: 12 Apr (77) SWC (JCR), max; 6 May (1) BEC (JCR, DWB), lrs.

Gull—Tern: Bonaparte's Gull: 3 May (1) OHL (LFK), lrs. Herring Gull: 13 Apr (1 ad) Laurel Hill L., LWC (DJS), very few LWC records. Forster's Tern: 9 Apr (4) Gray's Landing, LBL (DWB), ers; 2 May (19) CCNWR (JCR), max; 7 Jun (2) ROP (DJS), lrs. LEAST TERN: 18 May (3) CCNWR (JCR), very infrequent migrant in the region. Black Tern: 18 May (5) CCNWR (JCR), only report.

Cuckoo-Lark: Black-billed Cuckoo: 4 May (1) south-central SUC (JPC. DTC); 8 May (1) FTC (JCR); 12 May (1) CFC (MDH); 21 May (1) Bumpus Mills, SWC (DWB); only reports. SHORT-EARED OWL: 23, 30 Mar (5/7) DRU (JCR/CKC, ECC, MHI), only report; noted earlier at this site, but documentation not made available. Whip-poor-will: 30 Mar (1 calling) LEF (BHS), ers. Ruby-throated Hummingbird: 5 Apr (1 male) LEF (BHS), ers. Yellow-bellied Sapsucker: 3 May (1) LEF (BHS), lrs. Olive-sided Flycatcher: 26 Apr (1) LBL (JCR), ers; 24 May (4) S OVC (SJS, DJS), max; 31 May (1) near Sunrise, HIC (BHS, SJS), lrs. Acadian Flycatcher: 18 Apr (1) LEF (BHS), ers. Willow Flycatcher: 20 Apr (1 singing) WSB (DWB), ers; remarkably early. Least Flycatcher: 4 May (1) Percy Priest L. area (Audrey D. Perry, Ray Fleischer), ers; 26 May (1 singing in a small hillside orchard at c. 275 m elevation) S of Gainesboro, JAC (SJS, DJS), lrs; latter record possibly of a territorial male. Eastern Kingbird: 14 Mar (1 well seen in scope) Moores Ln. exit off I-65, WMC (CKC), ers; exceptionally early (within 2 days of NA early date). Scissor-tailed Flycatcher: 4 May (1 ad) Rt. 231, WLC (Kay Norris), only report. Horned Lark: reported from LWC, RUC, and SWC; see Summer 86 report.

Martin—Wren: Purple Martin: 4 Mar (1) Coleman L., DVC (LFK), ers; 10 Mar (1) Cannon Co. (James Vance, fide ALH). Tree Swallow: 26 Feb (4) DRU (JCR), ers. Northern Rough-winged Swallow: 17 Mar (2) MTC (JCR), ers. Bank Swallow: 30 May (1+) GSP (DTC, JPC), lrs; possibly breeding? Cliff Swallow: 27 Mar (10) CCSP (JCR), ers; 3-4 May (3150) CHC (CGD et al.), max; double the 1985 CHC max. Barn Swallow: 5 Mar (1) LOR (DJS), ers; new LWC early spring date. Blue Jay: 12 Apr (62) CCSP (JCR), max; no other reports of migrating jays were submitted although this phenomenon is easy to detect from mid- to late April. Red-breasted Nuthatch: 3 May (1) FTN (CGD et al.), lrs. BEWICK'S WREN: 7 Apr (1 singing) DOV (JCR), ers; see Summer 86 report. Marsh Wren: 13 Apr (1) CCSP (JCR), ers; 26 Apr - 18 May (up to 4) BEC (JCR), max (found in dry, brushy fields). Sedge Wren: 14 Mar (1) CCNWR (DWB), ers.

Kinglet-Vireo: Golden-crowned Kinglet: 9 Apr (2) LBL (JCR), lrs. Bluegray Gnatcatcher: 6 Mar (1) PIC (J. David Hassler), ers. Swainson's

Thrush: 6 Apr (1 stunned by window) Overbrook Dr., NASH (JOE), ers; new NA early spring date. Hermit Thrush: 21 April (2) Montgomery Bell S.P., DIC (JCR), lrs. Wood Thrush: 5 Apr (1) WSB (DWB), ers. Water Pipit: 8 May (1) CCSP (JCR), lrs. Cedar Waxwing: large flocks noted in NASH (Deane McGurk) and LWC (DJS) in Mar. White-eyed Vireo: 5 Apr (2/1) FTD/WSB (JCR/DWB), ers; 6 Apr (18) LBL (JCR), notable early max. Solitary Vireo: 8 May (1) CLA (AHH), lrs. Yellow-throated Vireo: 1 Apr (1) FTD (JCR), ers. Warbling Vireo: 9 Apr (1) CCNWR (JCR), ers. Philadelphia Vireo: 22 Apr (2) WSB (DWB), ers. Red-eyed Vireo: 31 Mar (1) Dunbar Cave S.P., MTC (EJW), ers.

Warbler-Chat: Blue-winged Warbler: 6 Apr (1/1) BAS/CCNWR (DFV, KAG/DWB), ers; 4 May (1 singing a typical Golden-winged song) Boyd Mill Pk., WMC (SJS, BHS), obviously, caution is necessitated in calling Golden-winged Warblers by song; do Golden-wings sing Blue-wings' songs? Golden-winged Warbler: 9 Apr (1) WSB (DWB), ers; exceptionally early. Tennessee Warbler: 9 Apr (6) WSB (DWB), ers. Orange-crowned Warbler: 23 Mar (1) DRU (JCR), ers; 28 Mar (1) BAR (DWB); 15-22 Apr (up to 2) Oaklands P., RUC (TJW, DMM, ALH), lrs; 16 Apr (1) RDL (SJS, JCR). Nashville Warbler: 12 Apr (1) BEC (JCR), ers. Northern Parula: 27 Mar (1 well described) Rivermont Dr., Clarksville, MTC (AHH), ers; 31 Mar (1) LEF (SJS), ties NA early spring date. Black-throated Green Warbler; decidedly more uncommon than in most years; however, note increase of this species during the past 6 years (Table 1). Yellow-throated Warbler: 25 Mar (1) WSB (DWB), ers. Prairie Warbler: 6 Apr (1/4) RDL/LBL (CKC/JCR), ers. Palm Warbler: 7 May (1) CCNWR (JCR), lrs. Worm-eating Warbler: 8 Apr (1) LEF (SJS), ers. Northern Waterthrush: 20 May (1) CCNWR (JCR), Irs. Louisiana Waterthrush: 14 Mar (1) BAS (KAG), ers. Kentucky Warbler: 15 Apr (2) Model Fire Tower, LBL (DWB), ers. Connecticut Warbler: 20 May (1) BAH (RWS), only report. Mourning Warbler: 4 May (1) PWP (DFV), ers; first sighting in the Warner Parks; 15 May (1) N HIC (BHS); 15 May (1) BAH (RWS); 30 May (1) LBL (DWB); only reports. Common Yellowthroat: 28 Mar (1) BAR (DWB), ers. Hooded Warbler: 6 Apr (1) LBL (JCR), ers. Yellow-breasted Chat: 26 Apr (5) LBL (JCR), ers.

Tanager—Longspur: Scarlet Tanager: 15 Apr (1) LBL (DWB), ers; 11-24 May (1 "orange variant") RDL (fide ATT); 31 May (another orange variant) Dunbar Cave S.P., MTC (AHH); probably not as uncommon as originally thought by the RDL contingent. BLACK-HEADED GROSBEAK: 4 May (1 female) Sherwood Dr., NASH (ATT), 5th NA record and first since 1975; it should be noted that females of this species and Rosebreasted Grosbeak are less easily separable than the field guides suggest. Indigo Bunting: 21 Mar (1) LOR (DJS, Don K. Simbeck), ers (possibly an overwintering bird). BACHMAN'S SPARROW: 26-27 Apr (1) FTC (JCR), ers; 7 May (2 singing) Fall River, Giles Co. (DJS), returnees to last year's site. Chipping Sparrow: 14 Mar (1) CFC (MDH), ers. Vesper Sparrow: 12 Apr (3) BEC (JCR), lrs. Lark Sparrow: 21 Apr (2) Simmons Bluff Rd., WLC (RVM et al.), ers; 6 May (1) CCNWR (JCR), only other report of a migrant; see Summer 86 report. Savannah Sparrow: 14 May (1) CCNWR (JCR), lrs. Grasshopper Sparrow: 4 May (2) Del Rio Pike, WMC (Thomas Guschke, Mary Anne Guschke), ers. LE CONTE'S SPARROW: 9 Mar (1) CCNWR (JCR), only report. SHARP-TAILED SPARROW: 18 May (1) BEC (JCR), only report. Fox Sparrow: 3 May (1) LWC (DJS), lrs; very late. Lincoln's Sparrow: 28 Mar (3) BAR (DWB), ers; 14 May (2) CCNWR (JCR), lrs. Dark-eyed Junco: 3 May (1) LWC (DJS), lrs. LAPLAND LONGSPUR: 8 Mar (3) CCNWR (DWB), only report.

Bobolink—Grosbeak: Bobolink: 21 Apr (2) WLC (RVM et al.), ers; 28 May (3) CCNWR (JCR), lrs. Rusty Blackbird: 4 May (1) SUC (DTC, JPC), lrs. BREWER'S BLACKBIRD: 7 Mar (2) St. Joseph, LWC (DJS), only report. Orchard Oriole: 9 Apr (2) 3 WSB (DWB), ers. Northern ("Baltimore") Oriole: 26 Apr (2/1) LBL/NASH (JCR, DWB/Paul B. Hamel), ers. Purple Finch: 18 May (1) Dunaway Dr., NASH (John N. and Heather Riggins), lrs; new NA late spring date. Pine Siskin: 15 May (1) N of Westpoint, LWC (DJS), very late for LWC; 22-23 May (2) Tullahoma, CFC (MDH), also very late; 27 May (1) DOV (JCR), lrs. Evening Grosbeak: 16 May (1) Lawndale Dr., Murfreesboro, RUC (ALH); 23-24 May (1) Old Fort St., Tullahoma, CFC (Chloe and Howell Peebles), lrs.

Corrigenda: Migrant 55:69, 1984: delete the entry for American Black Duck; Migrant 55:70, 1984: at the end of the entry for Bewick's Wren change "WLC (RVM)" to "RUC (RVM)"; Migrant 56:75, 1985: in the entry for Red-breasted Merganser change "23 May" to 23 Mar"; Migrant 56:76, 1985: in the entry for Bonaparte's Gull change "16 May" to "16 Mar" and in the entry for Ring-billed Gull at FRR change "11 May" to "11 Mar."

Addenda: Reports from this region were never submitted for the following seasons: Spring 80, Summer 80, Fall 80, Spring 82, and Summer 82. A number of important records occurred during those seasons and have subsequently come into the current compiler's hands. They are presented here, in the order of the 6th edition of the AOU Check-List, as a partial replacement for these missing reports. Some of the records are from the 1982 TOS Foray to White County, which covered both the Cumberland Plateau and Highland Rim portions of the county. Records from both physiographic regions are presented here. Should other important missing records be forthcoming, they, too, will be included in future addenda to this report.

American Bittern: 14 Jun 1980 (3) Goose Pond, Grundy County (Dan Jacobson, m.ob.), nesting suggested. Least Bittern: 14 Jun 1980 (2) Goose Pond, Grundy County (Dan Jacobson, m.ob.). Little Blue Heron: 1-16 Jun 1982 (2 ad) MC (DFV), Cattle Egret: 14 Apr 1980 (1) Old Hickory L. (LFK); 19 Apr 1982 (2) Madison, DVC (Richard Rimmer). Yellow-crowned Night-Heron: 28-31 May 1982 (2) Great Falls, WHC (TOS). WHITE IBIS: 1-2 Aug 1980 (2) Coleman L., DVC (LFK); 4-17 Aug 1980 (1) PPL (Audrey D. Perry). Blue-winged Teal: 25, 31 Jul 1980 (2 ad, 1 3/4 grown y) LWP (DTC, JPC), possible nesting indicated; 28 Jul 1982 (1) MC (DFV). Redhead: 24 May 1980 (1) Cedar Hill L., Madison, DVC (Richard Rimmer), new NA late spring date. Surf Scoter: 3 Apr 1980 (1 ad male) MC (DFV); from beginning of period until 27 Apr 1982 (1 imm male) RDL (CGD, m.ob.), new NA late spring date. MISSISSIPPI KITE: 22 Aug 1980 (1) Opryland, NASH (Margaret L. Mann), 4th NA record. Bald Eagle: 31 Mar 1982 (1) RDL (NTOS), few records at this locale; 8 May 1982 (1 imm) Gallatin (DTC, JPC, m.ob.). Cooper's Hawk: 28-31 May 1982 (2) WHC (TOS). Broadwinged Hawk: 24 Jun 1980 (2 ad, 2 y in nest) GSP (CPN, DTC, JPC), left nest 4 Jul 1980. Ruffed Grouse: 1 Jun 1980 (1) PIC (RCH); 28-31 May 1982 (1) Virgin Falls, WHC (TOS). Wild Turkey: 1 Jun 1980 (2) PIC (RCH). Sandhill Crane: 16 Mar 1980 (30) PIC (RCH); 18 Mar - 2 Apr 1980 (1) Dilton-Mankin Rd., RUC (ALH); 18 Mar 1982 (76) PIC (RCH); 31 May - 6 Jun 1982 (1) Curry Lane, NASH and Joelton, DVC (Joseph Martin, fide

Robert M. Hatcher), a sick or injured bird probably involved, as bird found dead last date. Spotted Sandpiper: 21 Jun 1980 (1 ad, 1 y) MC (DFV); 1 Jul 1982 (2 ad, 1 y) MC (DFV). MARBLED GODWIT: 26 Apr 1980 (2) MC (F. Joseph McLaughlin, m.ob.), 3rd NA record. Dunlin: 25 May 1980 (24) GSP (DTC, JPC), Buff-breasted Sandpiper: 6 Sep 1980 (21) CCNWR (Jeff Wilson, Joe Guinn), very large number. Common Snipe: 14 Jun 1980 (1) MC (DFV), few NA summer records, Wilson's Phalarope: 25 May 1980 (1) GSP (DTC, JPC); 6 Sep 1980 (1) CCNWR (Jeff Wilson, Joe Guinn). Monk Parakeet: 18 May 1980 (1, photos) Louisiana Ave., NASH (Dan Jacobson. m.ob.), probable escape. Ringed Turtle Dove: 8 Jun 1980 (2 ad, 1 y, nest with 3 eggs) Overton Rd., NASH (Dan Jacobson, m.ob.). Horned Lark: 28-31 May 1982 (3) WHC (TOS). Brown Creeper: 14 Jun 1980 (1) Love Circle, NASH (Jan C. Alexander); 17 Jul (1) Acklen Ave., NASH (F. Joseph McLaughlin); few NA summer records. Bewick's Wren: 24-25 Jun 1980 (1/1) Trousdale Co./Smith Co. (CPN); 8 Jul 1980 (1) 2 mi SE of Palmyra, MTC (David Snyder). Loggerhead Shrike: 28-31 May 1982 (15) WHC (TOS). Solitary Vireo: 28-31 May 1982 (1) WHC (TOS). Prothonotary Warbler: 18 May 1980 (pair nesting) PIC (RCH). Swainson's Warbler: 30 Apr 1980 (1 injured, recovered, banded) Austin Peav State Univ., MTC (David Snyder): 28-31 May 1982 (1) S of De Rossett, WHC (TOS). Black-throated Green Warbler: 28-31 May 1982 (5) WHC (TOS). Dickcissel: 28-31 May 1982 (5) WHC (TOS), Grasshopper Sparrow: 28-31 May 1982 (7) WHC (TOS), Song Sparrow: 28-31 May 1982 (120) WHC (TOS). Northern ("Baltimore") Oriole: May 1980 (singing male, nest) near Port Royal, MTC (Joe D. Allen): 28-31 May 1982 (1) WHC (TOS).

Locations: ACI - Ashland City, Cheatham Co.; BAH - Barnes Hollow, Putnam Co.; BAR - Barkley Waterfowl Management Unit, Stewart Co.; BAS - Basin Spring, Williamson Co.; BEC - Bear Creek Wildlife Management Unit, Stewart Co.; CCNWR - Cross Creeks National Wildlife Refuge, Stewart Co.; CCP Ponds - Carter's Creek Pike Ponds, Williamson Co.; CCSP - Cumberland City Steam Plant, Stewart Co.; CFC - Coffee Co.; CHC Cheatham Co.; CYC - Clay Co.; CLA - Clarksville, Montgomery Co.; DIC -Dickson Co.; DOV - Dover, Stewart Co.; DRU - Duck River Unit, Tennessee National Wildlife Refuge, Humphreys Co.; FRR - Franklin Reservoir, Williamson Co.; FTC - Fort Campbell Military Reservation, Montgomery and Stewart Cos.; FTD - Fort Donelson National Military Park, Stewart Co.; FTN - Fort Negley, Davidson Co.; GRP - Green's Pond, Sumner Co.; GSP -Gallatin Steam Plant, Sumner Co.; HIC - Hickman Co.; JAC - Jackson Co.; LBL - Land-Between-The-Lakes, Stewart Co.; LEF - Leiper's Fork, Williamson Co.; LOR - Loretto, Lawrence Co.; LWC - Lawrence Co.; LWP -Lewis Pond, Sumner Co.; MC - Metro Center, Davidson Co.; MOP -Monsanto Ponds, Maury Co.; MTC - Montgomery Co.; NA - Nashville Area (includes all of Cheatham, Davidson, Robertson, Rutherford, Sumner, Williamson, and Wilson Cos., as well as parts of Dickson and Trousdale Cos.); NASH - Nashville; OHL - Old Hickory Lake (in Davidson, Smith, Sumner, Trousdale, and Wilson Cos.); OVC - Overton Co.; PIC - Pickett Co.; PWP -Percy Warner Park, Davidson Co.; RDL - Radnor Lake State Natural Area, Davidson Co.; ROP - Robertson Pond, Lawrence Co.; RUC - Rutherford Co.; SUC - Sumner Co.; SWC - Stewart Co.; WHC - White Co.; WLC -Wilson Co.; WMC - Williamson Co.; WSB - Wiley's Spring Bay, Stewart Co.

STEPHEN J. STEDMAN, Rt. 6, Pinewood Rd., Franklin, TN 37064.

EASTERN RIDGE AND VALLEY REGION—The whole region is continuing to experience the effects of an extended drought. How this will affect the region's avifauna only time will tell. With this thought in mind, observers should watch very closely for the results and then report their impressions to the compilers. With our network of astute fieldworkers, comments will certainly be welcomed as we attempt to evaluate the effects of the weather.

In the Johnson City area, Glen Eller commented that during March, only 1.8 inches of rain was recorded, and in April, 0.9 inches. May rainfall was near normal, but it was very hot and humid. Knoxville experienced the driest spring on record, with 1.7 inches of rain in March, 2.3 inches in April, and 3.8 inches in May. Most of the May rainfall came during the last third of the month. March and May temperatures were near normal, while April was a few degrees below normal. The last frost at Knoxville was on 10 May, much later than average. At Chattanooga, the rainfall deficit measured 10 inches for the period and average temperatures were above normal during the period.

Waterfowl seemed to have moved through the region in normal fashion, as few comments were received as to their status.

The Blythe Ferry section of the Hiwassee River Area is rapidly gaining in importance for migrating Sandhill Cranes. Large numbers of cranes fed and roosted there in late February, and more birds were present in March. Before they departed, numbers peaked at 300 birds. Other sections of Hiwassee River Area, such as Armstrong Bend, Gunstocker, Candies, South Mouse and Rogers Creeks are probably equally utilized by cranes, but have not had the benefit of adequate documentation. For this reason, all observations should be reported. With the impinging human development all along the migration route, these locations will certainly gain in importance with each passing season.

Shorebirds are an interesting story. Under normal conditions, all of the TVA reservoirs are filled by the time the main spring migration has commenced. Thus, other than at Kingston Steam Plant, there is very limited available habitat. This season, mainstream reservoirs did not reach full summer pool levels until early June. While there was plenty of shoreline habitat available during the May migration period, most of the sites were too dry for the shorebirds. In late May, some birds were attracted to Savannah Bay mudflats and a few other areas when it rained.

There was a lack of strong frontal systems during the season, and, as a consequence, the passerine migration was unspectacular. Most of the birds arrived on schedule, with many of the later migrants not lingering as long as usual. As expected, individuals from last winter's northern finch invasion lingered into May, with some Pine Siskins still coming to a feeder on Signal Mountain at the end of the period.

Loon—Duck: Common Loon: 1-23 Mar (2-9) CHL (m.ob.); 16 Mar (2) FLL (KTOS); 1-2 Apr (2) FLL (PSP); 21 Apr (1) CHL (KHD); 13 May (1) BOL (RLK). Pied-billed Grebe: 1-16 May (4-37) CHL (m.ob.). Horned Grebe: 1-23 Mar (2-6) CHK (m.ob.). Double-crested Cormorant: 31 Mar (1) Maclellan Island, CHA (CDB); 12 Apr (8) FLL (PSP); 16 Apr (1) BOL (RLK); 26 Apr, 4 May (5) DOL (JAK); 10 May (2 flying N) KNX (MWA, LNA); 13 May (1) AUS (RLK). Great Egret: 1 Apr (1) CHL (KHD); 10, 24 Apr (1) AUS

(RLK); 31 May (1) POC (JCW). Snowy Egret: 16 Apr (1) BOL (RLK). Blackcrowned Night-Heron: 24 Apr (3) CHA (Randall L. Doyle, Linda Kay Doyle). Green-winged Teal: 3 May (300) HRA (KHD, LHD); 16 Apr (1) JNB (RLK), Am. Black Duck: 3 Mar (8) HRA (KHD, LHD), Mallard: 10 Mar (nest w/14 eggs) SAB (KHD). Northern Pintail: 26 May (1) HRA (KHD, LHD, RCS). Blue-winged Teal: 9-23 Mar (20-40) HRA (KHD, LHD); 4 May (2) KSP (KHD, LHD), Northern Shoveler, 4-12 May (1) KSP (KHD, LHD), Gadwall: 9 Mar (60) CHL (KHD); 15-28 Apr (1-10) SAB (KHD, LHD); 12 May (1) KSP (KHD, LHD). American Wigeon: 9-16 Mar (1-15) CHL (KHD, LHD); 2 Apr (1) HRA (KHD, LHD). Canvasback: 9 Mar (1) CHL (KHD, LHD). Redhead: 2-16 Mar (6-75) CHL (KHD, LHD). Ring-necked Duck: 2-23 Mar (1-211) CHL (KHD, LHD). Greater Scaup: 2-16 Mar (2-240) CHL (KHD, LHD). Lesser Scaup: 2-16 Mar (100-266) CHL (KHD, LHD); 4-12 May (4) KSP (KHD, LHD). Common Goldeneye: 2 Mar (2) CHL (KHD). Bufflehead: 4-23 Mar (4-16) CHL (KHD, LHD). Hooded Merganser: 9 Mar (31) HRA (KHD, LHD). Red-breasted Merganser: 16 Mar (28) CHL (KHD, LHD); 21 May (1) CHL, HLC (CPN). Ruddy Duck: 2-23 Mar (8-67) CHL (KHD, LHD).

Vulture—Merlin: Black Vulture: 30 Mar (nest, 2 eggs) Armstrong Bend, MEC (Paul H. Dietrich), same site as 1985 nesting. Osprey: 15 Mar (pair at nest) HRA, navigation light at Tenn. River Mile 501.3 (BJT); 11 Apr - 7 May (1+) BOL (LHTOS); 24 May (1) Catoosa WMA, CUC (JCR). Bald Eagle: 10 Mar (1) Watts Bar L. (T. Edward Beddow); 16 Mar (4) HRA (BJT); 20 Mar (1) Jefferson City (PSP); 21 Mar (1 imm) BOL (RLK). Sharpshinned Hawk: 15 May (1) SPW (GWM); 17 May (1) NOR (CPN) - both of these late migrants or breeding birds. Red-shouldered Hawk: total of 11 sites were active out of 17 new and old locations checked in Bradley, Meigs, Hamilton, Marion, McMinn, Rhea and Walker Co., Georgia (KHD, PGM, James D. Rowell, RAR). Broad-winged Hawk: 11 Apr (1) AUS (LHTOS). Red-tailed Hawk: active nest BYC (PGM, JDL); active nest Blythe Ferry, MEC (CWC). American Kestrel: nesting again White Pine, JEC (JAK); active nest w/4 young BYC (JDL, PGM). Merlin: 1 Apr (1) SPW (GWM); 3 May (1) KNX (ARH et al.).

Sora-Tern: Sora: 13, 21 Apr (2) AUS (RLK); 24 Apr (1) Alcoa Marsh (JAK). Sandhill Crane: 3-13 Mar (up to 300) HRA (CWC); 3 Apr (3) MEC (JCW); 16 May (1) S. Mouse Cr., HRA (Dan N. Williams). Black-bellied Plover: 23 May (1) KNC (MWA, LNA). Lesser Golden-Plover: 15 Mar (1) AUS (RLK); 17 Mar (2) SAB (LHD); 11 Apr (1) SAB (LHD, Rick Waldrop). Semipalmated Plover: 8 May (3) JNB (RLK); 19-31 May (1-4) SAB (LHD, KHD, RCS); 25 May (1) KSP (LHD, ACD, SLW, KHD). American Avocet: 26 Apr (4) Graysburg Hills Golf Course, GNC (Howard H. Earnest, Sr.); 28 Apr (25) Cherokee Lake (Mike Smith, fide LJF). Greater Yellowlegs: 15 Mar - 9 May (1+) AUS (RLK); 25 Apr (7) SAB (KHD). Lesser Yellowlegs: 18 Mar (1) SAB (LHD); 24 Apr (82) HRA (C. Richard Hughes); 28 Apr (28) SAB (LHD); 12 May (1) HRA (KHD, LHD). Willet: 24 Apr (27) SAB (KHD, LHD). Spotted Sandpiper: 24 May (1) BLC (JAK), lrs. Upland Sandpiper: 16 Apr (1), 20 Apr (2) KNC (MWA, LNA, RDH, ARH). Semipalmated Sandpiper: 5-16 Mar (1) SAB (KHD, LHD); 24 May (22) BLC (JAK); 25 May (100) KSP (LHD, ACD, SLW); 20-31 May (2-75) SAB (KHD, LHD, RCS). Western Sandpiper: 21 May (1) AUS (RC). Least Sandpiper: 19-22 May (8-9) SAB (KHD, LHD); 25 May (10) KSP (LHD, ACD, SLW, KHD). White-rumped Sandpiper: 25 May (10) KSP (LHD, ACD, SLW, KHD); 26 May (5) SAB (LHD, RCS, KHD). Baird's Sandpiper: 21 May (1) AUS (RC). Pectoral Sandpiper: 8-23 Mar (3-25) SAB (LHD); 15 Mar - 1 May (1-12) WGC (RLK et al.) - this represented a good migration for the area. Dunlin: 1-23 Mar (23-65) SAB (KHD, LHD); 24-25 May (1) KSP (MWA, LHD, ACD, SLW). Common Snipe: 1-23 Mar (6-57) SAB (KHD, LHD). American Woodcock: 3 Mar (6 singing) Amnicola Marsh, HLC (Paul C. Harris, R. John Henderson). Laughing Gull: 13 Mar (1 ad) Fort Loudoun Dam (JAK). Bonaparte's Gull: 9 Mar (2) CHL (KHD, LHD); 8 Apr (14) SAB (LHD); 16 Apr (1) BOL (RLK). Ring-billed Gull: 11 May (1) BOL (RLK); 22 May (9) CHL (CPN). Caspian Tern: 16 Apr (1) AUS (RLK), second ELI area spring record. Common Tern: 26 Apr (2) DOL (JAK). Forster's Tern: 24 Apr (7) FLL (PSP); 26 Apr (3) AUS (RLK, RC).

Barn-Owl-Nuthatch: Common Barn-Owl: bad news from White Pine, JEC. A nest tree used for over 20 years was cut, 2 young owls removed and given to TWRA; other know roost locations destroyed when a shed was torn down, and a tree limb blew down (JAK). Great Horned Owl: 24 Mar (active nest) Rogers Creek, McMinn Co. (KHD, LHD). Red-headed Woodpecker: 26 Mar (1) Summit, Hawkins Co. (Daniel J. Nieves): 16 Apr (1) JNC (RLK) - only reports. Olive-sided Flycatcher: 9 May (1) KNX (JCH, A. Boyd Sharp). Willow Flycatcher: 11 May (1) AUS (RLK), ers; 24 May (1) Roaring Fork Cr., GNC (WRN); 26 May (3) southern Fentress Co. (CPN). Least Flycatcher: 18 May (1) Cross Mountain, Campbell Co. (CPN). Horned Lark: 18 Apr (2 at nest) Grassy Cr., GNC (WRN); 23 May (1) UT Plant Sciences Farm, KNC (MWA), first breeding season report from this locality in some years; several on territory near Heritage High School, BLC (JAK). Tree Swallow: 1 Apr - 18 May (1+) WGC (LHTOS); 24 May (pair nesting in bluebird box) UT Plant Sciences Farm, KNC (MWA), first county nest record; 26 May (1) west Morgan Co. (CPN), probable nesting; pair again at DOL site (JAK). N. Rough-winged Swallow: 11 Mar (1) Maryville (JAK); 11 Mar (1) Raccoon Mountain, Marion Co. (LJF), ers. Cliff Swallow: 5 May (several at old nests) Goodfield Branch embayment, MEC (CPN); 6 May (8 new nests) TN 58 at Wolftever Cr., HLC (CPN) - first published reports from each location. Red-breasted Nuthatch: present in Knoxville area until at least 14 May (1) (JCH).

Gnatcatcher-Warbler: Blue-gray Gnatcatcher: 27 Mar (1) SPW (GWM). Veery: 30 May (1) KNX (JCH), late. Swainson's Thrush: 8 May (1) JNC (RLK) - only report from this area; 25 May (1) KNX (JCH), lrs. Hermit Thrush: 28 Apr (1) WGC (RLK), lrs. Water Pipit: 5-8 Mar (250-300) SAB (KHD, LHD): 26 Apr (90+) Tri-Cities Airport, Sullivan Co., (MD, Joy Kyle); 5 May (1) AUS (RLK); 11 May (4) KSP (MWS, LNA). Cedar Waxwing: 3 Mar (350) CHA (RAR); 16-23 May (10-20) Cravens House, Lookout Mtn. (KHD), feeding on ripening mulberries. Loggerhead Shrike: pair regular JNB (RLK et al.); 25 Apr (2 ad, 3 y) Hickory Valley Golf Course, CHA (RAR); 10 May (2 ad, 3 y) Snow Hill, HLC (KHD, LHD); 17 May (2 ad, 3 y) Davidson Rd., CHA (RAR). White-eyed Vireo: 5 Apr (1) KNX (RDH, ARH), ers. Solitary Vireo: 11 Mar (1) Raccoon Mt., Marion Co. (LJF), ers; 25 May (1) KNX (JCH), late report from non-breeding locale. Warbling Vireo: 28 Apr (1) SPW (GWM). Philadelphia Vireo: 13 May (1) JNC (SG). Red-eyed Vireo: 4 Apr (1) KNX (ARH, RDH), ers. Blue-winged Warbler: 7 Apr (1) SPW (GWM), ers; 5-31 May (1) CHA (RAR), first report of one singing on territory in HLC. "Brewster's" Warbler: 3 May (1) SPW (GWM); 4 May (1) KNX (James M. Campbell, Betty Reid Campbell, Frank L. Bills, RDH). Orange-crowned Warbler: 27 Apr (1) CHA (PDR). Black-throated Blue Warbler: 5 Apr (1) KNX (ARH), very early; 9 May (1), 13 May (2) SPW (GWM). Blackburnian Warbler: 5 Apr (1) KNX (ARH), early. Blackpoll Warbler: present in lower than average numbers in KNX area (CPN); 26 May (1) HRA (KHD, LHD, RCS), late. Black-and-white Warbler: 30 Mar (1) KNX (ARH, RDH), about average arrival. Worm-eating Warbler: 5 Apr (1) KNX (ARH, RDH), early. Ovenbird: 5 Apr (1) KNX (ARH, RDH), early. Connecticut Warbler: 24 May (1) BLC (JAK). Mourning Warbler: 22 May (1) KNX (JCH); 24 May (1) BLC (JAK). Hooded Warbler: 5 Apr (1) KNX (ARH, RCH), early. Wilson's Warbler: 8 May (1) KNX (CPN). Canada Warbler: 24 May (1) Catoosa WMA, CUC (JCR).

Grosbeak—Grosbeak: ROSE-BREASTED GROSBEAK: first 26 Apr (1) KNX (ARH, RDH, MWA, LNA), then migrants were present in low numbers: 11 May (pair building nest) Frozen Head, Morgan Co. (CPN); 23 May (2) HLC (KHD). Vesper Sparrow: 23 Mar (8 singing) HRA (KHD), LHD), Grasshopper Sparrow: 26 May (1) SE Putnam Co., (5) west Fentress Co., (1) west Morgan Co. (CPN). White-throated Sparrow: 20 May (1) KNX (James Mason), Irs. Bobolink: more reports than normal, from Knox, Anderson, Scott, Roane and Claiborne counties (MWA, ARH, LJF, CPN, GWM et al.). Purple Finch: 9 May (1) KNX (ARH); 23 May (1) JNC (MD), lrs. House Finch: several reports of young birds in JNC after mid-May (LHTOS); at CHA, winter flocks dispersed in March, and pair observed copulating on 24 Apr (RAR); 26 May (nest, 2 y and 3 eggs) SAB (KHD). Pine Siskin: 26 May (1) Oak Ridge (Wesley K. James); at feeder on Signal Mt. numbers ranged to 100+ at end of April, with lingering individuals noted to 31 May (JSL). Evening Grosbeak: 1 May (7) SPW (GWM); 4 May (3) KNX (KTOS); 10 May (1) JNC (SG); end of April numbers up to 100+ at feeder on Signal Mt., last date there 10 May (JSL).

Locations: AUS - Austin Springs, Washington Co.; BLC - Blount Co.; BOL - Boone Lake; BYC - Bradley Co.; CHA - Chattanooga; CHL - Chickamauga Lake; CUC - Cumberland Co.; DOL - Douglas Lake; FLL - Fort Loudoun Lake; GNC - Greene Co.; HLC - Hamilton Co.; HRA - Hiwassee River Area; JEC - Jefferson Co.; JNC - Johnson City; JNB - Jonesborough, Washington Co.; KSP - Kingston Steam Plant, Roane Co.; KNC - Knox Co.; KNX - Knoxville; MEC - Meigs Co.; NOR - Norris, Anderson Co.; POC -Polk Co.; SAB - Savannah Bay, Hamilton Co.; SPW - Speedwell, Claiborne Co.; WGC - Washington Co.

KENNETH H. AND LILLIAN H. DUBKE, 8139 Roy Lane, Ooltewah, TN 37363.

EASTERN MOUNTAIN REGION—The drought continued into this period, with monthly rainfall totals of 1.82 inches, 0.92 inches, and 4.88 inches for the months of March through May. The spring rainfall total was about 7 inches below normal. The most exciting find of the period was a Clapper Rail which was captured, held, and released in Elizabethton. If accepted by the TOS Certification Committee, this will be the first state record of this species.

Loon—Merganser: Common Loon: 18 Apr (1) LR (BHS, JCR), last report. Horned Grebe: 26 Apr (3, with 1 in breeding plumage) RNC (GDE, HHF), last report. Green-backed Heron: 13 Apr (1) ELI (HHF), first report. Yellow-crowned Night-Heron: first 27 Mar (1) ELI (GOW); 31 Mar (2 on nest) ELI (GDE). Canada Goose: brood of 5 y fledged at WGI (fide GDE).

Northern Pintail: 14 Mar (5) WTL (RC), last report. Blue-winged Teal: present in ELI area 21 Mar - 1 May (LHTOS). Northern Shoveler: 26 Apr (1) RNC (GDE, HHF); 21 May (3) WTL (MD, SG, CFW). Canvasback: 21 Mar (3) WTL (MD, SG, CFW). Lesser Scaup: 26 Apr (3) RNC (GDE, HHF), last report. Hooded Merganser: 17 Mar (2) RIL (GDE), last report. Redbreasted Merganser: 26 Apr (26) RNC (GDE, HHF).

Osprey-Swift: Osprey: present in ELI area 12 Apr - 7 May (LHTOS). Sharp-shinned Hawk: 8 May (1) TEL (RLK), only report since 12 Mar. Cooper's Hawk: 18 Apr (1) LR (BHS, JCR), last report. Red-shouldered Hawk: 24 May (1) GSMNP (RLK). Broad-winged Hawk: 13 Apr (2) RIL (GDE), first report. PEREGRINE FALCON: 3 young hacked in GSMNP during May (RLK et al.). Merlin: 9 Apr (1) LR (BHS, JCR). Wild Turkey: 26 May (1 ad with y) Shady Valley, Johnson Co. (Edward H. Schell). Sora Rail: 26 Apr (1) ERW (RPL, GWS). CLAPPER RAIL: 8 Apr (1 captured, released 17 Apr) ELI (GOW et al.), photographed, Round Table Note to be submitted. American Woodcock: 17 May (1) RNM, 5200 ft. elev. (RLK). Forster's Tern: 26 Apr (8) RNC (GDE, HHF); 4 May (4) WTL (RC). Blackbilled Cuckoo: 3 May (2) ELI (LHTOS), first report. Common Barn-Owl: 1-2 pairs at SHL area (fide GDE). Great Horned Owl: 2-3 pairs at SHL area (fide RPL). NORTHERN SAW-WHET OWL: 17 May (1) RNM (RLK). Chuck-will's-widow: 9 Apr (1) SHL (RPL), early. Chimney Swift: 7 Apr (3) ELI (HHF), first report.

Flycatcher-Vireo: Least Flycatcher: 3 May (5) RNM (LHTOS), first report. Alder Flycatcher: 17 May (6 singing males) RNM, elev. 5200+ ft. (RLK), Purple Martin: 9 Mar (2) ELI (fide GDE). Tree Swallow: 16 Mar (2) RNM, 5200 ft. elev. (RLK), first report. Common Raven: 20-26 May (2 ad, 4 y) GSMNP (RLK). House Wren: 13 Apr (1) ELI (HHF). Veery: first 3 May (4) RNM (LHTOS). Swainson's Thrush: 14 May (1) SHL (RPL), first report. Hermit Thrush: 17 May (1 singing) RNM, elev. 5200+ ft. (RLK). Water Pipit: 21 Mar (20+) WGI (SG). White-eyed Vireo: 10 Apr (1) SHL (RPL), first report, early. Yellow-throated Vireo: 6 Apr (1) SHL (RPL), first report, early. Warbling Vireo: 25 Apr (10 SHL (RPL).

Warbler-Grosbeak: Blue-winged Warbler: 17 May - EOP (1 territorial male) SHL (RPL). Golden-winged Warbler: 3 May (3) RNM (LHTOS), first report. Northern Parula: 15 Apr (1) RNC (SG), first report, early. Yellowrumped Warbler: 13 May (1) ELI (HHF), last report. Pine Warbler: first 14 Mar (1) WTL (RC). Louisiana Waterthrush: 3 Mar (1) TEL (PIK), very early. Mourning Warbler: 24 May (1) SHL (RPL). Blue Grosbeak: 26 Apr (1) ELI (GDE, HHF), first report. Grasshopper Sparrow: first 1 May (2) PWB (RC). Fox Sparrow: 21 Mar (1) ERW (GWS), last report. Bobolink: 7-13 May (7-11) PWB (RC). Purple Finch: last 26 Apr (2) ELI (GDE, HHF). House Finch: 10 May (nest, fledged 4 y) ELI (HHF). Red Crossbill: 14 Mar (1) WTL (RTC): 20 Apr (1) GSMNP (CPN et al.). Pine Siskin: 24-25 May (1) GSMNP (RLK); 2 May - EOP (3) ERW (GWS). Evening Grosbeak: 13 May (2) ELI (fide GDE).

Locations: ELI - Elizabethton; ERW - Erwin, Unicoi Co.; GSMNP -Great Smoky Mountains National Park; LOR - Look Rock, Blount Co.; PWB - Powder Branch, Carter Co.; RIL - Ripshin Lake, Carter Co.; RNC -Roans Creek area of Watauga Lake, Johnson Co.; SHL - South Holston Lake, Sullivan Co.; TEL - Tellico Ranger District, Cherokee Nat. Forest; WGI - Wagner's Island, Carter Co.; WTL - Watauga Lake.

GLEN D. ELLER, Rt. 9, Box 1340, Elizabethton, TN 37643.

OBSERVERS

LNA-Laurie N. Armstrong MWA-Marc W. Armstrong SDB-Sandra D. Bivens CDB—C. Del Blum DWB-Donald W. Blunk CHB-Carolyn H. Bullock CWC-Carl W. Campbell DCC-David C. Chaffin RC-Richard Clark BBC—Ben B. Coffey, Jr. LCC—Lula C. Coffey CKC-C. Kinian Cosner, Jr. DTC-Dot T. Crawford JPC-J. Paul Crawford ECC-E. Camille Crenshaw DAD-Dollyann Daily ACD-Angela C. Davis MD—Martha Dillenbeck HBD—Helen B. Dinkelspiel CGD-C. Gerald Drewry, Jr. KHD-Kenneth H. Dubke LHD-Lillian H. Dubke GDE-Glen D. Eller JOE-John O. Ellis HHF-Harry H. Farthing RPF-Robert P. Ford LJF-Linda J. Fowler SG—Sally Goodin KAG-Katherine A. Goodpasture MDH-Marguerite D. Hernandez AHH-Annie H. Heilman ALH-Anne L. Hettish ARH-Audrey R. Hoff RDH-Ron D. Hoff JCH-Joseph C. Howell MHI-Maxey H. Irwin DRJ-Daniel R. Jacobson PIK—Peter I. Kalla RLK-Richard L. Knight

JAK-Jon A. Koella LFK-Lee F. Kramer WNJ-William N. Jernigan RPL—Richard P. Lewis JDL-J. Dale Liner JSL-Jonnie Sue Lyons PGM-Paul G. Mascuch DMM—David M. McCarroll GWM-George W. McKinney RJM-Rocky J. Milburn, Jr. RVM-Ruth V. McMillan WRN-W. Ruth Nevius CPN-Charles P. Nicholson PSP-Paul S. Pardue WRP-William R. Peeples JCR-John C. Robinson PDR—Peter D. Robinson RAR-Robin A. Rudd RWS-Richard W. Simmers, Jr. DJS—Damien J. Simbeck BHS-Barbara H. Stedman SJS-Stephen J. Stedman RCS-Randy C. Stringer GWS-Glenn W. Swafford ATT-Ann T. Tarbell BJT-Bobby J. Thompson DFV-David F. Vogt MGW-Martha G. Waldron EJW-Ellen J. Walker GOW-Gary O. Wallace CFW-C. Frank Ward JCW-J. Craig Watson SLW-Susan L. Watson JRW-Jeff R. Wilson TJW-Terry J. Witt KTOS-Knoxville Chapter, TOS LHTOS-Lee R. Herndon Chapter, TOS MTOS—Memphis Chapter, TOS

PREPARATION OF COPY FOR PUBLICATION

The purpose of THE MIGRANT is the recording of observations and original information derived from the study of birds, primarily in the state of Tennessee or the area immediately adjacent to its borders. Articles for publication originate almost exclusively from T O.S. members.

Contributors should prepare manuscripts and submit them in a form acceptable to the printer, after editorial approval. Both articles and short notes are solicited but their format should be somewhat different.

Some suggestions to authors for the preparation of papers for publication are given herewith.

MATERIAL: The subject matter should relate to some phase of Tennessee Ornithology. It should be original, factual, concise, scientifically accurate, and not submitted for publication elsewhere.

TITLE: The title should be concise, specific, and descriptive.

STYLE: Recent issues of THE MIGRANT should be used as a guide in the preparation of manuscripts. Where more detail is needed reference should be made to the *Style Manual for Biological Journals* available from the American Institute of Biological Sciences, 1401 Wilson Boulevard, Arlington, Virginia 22209.

COPY: Manuscripts should be typed double spaced on 8½ x 11" paper with adequate margins, for editorial notations, and should contain only entries intended for setting in type, except the serial page number. Tabular data should be entered on separate sheets with appropriate title and column headings. Photographs intended for reproduction should be sharp with good contrast on glossy white paper in black and white (not in color). Instructions to the editors should be given on a separate sheet. Weights and measurements should be in metric units. Dating should be in "continental" form (e.g., 7 March 1981).

NOMENCLATURE: Common names should be capitalized followed by binomial scientific names in italics only after the first occurrence in the text for both regular articles and ROUND TABLE NOTES, and should conform to the A.O.U. Check-list 5th edition, 1957 and its Thirty-second Supplement. Trinomial should be used only after the specimen has been measured or compared with typical specimens.

BIBLIOGRAPHY: When there are more than five references in an article, they should be placed at the end of the article, otherwise they should be appropriately included in the text.

SUMMARY: Articles of five or more pages in length should be summarized briefly, drawing attention to the main conclusions resulting from the work performed.

IDENTIFICATION: Rare or unusual species identification to be acceptable must be accompanied by verifying evidence. This should include: date, time, light and weather conditions, exact location, habitat, optical equipment, distance, behavior of bird, comparison with other similar species, characteristic markings, experience of observer, other observers verifying observation and reference works consulted.

REPRINTS: Reprints are available on request. Reprint requests should accompany article at the time of submission. Billing to authors will be through the state T.O.S. Treasurer.

Books for review and articles for publication should be submitted to the editor. Seasonal reports and items should be forwarded to the appropriate departmental editor whose name and address will be found on the inside front cover.

CONTENTS

| THE 1986 SPRING FIELD DAYS. Richard L. Knight | 65 |
|---|----|
| ROUND TABLE NOTES | 70 |
| Second Tennessee Record of The Groove-billed Ani. William G. Chriswell | 70 |
| Nashville, Tennessee Television Tower Casualties, 1984. Katherine A. Goodpasture | 70 |
| Common Redpoll at Sewanee, Tennessee. James R. Peters | 72 |
| THE SEASON. Spring: 1 March—30 May 1986 | 73 |
| Western Coastal Plain Region. Martha G. Waldron | 74 |
| Highland Rim and Basin Region. Stephen J. Stedman | 75 |
| Eastern Ridge and Valley Region. Kenneth H. and Lillian H. Dubke | 83 |
| Eastern Mountain Region. Glen D. Eller | 86 |
| Observers | 88 |