

# ORNITOLOGIA NEOTROPICAL

EDITOR

ANDRÉ-ALEXANDER WELLER

## **INDEX TO VOLUME 21, 2010**

(Authors, key words, families, scientific names)

- Abundance, 599–610  
*Accipiter chionogaster*, 157–180  
*Accipiter erytronemius*, 263–270  
 Acosta Hospitaleche, C., 181–194  
*Actitis macularius*, 567–580  
*Agave salmiana*, 17–30  
*Agelaioides badius*, 1–16  
*Agelaius phoeniceus*, 17–30  
 Aguilar R., S. H., 87–103  
 Alarcón, P., 457–462  
 Albrieu, C., 283–294  
 Alexander Selkirk Island, 535–543  
 Alfaro, M., 361–370  
 Almeida, B. J. M., 567–580  
 Amat, J. A., 149–154  
*Amazilia beryllina*, 17–30, 519–533  
*Amazilia tzacatl*, 339–350  
*Amazilia violiceps*, 17–30  
*Amazona albifrons*, 599–610  
*Amazona autumnalis*, 599–610  
 American Kestrel, 263–270  
 Andes, 591–598, 611–614  
 Anggraini, K., 59–70  
 Anjos, L. dos, 203–213  
*Antilophia galeata*, 425–438  
*Anumbius anumbi*, 425–438  
*Aphrastura masafuerae*, 535–543  
 Apparent survival, 225–239  
*Aptenodytes forsteri*, 181–194  
*Aramides axillaris*, 109–129  
*Aratinga leucophthalma*, 425–438  
*Aratinga nana*, 599–610  
*Archilochus colubris*, 17–30  
*Ardea alba*, 383–396  
*Arenaria interpres*, 567–580  
 Arendt, W.-J., 409–423  
 Areta, J. I., 71–86, 351–359  
 Argentina, 1–16, 263–270, 283–294, 351–359, 445–451, 507–518  
*Arremon brunneinucha*, 339–350  
*Arremon flavirostris*, 425–438  
 Arthropods, 497–506  
 Association, 299–303  
 Atlantic forest, 71–86, 397–408, 425–438  
*Atlapetes albinucha*, 339–350  
*Angastes scutatus*, 425–438  
*Aulacorhynchus prasinus*, 339–350  
*Automolus leucophthalmus*, 425–438  
 Avalos, V. d. R., 195–202  
 Avian ecology, 519–533  
 Baja California Peninsula, 545–554  
 Bala, L.-O., 445–451  
 Bañados del Río Dulce, 383–396  
 Barn Swallow, 105–108  
*Basileuterus culicivorus*, 339–350, 425–438  
*Basileuterus flaveolus*, 425–438  
*Basileuterus hypoleucus*, 425–438  
*Basileuterus leucoblepharus*, 1–16, 425–438  
*Basileuterus rufifrons*, 339–350  
*Basileuterus tristriatus*, 339–350  
 Bay-ringed Tyrannulet, 397–408  
 Beautiful Jay, 611–614  
 Beltrán, J.-W., 409–423  
 Beltzer, A. -H., 1–16  
 Berduc, A. -J., 1–16  
 Berón, M. P., 215–224  
 Bertolin, M.-L., 457–462  
 Bianchi-Ballesteros, G., 305–309  
 Bianchi-Pérez, G., 305–309  
 Bioacoustic, 203–213  
 Bird communities, 339–350, 519–533  
 Bird migration, 105–108  
 Black-and-white Monjita, 299–303  
 Blackburn, G. A., 131–147  
 Bock, B. C., 271–281  
 Bodrati, A., 71–86, 351–359, 397–408  
 Bolivia, 39–46  
 Borges, F. J. A., 581–590  
 Bosque exótico, 1–16  
 Bosque nativo, 1–16  
 Botero-Delgadillo, E., 464–477  
 Botia-Becerra, J., 497–506  
 Brazil, 425–438, 453–456, 567–580  
 Breeding biology, 225–239, 361–370  
 Breeding diet, 457–462  
 Breeding period, 409–423  
 Breeding sites, 555–566  
 Breeding success, 439–443  
 Brood parasitism, 305–309  
 Brunetta, B., 203–213

- Bubo magellanicus*, 591–598  
*Bubulcus ibis*, 383–396  
 Bullock, S., 59–70  
*Burhinus bistriatus*, 149–154  
*Buteo albicaudatus*, 263–270  
*Buteo magnirostris*, 1–16, 263–270  
*Buteo swainsoni*, 263–270  
*Butorides striatus*, 383–396  
 Caballero-Sadi, D., 361–370  
*Cacicus solitarius*, 1–16  
 Calabuig, C. P., 555–566  
*Calidris alba*, 567–580  
*Calidris canutus rufa*, 445–451  
*Calidris fuscicollis*, 567–580  
*Calidris minutilla*, 545–554  
*Calidris pusilla*, 567–580  
*Calothorax lucifer*, 17–30  
*Campostoma obsoletum sclateri*, 109–129  
*Campostoma obsoletum*, 1–16  
*Campylopterus largipennis*, 425–438  
*Campylorhamphus trochilirostris*, 1–16  
*Campylorhynchus brunneicapillus*, 17–30  
*Campylorhynchus fasciatus*, 109–129  
*Campylorhynchus jocosus*, 17–30  
 Cangrejos, 215–224  
*Caprimulgus longirostris*, 425–438  
 Capture probability, 225–239  
*Caracara plancus*, 1–16, 263–270  
 Carmona, R., 545–554  
*Carpodacus mexicanus*, 17–30, 519–533  
 Casaux, R.-J., 457–462  
*Catharacta antarctica*, 181–194  
*Catharacta macormicki*, 181–194  
*Cathartes aura*, 263–270  
*Catharus aurantiirostris*, 339–350  
*Catharus frantzii*, 225–239  
*Catharus fuscescens*, 339–350  
*Catharus minimus*, 339–350  
*Catharus ustulatus*, 339–350  
*Cecropia pachystachya*, 59–70  
*Celeus obrieni*, 453–456  
 Central America, 225–239  
 Central Brazil, 581–590  
 Cerrado, 39–46, 581–590  
*Chaetocercus berlepschi*, 313–322  
*Chamaepetes unicolor*, 339–350  
 Chapman, C. A., 59–70  
*Charadrius collaris*, 567–580  
*Charadrius semipalmatus*, 567–580  
*Charadrius wilsonia*, 567–580  
 Chestnut-breasted Wren, 49–59  
 Chick mortality, 479–487  
 Chile, 535–543  
 Chimango Caracara, 263–270  
*Chionis alba*, 181–194  
*Chiroxiphia caudata*, 425–438  
*Chiroxiphia linearis*, 339–350  
*Chlorophonia callophrys*, 339–350  
*Chlorophonia cyanea*, 71–86  
*Chlorospingus ophthalmicus*, 339–350  
*Chlorostilbon aureoventris*, 1–16  
*Chlorostilbon lucidus*, 425–438  
 Christiansen, M. B., 59–70  
 Cinereous Harrier, 263–270  
*Circus buffoni*, 263–270  
*Circus cinereus*, 263–270  
 Cirignoli, S., 591–598  
 Climate change, 105–108  
 Cloud forest, 131–147, 225–239, 489–495  
 Coates-Estrada, R., 599–610  
*Coccyzus lansbergi*, 109–129  
*Coccyzus melacoryphus*, 1–16  
 Cockle, K., 397–408  
*Colaptes auratus*, 17–30  
*Colaptes melanochloros*, 1–16, 425–438  
*Colaptes rubiginosus*, 339–350  
*Colibri thalassinus*, 339–350  
 Colombia, 49–59, 497–506  
 Colonial waterbirds, 383–396  
 Coloniality, 479–487  
 Colonies, 271–281  
 Color-banding, 39–46  
 Columbidae, 581–590  
*Columbina cruziana*, 109–129  
*Columbina inca*, 17–30, 519–533  
*Columbina picui*, 1–16  
*Columbina squammata*, 581–590  
*Columbina talpacoti*, 581–590  
*Commiphora leptophloeos*, 59–70  
*Conopophaga lineata*, 425–438

- Conservation, 371–381, 464–477, 599–610  
*Contopus fumigatus*, 489–495  
*Contopus sordidulus*, 339–350  
*Contopus virens*, 339–350  
 Corvid, 203–213  
*Coryphospingus cucullatus*, 1–16  
*Corythopsis delalandi*, 425–438  
*Coscoroba coscoroba*, 181–194, 555–566  
 Costa Rica, 149–154, 339–350  
 Cotingidae, 195–202  
 Cracidae, 31–38  
*Cranioleuca pyrrhophia*, 1–16  
 Crustaceans, 457–462  
*Crypturellus parvirostris*, 425–438  
*Crypturellus soui*, 339–350  
 Curassow, 31–38  
*Cyanocompsa brissonii*, 1–16  
*Cyanocorax chrysops*, 203–213  
*Cyanocorax cyanomelas*, 39–46  
*Cyanocorax morio*, 339–350  
*Cyanocorax yncas*, 305–309  
*Cyanolyca cucullata*, 339–350  
*Cyanolyca pulchra*, 611–614  
*Cyclarhis gujanensis*, 1–16, 109–129  
*Cyananthus latirostris*, 17–30  
*Cyphorhinus thoracicus*, 49–59  
*Cypseloides fumigatus*, 351–359  
*Cypseloides major*, 351–359  
*Cypseloides rothschildi*, 351–359  
*Cypseloides senex*, 351–359  
*Cyrtograpsus angulatus*, 215–224  
*Dacnis cayana*, 425–438  
 Davanço, P. V., 251–261  
 De Labra, M. A., 599–610  
 De Oliveira, L. S., 251–261  
 DeGroot, L. W., 39–46  
 Demongin, L., 439–443  
*Dendroica coronata*, 17–30  
*Dendroica virens*, 339–350  
*Dendrotyx barbatus*, 87–103  
*Dendrotyx leucophrys*, 339–350  
 Diet, 215–224, 283–294, 445–451, 464–477,  
 497–506, 591–598  
*Diglossa baritula*, 17–30  
*Dilodendron bipinnatum*, 59–70  
*Diomedea epomophora*, 181–194  
*Diomedea melanophrys*, 181–194  
 Dispersal, 39–46  
 Distribution, 351–359  
*Dives narszeniczi*, 109–129  
 Dominino, J., 383–396  
 Dry forests, 59–70  
*Dryophila ferruginea*, 425–438  
*Dryophila ochropyga*, 425–438  
 Duca, C., 581–590  
 Dyrz, A., 489–495  
*Dysithamnus mentalis aequatorialis*, 109–129  
*Dysithamnus mentalis*, 425–438  
 Echeverry-Galvis, M.-A., 497–506  
 Ecology, 464–477, 489–495, 611–614  
 Egg, 489–495  
 Eggs, 109–129, 157–180, 581–590, 615–618  
*Elaenia frantzii*, 339–350  
*Elaenia mesoleuca*, 425–438  
*Elaenia obscura*, 425–438  
*Elaenia parvirostris*, 1–16  
*Elaenia* sp., 425–438  
*Elanoides forficatus*, 263–270  
*Elanus leucurus*, 263–270  
 Elliptic Fourier Analysis, 181–194  
 El Salvador, 157–180  
*Emberizoides ypiranganus*, 425–438  
*Embernagra longicauda*, 425–438  
*Empidonax flavescens*, 339–350  
*Empidonax flaviventris*, 339–350  
*Empidonax* sp., 17–30  
*Empidonax trailli*, 339–350  
 Endemic, 109–129  
 Ensembles de aves, 1–16  
 Environment, 181–194  
 Environmental variables, 105–108  
*Eremophila alpestris peregrina*, 497–506  
 Escalante, P., 599–610  
 Esmeraldas Woodstar, 313–322  
 Espinal, 1–16  
 Estades, C. F., 535–543  
*Eudiptes chrysocome*, 439–443  
*Eugenes fulgens*, 17–30  
*Euphonia cyanocephala aureata*, 71–86  
*Euphonia cyanocephala*, 71–86

- Euphonia elegantissima*, 71–86  
*Euphonia musica*, 71–86  
 European starling, 507–518  
*Euscarthmus meloryphus*, 1–16, 109–129  
 Exotic plantation, 339–350  
 Extinction, 31–38  
*Falco sparverius*, 263–270, 425–438  
 Falkland Islands, 439–443  
 Fandino, B., 1–16  
 Favero, M., 215–224  
 Fawn-breasted Tanager, 615–618  
 Feeding ecology, 59–70  
 Ferrari, S. F., 283–294, 567–580  
*Ficus calyptrocerus*, 59–70  
 Fiorini, V.-D., 507–518  
 Flight range, 371–381  
*Florisuga fusca*, 71–86  
 Flushing behavior, 149–154  
 Food, 157–180  
 Food availability, 283–294  
 Food items, 425–438  
 Food preference, 241–250  
 Foraging behavior, 464–477  
 Foraging, 497–506  
*Formicivora serrana*, 425–438  
 Fragmentation, 599–610  
 França, L. C., 581–590  
 Francisco, M. R., 251–261  
 Frugivory, 59–70  
 Fruit abundance, 409–423  
 Furnariidae, 535–543  
*Furnarius figulus*, 425–438  
*Furnarius rufus*, 1–16  
*Galbula ruficauda*, 425–438  
 Gantz, A., 241–250  
 García, G., 479–487  
 Gatto, A., 479–487  
 Gaviota de Olrog, 215–224  
 Genetic variability, 271–281  
*Geothlypis aequinoctialis*, 425–438  
*Geotrygon montana*, 425–438  
 Giant Cowbird, 305–309  
*Glaucidium costaricanum*, 339–350  
 Gómez-Aíza, L., 17–30  
 Gonçalves, R. M. M., 425–438  
 Gonzalez, C., 271–281  
 González, F.-R., 591–598  
 Grau, E. T., 31–38  
 Green, A. J., 555–566  
 Green Jay, 305–309  
 Greeney, H. F., 313–322, 489–495  
 Gressler, D. T., 581–590  
 Guanacaste, 149–154  
*Gnarea guidonia*, 59–70  
*Gnibourtia hymenaeifolia*, 59–70  
*Gnira gnira*, 1–16  
 Habitat, 299–303  
 Habitat occupancy, 203–213  
 Habitat use, 599–610  
*Haplospiza unicolor*, 425–438  
 Harris, J. B. C., 313–322  
 Harris's Hawk, 263–270  
 Heming, N. M., 581–590  
*Hemitriccus diops*, 425–438  
*Hemitriccus margaritaceiventer*, 1–16  
*Hemitriccus nidipendulus*, 425–438  
*Henicorbina leucophrys*, 339–350  
 Hernández, M.-A., 445–451  
 Hickman, B. R., 313–322  
 High altitude grasslands, 591–598  
 Hillman, G., 383–396  
*Hilocybta mustelina*, 339–350  
*Himantopus melanurus*, 181–194  
*Hirundinea ferruginea*, 425–438  
*Hirundo rustica*, 105–108  
 Hodum, P. J., 535–543  
 Home range, 409–423  
 Honduras, 157–180  
 Horned Lark, 497–506  
 Hudsonian Godwit, 283–294  
 Hybrids, 31–38  
*Hydropsalis torquata*, 425–438  
*Hylocharis leucotis*, 17–30  
*Hylophilus amaurocephalus*, 425–438  
 Ibañez, L., 181–194  
*Icterus pustulatus*, 17–30  
*Icterus abeillei*, 519–533  
*Icterus bullockii*, 17–30  
*Icterus cayanensis*, 1–16  
*Icterus galbula*, 17–30

- Icterus graceannae*, 109–129  
*Icterus parisorum*, 17–30  
*Icterus spurius*, 17–30  
*Icterus wagleri*, 17–30  
 Incubation, 313–322  
 Incubation period, 109–129  
 Invasive species, 507–518  
 Island conservation, 535–543  
 Jahn, A. E., 39–46  
 Jayat, J.-P., 591–598  
 Jenner, T., 157–180  
 Jiménez, S., 361–370  
 Juan Fernández archipelago, 535–543  
 Juhant, M. A., 263–270  
 Juiña, M. E., 313–322  
 Juvenile plumage, 453–456  
 Juveniles, 313–322  
 Kaempfer's Woodpecker, 453–456  
 Kattan, G.-H., 49–59  
 Kirwan, G. M., 397–408  
*Knipolegus nigerrimus*, 425–438  
 Knowlton, J. L., 109–129  
 Komar, O., 225–239  
 Krüger, L., 299–303  
*Lampornis calolaemus*, 339–350  
*Lampornis clemenciae*, 17–30  
*Lanius ludovicianus*, 17–30  
 Laporta, P., 361–370  
*Larus atlanticus*, 215–224  
*Larus dominicanus*, 361–370, 479–487  
*Larus maculipennis*, 181–194  
*Lathrotriccus euleri*, 1–16, 425–438  
 Laufenberg, D., 611–614  
 Ledezma, J. P., 39–46  
 Leite, G. A., 453–456  
 Leite, L. O., 425–438  
 Lenzi, J., 361–370  
 León, J., 383–396  
*Lepidocolaptes affinis*, 339–350  
*Lepidocolaptes angustirostris*, 1–16  
*Leptopogon amarocephalus*, 425–438  
*Leptotila verreauxi*, 1–16, 339–350  
*Leptotila verreauxi decolor*, 109–129  
 Levey, D. J., 39–46  
 Lima, C.-A., 425–438  
*Limosa haemastica*, 283–294  
 Lined Seedeater, 251–261  
 Lizarralde, Z., 283–294  
*Lochmias nematura*, 425–438  
 Logistic regression, 299–303  
 Longitudinal migration system, 71–86  
 Long-winged Harrier, 263–270  
 Lopera, J. C., 271–281  
 Lopes, L. E., 581–590  
 Macedo, R. H., 615–618  
 MacGregor-Fors, I., 105–108, 519–533  
 Machalilla National Park, 109–129  
*Macronectes giganteus*, 181–194  
 Macuiltépetl Ecological Park, 87–103  
 Maglianesi, M.-A., 339–350  
 Mamani, A. M., 39–46  
*Manacus manacus*, 425–438  
 Mangeaud, A., 383–396  
 Manica, L. T., 581–590  
 Mar Chiquita, 383–396  
 Marcelino, D. G., 453–456  
*Margarops fuscatus*, 409–423  
 Marini, M. Â., 581–590  
 Martin, T. E., 131–147  
*Megascops choliba*, 425–438  
*Melanerpes aurifrons*, 17–30  
*Melanerpes cactorum*, 1–16  
*Melanerpes hypopolius*, 17–30  
*Melanopareia elegns*, 109–129  
*Melanotis caerulescens*, 17–30  
*Melospiza fusca*, 519–533  
*Melospiza leucotis*, 339–350  
 Menegheti, J. O., 555–566  
 Mexico, 87–103  
 Mexico City, 519–533  
 Michelutti, P., 383–396  
*Micrastur ruficollis*, 339–350  
 Migration, 371–381, 545–554, 567–580  
 Migration route, 555–566  
 Migration watchsite, 263–270  
*Milvago chimango*, 263–270  
*Mimus longicaudatus*, 109–129  
*Mimus polyglottos*, 17–30  
*Mionectes olivaceus*, 339–350  
*Mionectes rufiventris*, 425–438

- Miranda, M. V., 323–337  
 Misiones province, 351–359  
 Mist nets, 131–147, 323–337  
 Mistle-toe, 71–86  
*Mitu mitu*, 31–38  
*Mitu tuberosum*, 31–38  
*Mniotilta varia*, 339–350  
*Molothrus aeneus*, 17–30  
*Molothrus ater*, 17–30  
*Molothrus bonariensis*, 1–16  
*Molothrus oryzivora*, 305–309  
*Molothrus rufoaxillaris*, 1–16  
 Molt, 157–180, 225–239  
 Molting sites, 555–566  
*Momotus momota*, 339–350  
 Monte, 591–598  
 Monterrubio Rico, T. C., 599–610  
 Mountain savanna, 195–202  
 mtDNA, 271–281  
 Muriel, R. A., 555–566  
*Muscipipra vetula*, 71–86  
 Musmeci, R.-L., 445–451  
*Myadestes melanops*, 339–350  
*Myiarchus crinitus*, 339–350  
*Myiarchus swainsoni*, 1–16  
*Myiarchus tuberculifer*, 339–350  
*Myiobius barbatu*s, 425–438  
*Myioborus miniatus*, 339–350  
*Myiodynastes maculatus*, 1–16, 425–438  
*Myiopagis viridicata*, 425–438  
*Myiophobus fasciatus*, 1–16, 109–129  
*Myiopsitta monachus*, 1–16  
*Myiozetetes cayanensis*, 425–438  
*Myrmia micrura*, 109–129  
*Myrmorchilus strigilatus*, 425–438  
 Nacimiento, J. X. do, 371–381  
 Nearctic shorebirds, 567–580  
*Nectandra cissiflora*, 59–70  
*Neobelice granulata*, 215–224  
*Neopelma pallescens*, 425–438  
 Neotropics, 519–533  
 Nest, 157–180, 489–495, 615–618  
 Nest construction, 313–322  
 Nest construction behavior, 489–495  
 Nest crypsis, 149–154  
 Nest density, 157–180  
 Nest site selection, 195–202  
 Nest sites, 351–359  
 Nesting, 397–408  
 Nesting behaviour, 251–261  
 Nesting ecology, 149–154  
 Nesting success, 383–396, 195–202  
 Nestling, 489–495, 611–614, 615–618  
 Nestling period, 109–129  
 Nests, 109–129, 581–590  
*Nonnula rubecula*, 425–438  
 Northwestern Argentina, 323–337, 591–598  
 Null model, 299–303  
*Numenius phaeops*, 567–580  
*Nyctibius griseus*, 425–438  
*Nycticorax nycticorax*, 383–396  
*Nyctidromus albicollis*, 425–438  
 Oilbirds, 271–281  
 Ortega-Álvarez, R., 519–533  
 Ortiz, P.-E., 591–598  
 Otero, L.-D., 305–309  
*Pachyramphus albogriseus*, 109–129  
*Pachyramphus polychopterus*, 1–16  
*Pachyramphus validus*, 425–438  
*Pachyramphus viridis*, 1–16  
 Páez, C.-A., 464–477  
 Pagot, M., 383–396  
 Paiva, L. V., 581–590  
 Palkachupa, 195–202  
*Parabuteo unicinctus*, 263–270  
 Pardiñas, U.-F.-J., 591–598  
 Parental care, 251–261, 489–495, 611–614  
*Parula pitiayumi*, 1–16  
*Passer domesticus*, 17–30, 519–533  
 Pastures, 241–250  
*Patagioenas cayennensis*, 581–590  
*Patagioenas picazuro*, 581–590  
*Patagioenas maculosa*, 1–16  
*Patagioenas picazuro*, 1–16  
 Patagonia, 283–294, 457–462, 479–487  
 Patino, J., 555–566  
 Pearly-eyed Thrasher, 409–423  
 Pearman, M., 351–359  
 Península Valdés, 445–451  
 Pereira, A. I., 149–154

- Pereira, S. L., 31–38  
 Peres, C. A., 59–70  
 Petry, M. V., 299–303  
*Phacellodomus erythrophthalmus*, 425–438  
*Phacellodomus ruber*, 1–16  
*Phacellodomus rufifrons*, 425–438  
*Phaeomyias murina*, 425–438  
*Phaeomyias tumbezana*, 109–129  
*Phaethornis eurynome*, 425–438  
*Phaethornis idaliae*, 425–438  
*Phaethornis pretrei*, 425–438  
*Phaethornis ruber*, 425–438  
*Phaethornis squalidus*, 425–438  
*Phalacrocorax atriceps*, 457–462  
*Phalacrocorax brasilianus*, 383–396  
 Phenology, 555–566  
*Pheucticus chrysogaster*, 109–129  
*Pheucticus melanocephalus*, 17–30, 519–533  
*Phibalura flavirostris*, 71–86  
*Phibalura flavirostris boliviana*, 195–202  
*Philydor rufum*, 425–438  
*Phoenicopterus chilensis*, 181–194  
*Phoradendron*, 71–86  
*Phylloscartes sylvicolus*, 397–408  
*Phylloscartes ventralis*, 425–438  
*Picoides scalaris*, 17–30  
*Picumnus albosquamatus*, 425–438  
 Piedmont forest, 323–337  
 Pinheiro, R. T., 453–456  
*Pipilo fuscus*, 17–30  
*Pipraeidea melanonota*, 615–618  
*Piranga ludoviciana*, 17–30  
*Pitangus sulphuratus*, 1–16, 425–438  
 Pittalunga, S., 283–294  
*Platypsaris homochrous*, 109–129  
*Platyrinchus mystaceus*, 425–438  
 Plencovich, G., 383–396  
 Plush-crested Jay,  
*Pluvialis squatarola*, 567–580  
*Pogonotriccus*, 397–408  
 Point counts, 131–147  
 Poisbleau, M., 439–443  
*Polioptila caerulea*, 17–30  
*Polioptila dumicola*, 1–16  
*Polioptila plumbea*, 109–129  
 Politi, N., 323–337  
*Poospiza hispaniolensis*, 109–129  
*Poospiza melanolenca*, 1–16  
 Population density, 49–59  
 Population structure, 271–281  
 Population trend, 225–239  
 Porro, G., 457–462  
 Post-breeding period, 409–423  
 Pozzi, C., 383–396  
 Predation, 361–370  
 Predation, 479–487  
 Prey selection, 283–294  
*Protium heptaphyllum*, 59–70  
*Psaltiriparus minimus*, 519–533  
*Pseudoseisura lophotes*, 1–16  
 Psittacidae, 599–610  
 Puna, 591–598  
 Punta Rasa, 263–270  
 Purplish Jay, 39–46  
*Pygoscelis adeliae*, 181–194  
*Pygoscelis antarctica*, 181–194  
*Pygoscelis papua*, 181–194  
*Pyriglena leucoptera*, 425–438  
*Pyrocephalus rubinus*, 109–129  
*Pyrrhura viridicata*, 464–477  
 Quillfeldt, P., 439–443  
*Quiscalus mexicanus*, 17–30  
 Radio telemetry, 39–46  
 Ragusa-Netto, J., 59–70  
 Rain forest, 409–423  
 Ramoni-Perazzi, P., 305–309  
 Ramphastidae, 59–70  
*Ramphastos toco*, 59–70  
 RAPD, 271–281  
 Raptor migration, 263–270  
 Rebolo Ifrán, N., 507–518  
 Red Knots, 445–451  
*Regulus calendula*, 17–30  
 Reintroduction, 31–38  
 Reproductive biology, 251–261  
 Reproductive stages, 409–423  
*Rhodospingus cruentus*, 109–129  
 Ring recovery, 555–566  
 Río Gallegos Estuary, 283–294  
 Rivera, L. O., 323–337



- Roadside Hawk, 263–270  
 Roesler, I., 351–359  
 Rodríguez, A., 383–396  
*Rostrhamus sociabilis*, 263–270  
 Ruelas I., E., 87–103  
 Rufous-thighed Hawk, 263–270  
 Saffron-cowled Cowbird, 299–303  
*Sakesphorus bernardi bernardi*, 109–129  
 Salt gland furrow, 181–194  
*Saltator aurantiirostris*, 1–16  
*Saltator coerulescens*, 1–16  
*Saltator similis*, 1–16, 425–438  
*Saltator striatipectus*, 109–129  
 Santos, P. J. P. dos, 371–381  
 Sauma, L., 545–554  
 Sáyago, R., 105–108  
*Schefflera morototoni*, 59–70  
*Schiffornis virescens*, 425–438  
*Schistochlamys ruficapillus*, 425–438  
*Sclerurus scansor*, 425–438  
 Sea birds, 181–194  
 Seasonal abundance, 567–580  
 Seasonality, 39–46  
*Selasphorus platycercus*, 17–30  
 Sergipe, 567–580  
*Serpophaga subcristata*, 1–16  
 Sex and age rates, 545–554  
*Sicalis flaveola*, 1–16  
 Silva, L., 479–487  
 Simcox, W., 131–147  
 Siqueira, P.-R., 425–438  
*Sittasomus griseicapillus*, 425–438  
 Snail Kite, 263–270  
 Sooty Swift, 351–359  
 Soto-Werschitz, I.-A., 305–309  
 Sousa, L. M. S., 251–261  
 South America, 251–261, 555–566  
 Southern Caracara, 263–270  
 Southern Chile, 241–250  
 Spatial distribution, 445–451  
 Species loss, 599–610  
*Spheniscus magellanicus*, 181–194  
*Spinus psaltria*, 17–30, 519–533  
*Spizella atrogularis*, 17–30  
*Sporophila lineola*, 251–261, 425–438  
*Sporophila peruviana*, 109–129  
 Starling density estimate, 507–518  
*Steatornis caripensis*, 271–281  
*Stephanophorus diadematus*, 425–438  
*Sterculia striata*, 59–70  
*Sterna hirundo*, 181–194  
 Stopover ecology, 283–294  
 Strange, I. J., 439–443  
*Streptoprocne biscutata*, 351–359, 425–438  
*Streptoprocne zonaris*, 351–359  
 Strewé, R., 295–298  
*Sturnus vulgaris*, 17–30, 507–518  
 Swainson's Hawk, 263–270  
 Swallow-tailed Cotinga, 195–202  
 Swallow-tailed Kite, 263–270  
*Synallaxis albescens*, 1–16  
*Synallaxis cinerascens*, 425–438  
*Synallaxis frontalis*, 1–16  
*Synallaxis ruficapilla*, 425–438  
*Synallaxis stictothorax stictothorax*, 109–129  
*Syndactyla rufosuperciliata*, 425–438  
*Tachyeres patachonicus*, 181–194  
*Tachyphonus coronatus*, 425–438  
 Tambussi, C., 181–194  
*Tangara cayana*, 425–438  
*Tapera naevia*, 1–16  
*Taraba major*, 1–16  
 Tartara, M.-A., 457–462  
 Taylor, J., 225–239  
 Tello, S.-A., 49–59  
 Temperature, 105–108  
*Tersina viridis*, 425–438  
 Teta, P., 591–598  
*Thalasseus maximus*, 361–370, 479–487  
*Thalasseus sandvicensis eurygnathus*, 361–370, 479–487  
*Thalurania glaucopis*, 425–438  
*Thamnophilus caerulescens*, 1–16, 425–438  
*Thamnophilus pelzelni*, 425–438  
*Thamnophilus ruficapillus*, 425–438  
*Theristicus melanopis*, 241–250  
 Thermal environment, 149–154  
*Thraupis sayaca*, 1–16  
 Threatened populations, 457–462  
*Thryomanes bewickii*, 17–30, 519–533

- Thryothorus sclateri*, 109–129  
 Toco Toucan, 59–70  
*Todirostrum poliocephalum*, 425–438  
*Tolmomyias sulphurescens*, 109–129, 425–438  
 Tomasevic, J. A., 535–543  
 Torres, R., 383–396  
*Toxostoma curvirostre*, 17–30  
*Trichothraupis melanops*, 425–438  
*Tringa flavipes*, 567–580  
*Tringa melanoleuca*, 567–580  
*Tringa semipalmata*, 567–580  
*Troglodytes aedon*, 1–16  
*Trogon surrucura*, 425–438  
 Trophic guilds, 425–438  
 Tropical evergreen forest, 599–610  
 Tropical Kingbird, 39–46  
*Turdus albicollis*, 425–438  
*Turdus amaurochalinus*, 1–16  
*Turdus flavipes*, 71–86  
*Turdus leucomelas*, 425–438  
*Turdus migratorius*, 519–533  
*Turdus reevei*, 109–129  
*Turdus rufiventris*, 1–16, 425–438  
*Turdus rufopalliatus*, 519–533  
 Turkey Vulture, 263–270  
*Tyrannus melancholicus*, 39–46  
*Tyrannus vociferans*, 17–30  
 Understory birds, 323–337  
 Urban avifauna, 87–103  
 Urban ecology, 519–533  
 Urbanization, 519–533  
*Vanellus chilensis*, 181–194, 567–580  
 Vasconcelos, M.-F., 425–438  
 Venezuela, 615–618  
 Venezuelan Andes, 305–309  
*Veniliornis mixtus*, 1–16  
 Veracruz, 87–103  
 Verhelst, J.-C., 464–477  
*Vermivora celata*, 17–30  
*Vireo olivaceus*, 1–16, 109–129  
 Vocalizations, 157–180  
 Wajntal, A., 31–38  
 Water level, 383–396  
 Weather conditions, 439–443  
 Western Brazil, 59–70  
 White-breasted Hawk, 157–180  
 White-rumped Sandpiper, 371–381  
 White-tailed Hawk, 263–270  
 White-tailed Kite, 263–270  
 Wind, 383–396  
 Woodward, C., 611–614  
 Wunderle, J.-M. Jr., 409–423  
*Xanthocephalus flavus*, 299–303  
*Xiphorhynchus fuscus*, 425–438  
*Xolmis dominicanus*, 299–303  
 Yorio, P., 479–487  
 Yungas, 323–337  
*Zenaida asiatica*, 17–30  
*Zenaida auriculata*, 1–16  
*Zenaida meloda*, 109–129  
*Zonotrichia capensis*, 1–16, 425–438  
 Zuria, I., 17–30

## REVIEWERS FOR ORNITOLOGÍA NEOTROPICAL, VOLUME 20

Each manuscript submitted for publication in *Ornitología Neotropical* is generally reviewed by two specialists, often including a member of the editorial board, and frequently by a third one. The following individuals kindly reviewed one or more (\*) manuscripts for Volume 21: J. L. B. Albuquerque, J. A. Amat, M. J. Andersen, L. d. Anjos\*, J. I. Areta\*, E. Barbieri, J. Bates, N. Bayly, L. Bellis, I. Berkunsky, C. Bianchi, K. Bildstein, J. Blake\*, D. Blamires, D. Brighsmith, M. d. L. Brooke, L. Bugoni, G. S. Cabanne, D. Calderon, P. Capllonch, A. Castillo, O. Chassot, M. Codesido, R. Covacado, V. Cueto, R. Curry, M. P. Dantas Santos, K. Delhey, A. Di Giacomo, S. Duijns, R. Durães, M. Efe, K. Eisermann, J. C. Eitniecear\*, M. Erwin, B. Fandiño, G. Fernandez, K. Fierro-Calderón, R. Fraga\*, M. Francisco, R. Franke, A. Gatto, H. Gomez de Silva, F. González-García, J. A. González-Oreja\*, J. A. Graves, F. Hayes\*, A. B. Hennessey, G. Hernandez Volpato, F. Hertel, T. Hetmanski, E. Höfling, J. T. Ibarra, J. P. Isaach\*, A. Jahn, O. Jahn, L. Joseph, J. Karubian, P. Kennedy, G. M. Kirwan, N. Krabbe, P. Lau, L. Leveau\*,

F. Mallet-Rodriguez, M. Marin, P. McGowan, R. McNeil\*, S. McGehee, I. McGregor-Fors, E. Mellink\*, L. B. Mendonça, E. Miller, T. Monterrubio, A. Morales, M. C. Morera, G. Murga, G. M. Nachtigall, J. L. Nascimento, A. Navarro Siguenza\*, E. Nol, D. Oehler, F. Olmos, E. Palacios, V. Piacentini, G. Pohland, R. Prum, J. Ramos, M. Raposo\*, J. Rappole\*, J. Rau, C. Rengifo C., A. Rinderknecht, W. D. Robinson\*, O. Rojas Soto, G. Rompré, J. J. Roper, E. Ruelas Inzunza\*, M. Russello, I. Ruvalcaba-Ortega\*, L. Sánchez-González, M. A. Santos Alves, A. Sapoznikow, R. Schlatter, A. Schmitz\*, K.-L. Schuchmann\*, L. F. Silveira, E. Stolen, J. M. Thiollay, L. Tomialojc, A. Trejo, D. Tubelis\*, I. D. Valencia, M. F. Vasconcelos\*, J. H. Vega-Rivera\*, C. Vereá, B. Villa-Bonilla, F. Villela\*, A.-A. Weller\*, J. Wiley\*, M. Yasue, P. Yorio\*.

Some of the abstracts and/or manuscripts were kindly corrected by J. C. Eitniecear and J. Rappole for English, by E. Ruelas I. for Spanish, and by L. d. Anjos, M. Raposo, and M. F. Vasconcelos for Portuguese. Page-setting was done by R. McNeil and A.-A. Weller.

