

of shorebird biology and their use in coastal wetland assessment and conservation should keep the front of research moving significantly.

ACKNOWLEDGMENTS

I thank many colleagues and friends for significant help in this venture. Those I can remember and have to mention especially include George J. Divoky and the Pacific Seabird Group for creating the occasion for a shorebird symposium and for their warm support throughout; L. Richard Mewaldt, local program chairman for the 1977 meeting at Asilomar, and his staff for generous and efficient help in planning and running of the meeting itself; M. R. Foster, T. R. Howell, J. R. Jehl, Jr., and O. P. Pearson for advice on Latin American activities in shorebird work; C. Hand, J. D. Cubit, J. W. Hedgpeth, and W. J. North for help with background information useful for my introductory paper in particular as well as the symposium in general; Peter C. Lent and the BLM-NOAA offices in Boulder, Colorado, for travel support of speakers from Great Britain; Daniel W. Anderson and the PSG's publication committee for counsel on planning the symposium for publication; Ralph J. Raitt, editor for *Studies in Avian Biology*, for guidance and assistance in readying manuscripts for the printer; J. P. Myers for listening to me and giving useful reactions at all stages of this symposium; and last but definitely not least, the Fish and Wildlife Service and its Deputy Associate Director, Harvey K. Nelson, for support of the Pacific Seabird Group through a grant contributing significantly to the cost of publication.

Frank A. Pitelka

September, 1978