

CONTENTS

AUTHOR ADDRESSES	xvii
ABSTRACT	1
TAXONOMY	5
GENETIC EVIDENCE FOR KNOT SUBSPECIES	7
ECOLOGICAL AND MORPHOLOGICAL EVIDENCE FOR SUBSPECIES IN NORTH AMERICA	9
EVIDENCE FOR DISCRETE WINTERING POPULATIONS	9
PHYSICAL DESCRIPTION	11
FIRST BASIC PLUMAGE	11
DEFINITIVE ALTERNATE PLUMAGE	13
ALTERNATE PLUMAGE	13
FEMALES	13
MALES— <i>CALIDRIS CANUTUS RUFUS</i>	13
MALES— <i>CALIDRIS CANUTUS CANUTUS</i>	13
MALES— <i>CALIDRIS CANUTUS ROGERSI</i>	13
MALES— <i>CALIDRIS CANUTUS ISLANDICA</i>	14
MALES— <i>CALIDRIS CANUTUS ROSELAARI</i>	14
FIRST ALTERNATE PLUMAGE	14
JUVENILE PLUMAGE	14
HATCHLINGS	14
DISTRIBUTION IN TIME AND SPACE	14
THE ANNUAL CYCLE	14
BREEDING RANGE	15
WINTER (NON-BREEDING) RANGE	17
MIGRATORY RANGE AND MAJOR STOPOVER AREAS	19
BIOLOGY AND NATURAL HISTORY	25
REPRODUCTION	25
<i>Mating displays</i>	26
<i>Sexual behavior</i>	26
<i>Nest sites</i>	27

<i>Number of broods</i>	27
<i>Clutch size</i>	27
<i>Incubation period</i>	27
<i>Nestling period</i>	27
<i>Nesting success</i>	28
MORTALITY	28
LONGEVITY	28
SITE FIDELITY	28
MIGRATION	28
FEEDING HABITS	29
HABITAT	30
PREFERRED MICROHABITATS	30
BREEDING HABITAT	30
MIGRATION AND STOPOVER HABITAT – CANADA	30
MIGRATION AND STOPOVER HABITAT – NORTHEASTERN U.S.	30
MIGRATION AND STOPOVER HABITAT – DELAWARE BAY, U.S.	30
<i>Habitats important for Red Knots in Delaware Bay</i>	33
<i>Red Knot feeding ecology in Delaware Bay</i>	36
<i>Mapping horseshoe crab spawning habitat suitability</i>	39
<i>Mapping critical Red Knot habitat</i>	41
<i>Evidence of decline in both the population of horseshoe crabs and the availability of their eggs for Red Knots</i>	44
MIGRATION AND STOPOVER HABITAT – SOUTHEASTERN UNITED STATES	47
MIGRATION AND STOPOVER HABITAT – PANAMA	48
MIGRATION AND STOPOVER HABITAT – MARANHÃO, BRAZIL	48
MIGRATION AND STOPOVER HABITAT – LAGOA DO PEIXE, BRAZIL	48
MIGRATION AND STOPOVER HABITAT – PAMPAS REGION, ARGENTINA	49
MIGRATION AND STOPOVER HABITAT – PATAGONIAN SHORELINE, ARGENTINA	50
MIGRATION AND STOPOVER HABITAT – URUGUAY	50
WINTER HABITAT – UNITED STATES	50

WINTER HABITAT — CHILE	50
WINTER HABITAT — ARGENTINA	52
WINTER HABITAT — BRAZIL	52
WINTER HABITAT — PANAMA	52
POPULATION SIZE AND TRENDS	52
RED KNOT POPULATIONS OF THE AMERICAS	53
WINTERING POPULATION TRENDS IN <i>CALIDRIS CANUTUS RUFA</i>	56
PASSAGE POPULATION TRENDS	58
BREEDING AREA POPULATION TRENDS	62
SUMMARY OF POPULATION TRENDS	63
<i>Population size and trends of Calidris canutus roselaari</i>	64
GEOGRAPHIC AREA SUMMARIES	65
<i>Chile</i>	65
<i>Argentina</i>	65
<i>Brazil</i>	70
<i>Panama</i>	73
<i>United States Fish and Wildlife Service Region 2 — Texas</i>	73
<i>United States Fish and Wildlife Service Region 4 — Florida</i>	74
<i>United States Fish and Wildlife Service Region 4 — Georgia</i>	75
<i>United States Fish and Wildlife Service Region 4 — South Carolina</i>	77
<i>United States Fish and Wildlife Service Region 4 — North Carolina</i>	79
<i>United States Fish and Wildlife Service Region 5 — Virginia</i>	82
<i>United States Fish and Wildlife Service Region 5 — Maryland</i>	85
<i>United States Fish and Wildlife Service Region 5 — New Jersey and Delaware</i>	85
<i>United States Fish and Wildlife Service Region 5 — New York</i>	89
<i>United States Fish and Wildlife Service Region 5 — Connecticut</i>	89
<i>United States Fish and Wildlife Service Region 5 — Rhode Island</i>	89
<i>United States Fish and Wildlife Service Region 5 — Massachusetts</i>	90
<i>United States Fish and Wildlife Service Region 5 — New Hampshire</i>	91
<i>United States Fish and Wildlife Service Region 5 — Maine</i>	91

<i>Canada</i>	93
THREATS	95
THREATS TO HABITATS IN DELAWARE BAY DURING SPRING MIGRATION	95
THREATS TO HABITAT IN MASSACHUSETTS	97
THREATS TO HABITAT IN NORTH CAROLINA	97
THREATS TO HABITAT IN SOUTH CAROLINA	97
THREATS TO HABITAT IN FLORIDA	97
THREATS TO HABITAT IN BRAZIL	97
THREATS TO HABITAT AT MIGRATION STOPOVER SITES ALONG THE ATLANTIC COAST OF PATAGONIA, ARGENTINA	98
THREAT OF OIL POLLUTION AND POSSIBILITY OF OTHER UNIDENTIFIED FACTORS AFFECTING THE PRINCIPAL C. C. RUFA NON-BREEDING SITE AT BAHÍA LOMAS, CHILE	98
OIL POLLUTION THREAT AND HUMAN DISTURBANCE AT THE ONLY OTHER MAJOR NON-BREEDING SITE AT RÍO GRANDE, ARGENTINA	98
THREATS TO RED KNOT HABITAT IN CANADA	99
OVER UTILIZATION FOR COMMERCIAL, RECREATIONAL, SCIENTIFIC, OR EDUCATIONAL PURPOSES	99
HAVE SCIENTIFIC STUDIES CONTRIBUTED TO THE RED KNOT’S DECLINE?	99
DO SCIENTIFIC STUDIES CAUSE SIGNIFICANT DISTURBANCE TO RED KNOTS	100
STEPS TO MINIMIZE DISTURBANCE BY RESEARCH ACTIVITIES	100
DISEASE OR PREDATION	100
THE INADEQUACY OF EXISTING REGULATORY MECHANISMS	102
INADEQUACIES OF THE FEDERAL AND REGIONAL REGULATORY SYSTEM	102
INADEQUACIES OF REGULATORY SYSTEMS IN INDIVIDUAL STATES	104
CURTAILMENT OF HABITAT USE FROM DISTURBANCE BY PEOPLE AND DOGS	105
COMPETITION FROM GULLS	107
RISKS ASSOCIATED WITH SMALL POPULATION SIZE	108
WEATHER-RELATED THREATS TO RED KNOTS	108
SUMMARY OF LAND OWNERSHIP AND EXISTING HABITAT PROTECTION FOR POPULATIONS	108

PAST AND CURRENT CONSERVATION AND HABITAT MANAGEMENT	
ACTIVITIES UNDERTAKEN TO BENEFIT THE SPECIES	108
THE RAMSAR CONVENTION ON WETLANDS	108
WESTERN HEMISPHERE SHOREBIRD RESERVE NETWORK (WHSRN)	108
IMPORTANT BIRD AREAS PROGRAM	109
CONVENTION ON THE CONSERVATION OF MIGRATORY SPECIES OF WILD ANIMALS 1979	109
NATIONAL WILDLIFE REFUGES	109
THE INTERNATIONAL SHOREBIRD SURVEY AND PROGRAM FOR REGIONAL AND	
INTERNATIONAL SHOREBIRD MONITORING	109
DELAWARE BAY—INCREASED AVAILABILITY OF HORSESHOE CRAB EGGS	110
DELAWARE BAY—PROTECTION OF ROOSTING SITES	110
DELAWARE BAY—REDUCE DISTURBANCE BY MINIMIZING RESEARCH ACTIVITIES	111
DELAWARE BAY—MONITOR NUMBERS OF MIGRATORY SHOREBIRDS ON THE	
DELAWARE BAY STOPOVER	111
DELAWARE BAY—PAST AND CURRENT MANAGEMENT ACTIONS FOR SHOREBIRD	
POPULATIONS	112
DELAWARE BAY—PAST AND CURRENT MANAGEMENT ACTIONS FOR THE HORSESHOE	
CRAB POPULATIONS	113
DELAWARE BAY—MANAGEMENT PLANS	114
DELAWARE BAY—HABITAT PROTECTION	114
DELAWARE BAY—BAIT BAGS	114
NON-BREEDING AND STOPOVER AREA MANAGEMENT AND CONSERVATION	114
<i>South America</i>	114
<i>United States – Florida</i>	117
<i>United States – Georgia</i>	117
<i>United States – South Carolina</i>	117
STOPOVER HABITAT MANAGEMENT	117
<i>United States – North Carolina</i>	117
<i>United States – Virginia</i>	117
<i>United States – Maryland</i>	118

<i>United States – New Jersey</i>	118
<i>United States – Delaware</i>	119
<i>United States – New York</i>	120
<i>United States – Connecticut</i>	120
<i>United States – Rhode Island</i>	120
<i>United States – Massachusetts</i>	120
<i>United States – New Hampshire</i>	121
<i>United States – Maine</i>	121
<i>Panama</i>	121
<i>Canada</i>	121
BREEDING HABITAT MANAGEMENT	121
OTHER MANAGEMENT CONSIDERATIONS AND OPPORTUNITIES	121
MONITORING EFFECTS AND MANAGEMENT ACTIVITIES	122
CONSERVATION GOALS AND THE SURVEYS, MONITORING, RESEARCH, AND MANAGEMENT NEEDED TO SUPPORT THEM	123
SURVEY NEEDS	124
<i>South America – overall</i>	124
<i>South America – Argentina</i>	124
<i>South America – Chile</i>	124
<i>South America – Brazil</i>	125
<i>Caribbean countries and northern South America</i>	125
<i>Mexico</i>	125
<i>United States – Delaware Bay</i>	125
<i>United States – Virginia</i>	125
<i>United States – North Carolina, South Carolina, Georgia, and Florida</i>	125
<i>United States – Alaska</i>	125
<i>United States – California</i>	125
<i>United States – Washington</i>	125
MONITORING NEEDS	126
<i>Overall</i>	126

<i>South America – Argentina</i>	126
<i>South America – Chile</i>	126
<i>South America – Brazil</i>	126
<i>United States – Delaware Bay</i>	127
<i>United States – Virginia</i>	127
<i>United States – North Carolina and South Carolina</i>	127
<i>United States – Florida and Georgia</i>	127
<i>United States – other Sites on the United States East Coast</i>	127
<i>Arctic – Canada and Alaska</i>	127
RESEARCH NEEDS	128
<i>Broad-scale research topics</i>	128
<i>Country-specific research needs – Argentina</i>	129
<i>Country-specific research needs – Chile</i>	130
<i>Country-specific research needs – Brazil</i>	130
<i>Country-specific research needs – Mexico</i>	130
<i>Country-specific research needs – United States</i>	130
MANAGEMENT NEEDS	131
UPDATE TO THE STATUS OF THE RED KNOT (<i>CALIDRIS CANUTUS</i>) IN	
THE WESTERN HEMISPHERE, FEBRUARY 2008	131
TAXONOMIC STATUS	132
POPULATION STATUS OF <i>CALIDRIS CANUTUS ROSELAARI</i>	132
<i>Summary</i>	133
POPULATION STATUS OF <i>CALIDRIS CANUTUS RUFUS</i>	133
<i>Tierra del Fuego population</i>	133
<i>Maranhão population</i>	134
<i>Florida population</i>	134
<i>Are the counts accurate?</i>	134
<i>Could the birds have moved elsewhere?</i>	134
MASS GAIN IN DELAWARE BAY	135
HORSESHOE CRABS AND THEIR EGGS IN DELAWARE BAY	135

RECOMMENDATIONS	136
ACKNOWLEDGMENTS	137
LITERATURE CITED	138

TABLES

TABLE 1. NUMBER OF CITATIONS OF THE MOST EXTENSIVELY STUDIED SHOREBIRDS IN THE WORLD (THOMAS ET AL. 2003).....	5
TABLE 2. POPULATION ESTIMATES OF THE SIX SUBSPECIES OF THE RED KNOT (<i>CALIDRIS CANUTUS</i>)	6
TABLE 3. ESTIMATES OF F_{ST} FOR POPULATION DIFFERENTIATION IN RED KNOTS (BELOW DIAGONAL) CALCULATED USING mtDNA CONTROL REGION SEQUENCES	7
TABLE 4. MEAN WING CHORD AND CULMEN MEASUREMENTS FROM MUSEUM SPECIMENS OF RED KNOTS TAKEN FROM WESTERN HEMISPHERE LOCATIONS	12
TABLE 5. BODY MASS OF WESTERN HEMISPHERE RED KNOTS AT DIFFERENT STAGES OF NORTH AND SOUTH MIGRATION	12
TABLE 6. HABITAT TYPES UTILIZED BY FORAGING RED KNOTS ON BREEDING GROUNDS (B), SPRING MIGRATION (S), FALL MIGRATION (F), AND WINTERING GROUNDS (W)	31
TABLE 7. THE AMOUNT OF TIME (HR) NEEDED TO ACHIEVE THE DAILY HORSESHOE CRAB EGG CONSUMPTION OF 24,000 FRESH EGGS (HARAMIS ET AL. 2007) IN RELATION TO EGG AVAILABILITY (EGGS M^{-2}) DEPENDENT ON WHETHER EGGS ARE: (A) FREELY AVAILABLE ON THE SAND SURFACE OR (B) BURIED WITHIN THE TOP 5 CM OF SAND	38
TABLE 8. CHARACTERIZATION AND LENGTHS OF THE DELAWARE BAY SHORELINE	39
TABLE 9. LENGTH OF SHORELINE IN DELAWARE AND NEW JERSEY ACCORDING TO SUITABILITY FOR HORSESHOE CRAB SPAWNING	40
TABLE 10. LENGTH AND PERCENTAGE OF EACH SPAWNING HABITAT SUITABILITY CATEGORY IN CONSERVATION OWNERSHIP	42
TABLE 11. TOTAL NUMBERS OF ALL BENTHIC INVERTEBRATES COLLECTED DURING TRANSECT SAMPLING ON METOMPKIN (MET), PARRAMORE (PARR), AND FISHERMAN ISLANDS (FISH), VIRGINIA, IN MAY 2000 (TRUITT ET AL. 2001)	48
TABLE 12. NUMBERS OF EACH INVERTEBRATE SPECIES COUNTED DURING TRANSECT SAMPLING ON METOMPKIN, PARRAMORE, AND FISHERMAN ISLANDS, VIRGINIA IN MAY 2000 (TRUITT ET AL. 2001)	49
TABLE 13. INVERTEBRATES RECORDED IN TRANSECT SAMPLING AT BAHÍA LOMAS, CHILE, AND THE RELATIVE ABUNDANCE OF EACH	52
TABLE 14. RECENT POPULATION ESTIMATES OF RED KNOTS WINTERING IN THE NEW WORLD	56
TABLE 15. WINTERING SITES OF RED KNOTS IN ARGENTINA (LOCATIONS GIVEN WITH PROVINCES IN PARENTHESES)	68
TABLE 16. STOPOVER SITES USED BY RED KNOTS IN ARGENTINA DURING NORTHWARD MIGRATION	69
TABLE 17. TOWNS AND PROVINCES IN WHICH RED KNOTS HAVE BEEN OBSERVED IN ARGENTINA	70
TABLE 18. NUMBER OF RED KNOTS COUNTED ON THE NORTH COAST OF MARANHÃO, BRAXIL, IN APRIL AND MAY 1995 (I. SERRANO ET AL., UNPUBL. DATA)	71

TABLE 19. NUMBER OF RED KNOTS RECORDED ON THE COAST OF MARANHÃO, BRAZIL	71
TABLE 20. COUNTS OF RED KNOTS AT THREE SITES IN PANAMA BETWEEN 5 JANUARY AND 15 APRIL 2002 (BUEHLER 2002)	73
TABLE 21. RECORDS OF RED KNOTS ON THE TEXAS COAST DURING 1980–1996 (SKAGEN ET AL.1999)	74
TABLE 22. LOWEST AND HIGHEST NUMBER OF RED KNOTS COUNTED ON THE COAST OF FLORIDA IN FOUR COUNTS BETWEEN 16 DECEMBER 1993 AND 1 MARCH 1994, ORDERED ACCORDING TO THE MAXIMUM NUMBER RECORDED	75
TABLE 23. IMPORTANT RED KNOT STOPOVER AND WINTER LOCATIONS IN GEORGIA	76
TABLE 24. RED KNOT SURVEYS CONDUCTED BETWEEN 2000 AND 2004 IN THE CAPE ROMAIN REGION OF CHARLESTON, SOUTH CAROLINA	78
TABLE 25. STOPOVER AND POTENTIAL WINTERING AREAS FOR RED KNOTS IN NORTH CAROLINA	81
TABLE 26. NUMBER OF RED KNOTS OBSERVED DURING THE 2001–2002 INTERNATIONAL SHOREBIRD SURVEY ON CLAM SHOAL, NORTH CAROLINA	82
TABLE 27. RESULTS OF AERIAL SURVEYS CONDUCTED AT LOW TIDE FOR RED KNOTS ALONG OUTER BEACH SURF ZONE OF VIRGINIA BARRIER ISLANDS FROM VIRGINIA–MARYLAND BORDER TO MOUTH OF CHESAPEAKE BAY IN 1995, 1996, AND 2005 (B. WATTS, PERS. COMM.; B. TRUITT, PERS. COMM.)	82
TABLE 28. RESULTS OF AERIAL SURVEYS CONDUCTED AT LOW TIDE FOR RED KNOTS ALONG OUTER BEACH SURF ZONE OF VIRGINIA BARRIER ISLANDS FROM VIRGINIA–MARYLAND BORDER TO MOUTH OF CHESAPEAKE BAY IN 1995 (B. WATTS, PERS. COMM.; B. TRUITT, PERS. COMM.)	83
TABLE 29. RESULTS OF AERIAL SURVEYS CONDUCTED AT LOW TIDE FOR RED KNOTS ALONG OUTER BEACH SURF ZONE OF VIRGINIA BARRIER ISLANDS FROM VIRGINIA–MARYLAND BORDER TO MOUTH OF CHESAPEAKE BAY IN 1996 (B. WATTS, PERS. COMM.; B. TRUITT, PERS. COMM.)	83
TABLE 30. GROUND SURVEYS FOR RED KNOTS ON CHINCOTEAGUE NATIONAL WILDLIFE REFUGE, 1992–2003	84
TABLE 31. RESULTS OF SHOREBIRD SURVEYS CONDUCTED ON FISHERMAN ISLAND NATIONAL WILDLIFE REFUGE, VIRGINIA, FOLLOWING INTERNATIONAL SHOREBIRD SURVEY PROTOCOL IN 2004	85
TABLE 32. PEAK COUNTS OF RED KNOTS OBSERVED IN AERIAL SURVEYS OF DELAWARE BAY SHORELINE	86
TABLE 33. NUMBERS OF RED KNOTS ROOSTING AT HIGH WATER BY DAY AND AT DUSK AT STONE HARBOR POINT, NEW JERSEY, DURING MAY 2001 (H. P. SITTERS, UNPUBL. DATA)	87
TABLE 34. OBSERVATIONS OF RED KNOTS FEEDING IN THE STONE HARBOR AREA WETLANDS DURING MAY 2001 (H. P. SITTERS, UNPUBL. DATA)	88
TABLE 35. SUMMARY OF RED KNOT EVENING AND NIGHT COUNTS AND NUMBER OF RADIO- TAGGED RED KNOTS AT HEREFORD INLET, STONE HARBOR, NEW JERSEY, DURING 19–31 MAY 2005 (H. P. SITTERS, UNPUBL. DATA)	88
TABLE 36. SUMMARY OF SPRING AND FALL MIGRATION SURVEYS FOR RED KNOTS IN RHODE ISLAND BETWEEN 1982 AND 2004 (C. RAITHEL, PERS. COMM.)	90
TABLE 37. RED KNOT SURVEY DATA FOR MAINE (1980–2004)	94
TABLE 38. AERIAL SURVEY COUNTS OF GULLS ON THE ATLANTIC COAST OF NEW JERSEY (D. JENKINS, UNPUBL. DATA)	107

TABLE 39. COUNTS OF RED KNOTS DURING THE NORTHERN WINTERS OF 2004–2005 TO 2007–2008 IN TIERRA DEL FUEGO (ARGENTINA AND CHILE), MARANHÃO (BRAZIL), AND ON THE WEST COAST OF FLORIDA (NC = NO COUNT). WHERE NO COUNT OCCURRED, THE TOTALS ROW USES THE PREVIOUS YEARS’ COUNT FOR MARANHÃO AND THE SUCCEEDING YEAR’S COUNT FOR FLORIDA (SEE TEXT)	133
TABLE 40. POPULATION PARAMETERS OF HORSESHOE CRABS IN DELAWARE BAY FOR 2004–2007	136
TABLE 41. DENSITY OF HORSESHOE CRAB EGGS IN THE TOP 5 CM OF SAND IN THE BEACHES OF DELAWARE BAY DURING MAY AND JUNE 2004–2007 IN NEW JERSEY AND DELAWARE (SURVEYS CONDUCTED RESPECTIVELY BY THE NEW JERSEY AND DELAWARE DIVISIONS OF FISH AND WILDLIFE)	136

FIGURES

FIGURE 1. Worldwide distribution of the six recognized subspecies of the Red Knot. All breeding areas (dark gray shading) are on high-arctic tundra where the adults spend June–July. After their long-distance migrations (arrows), they spend the non-breeding season (August–May) mainly in intertidal soft-sediment habitats (dots, which are scaled according to population size). This map was prepared in 2003 and revised according to recent studies described in this review. Note that it is uncertain whether the Red Knots that winter in Northern Brazil and/or Florida are <i>Calidris canutus roselaari</i> , but some birds presumed to be <i>C. c. roselaari</i> winter on the coast of California and Baja California (map drawn by Dick Visser, provided by Jan van Gils, and reproduced with their permission)	6
FIGURE 2. Minimum spanning network showing the relationships between haplotypes from the mitochondrial control region of Red Knots. Ovals represent haplotypes and connecting lines represent a single base pair change between haplotypes. Small open circles on lines represent multiple base pair changes between haplotypes	7
FIGURE 3. Observed and expected mismatch distributions of mitochondrial control region sequences in Red Knots. Red Knots closely match the pattern expected under population growth in the recent past	8
FIGURE 4. Biometrical variation of male and female Red Knots in the wintering populations of Maranhão (Brazil), Florida, and Bahia Lomas (Tierra del Fuego) indicated by (A) bill length and (B) body mass (mean \pm 95% confidence intervals (data are from Baker et al. 2005a and Niles et al. 2006)	10
FIGURE 5. Diagrammatic representation of the annual cycle of a typical Tierra del Fuego wintering Red Knot (<i>Calidris canutus rufa</i>) in terms of latitudinal location and date. Horizontal lines represent periods when birds stay on the breeding or wintering grounds or stopover while on migration; dotted lines represent largely non-stop migratory flights	
FIGURE 6. Predicted Red Knot nesting habitats based on land cover types in the Canadian Arctic and point locations of Red Knots obtained by radio telemetry (Red Knot data from New Jersey Department of Environmental Protection Endangered and Nongame Species Program; potential Red Knot habitat data from Grant F. Walton Center for Remote Sensing and Spatial Analysis (CRSSA) Rutgers University; boundary data from GeoCratis Canada)	15
FIGURE 7. Red Knot wintering areas in the Western Hemisphere. Each area boxed in the left map is shown in greater detail and delineated in black	16

FIGURE 8. International Shorebird Survey Data (ISS) showing distribution of Red Knots in winter in the U.S. before year 2000 (upper) and during 2000–2004 (lower). The level of ISS survey effort declined after 2000; therefore, the differences in numbers before and since 2000 may partly represent reduced survey effort (B. Harrington, pers. comm.)	18
FIGURE 9. International Shorebird Survey (ISS) Data showing distribution of Red Knots during fall migration in the U.S. before year 2000 (upper) and during 2000–2004 (lower). The level of ISS survey effort declined after 2000; therefore, the differences in numbers before and since 2000 may partly represent reduced survey effort (B. Harrington, pers. comm.)	20
FIGURE 10. Critical stopover sites used by Red Knots during northward and southward migration in South America	21
FIGURE 11. International Shorebird Survey (ISS) Data showing distribution of Red Knots during spring migration in the U.S. before 2000 (upper) and during 2000–2004 (lower). The level of ISS survey effort declined after 2000; therefore, the differences in numbers before and since 2000 may partly represent reduced survey effort (B. Harrington, pers. comm.)	22
FIGURE 12. Critical breeding, migration stopover, and wintering habitat for the Red Knot <i>Calidris canutus rufa</i> . Numbers on the map correspond with the numbers in Table 6	23
FIGURE 13. Annual landings of horseshoe crabs in Virginia, Maryland, Pennsylvania, Delaware, New Jersey, and New York, 1990–2006 (Morrison et al. 2004). Most states had mandatory reporting by 1996 and all did by 1998, so landings data prior to 1998 may be underrepresented	25
FIGURE 14. Migration route of <i>Calidris canutus rufa</i> between its wintering grounds on Tierra del Fuego, South America, stopover areas along the Patagonian Coast of Argentina, and in the northeastern United States, and breeding grounds in the Canadian Arctic	29
FIGURE 15. Map of the Delaware Bay (New Jersey and Delaware) showing some of the most important refueling sites for Red Knots	34
FIGURE 16. Map of horseshoe crab spawning habitat suitability with location of protected conservation lands. Several key locations have been annotated: (A) Slaughter Beach, (B) Cape May National Wildlife Refuge, (C) Fortescue, and, (D) Broadkill Beach. Protected Lands GIS Data Sources: NJDEP, NJ Green Acres, TNC-NJ Chapter, DE Parks and Recreation	35
FIGURE 17. Functional responses relating the intake rate (eggs s ⁻¹) achieved by Red Knots to the density of (upper) eggs present on sand surface and (lower) eggs buried and mixed in the top 5 cm of sand	37
FIGURE 18. Density of eggs on the sand surface or buried and mixed in the sediment (down to 5 cm) will determine whether it is most profitable to peck or probe	38
FIGURE 19. Map of horseshoe crab spawning habitat suitability on Delaware Bay based on beach sediment and development characteristics (Lathrop and Allen 2005). Note that this mapping does not include consideration of beach morphology or wave energy characteristics that may be also be important in determining the suitability of the beach as horseshoe crab spawning habitat or other human disturbance or habitat factors that might influence bird usage	41

FIGURE 20. Map of percent Red Knot use for spring stop over between 1986–1990 and 2001–2005. Survey data summed across the 5-yr period and percent of total calculated for each beach segment (K. Clark, unpubl. data)	43
FIGURE 21. Number of horseshoe crabs in 30-foot trawls in Delaware Bay during May 1990–2006 (S. Michels, pers. comm.). The declining trend is highly significant ($r^2 = 0.65$, $P < 0.001$)	45
FIGURE 22. Density of horseshoe crabs eggs in the upper 5 cm of sand in the Delaware Bay beaches of New Jersey during late May 1985–2006. The declining trend is highly significant ($r^2 = 0.56$, $P = 0.002$). Source: 1985–1999 (M. L. Botton, pers. comm.; R. E. Loveland, pers. comm.); 2000–2006 (NJENSP, unpubl. data). Confidence intervals are not plotted because the raw data are not available for the earlier period and for the later period they are very small in relation to the scale. All data points relate to 2–6 sampling dates spread over May and early June and to core samples taken along transects between the high and low tide lines at 3-m intervals	45
FIGURE 23. Mean of weekly aerial counts of Red Knots in New Jersey and Delaware in May 2002–2005	46
FIGURE 24. Mean densities of horseshoe crab eggs in the upper 5 cm of sand from beach transects sampled once in late May and once in early June at six sites on the Delaware shore of Delaware Bay during 2002–2004 ordered from north (Port Mahon) to south (Slaughter Beach) (Weber 2003, 2004). At each site on each sampling date, 20 core samples were taken along each of two transects covering 83% of the distance between the nocturnal high tide line and the tidal flat. Only the means for both transects are given by Weber so confidence intervals are not available	47
FIGURE 25. Location of Bahía Lomas in Tierra del Fuego, Chile	51
FIGURE 26. The intertidal distribution pattern of <i>Darina solenoides</i> at Bahía Lomas, Chile	52
FIGURE 27. Nitrogen and carbon assimilated by Red Knots from ingested <i>Darina solenoides</i> (C. Espoz, unpubl. data)	53
FIGURE 28. Tierra del Fuego and Bahía Lomas, Chile, the primary wintering grounds of <i>Calidris canutus rufa</i>	54
FIGURE 29. The number of Red Knots counted at Cape Romain National Wildlife Refuge, South Carolina, 2000–2004 (Cape Romain National Wildlife Refuge, South Carolina DNR, unpubl. data)	55
FIGURE 30. The number of Red Knots spending the austral summer in southern South America according to aerial counts made during the <i>Atlas of Nearctic shorebirds on the coast of South America</i> project (Morrison and Ross 1989) in 1985 and during 2000–2006. Grey sections are numbers at Bahía Lomas, black sections are other sites in Tierra del Fuego (mainly Río Grande) and southern Chilean Patagonia and white sections are other sites farther north along the coast of Argentina. No counts were made north of Tierra del Fuego in 2000, 2001, or 2005 because reports by ground observers (Ferrari et al. 2002, Escudero et al. 2003) showed that very few Red Knots wintered at any of the sites at which they had previously been reported	57
FIGURE 31. Peak numbers of Red Knots during northward passage at (upper) Bahía San Antonio, Argentina 1990–2005 (P. M. González, unpubl. data) and (lower) Lagoa do Peixe, Brazil 1995–2003 (I. Serrano, unpubl. data). Counts	

at Bahía San Antonio were mostly carried out on a weekly basis throughout February to April. Counts at Lagoa do Peixe were obtained during expeditions that covered the peak spring passage in April.....	58
FIGURE 32. Peak counts of Red Knots in Delaware Bay May 1982–2006 as shown by weekly aerial counts. (NJAS (1982–1983), NJENSP (1986–2005). Also shown are simultaneous counts from other U.S. East Coast sites (mainly Virginia), the 1985 South America winter count (Morrison and Ross 1989), the authors’ estimate of the total range over which the U.S. East Coast flyway population fluctuated (range enclosed by dashed lines) and the estimates of the flyway population in 1999 of 60,000 (Baker et al. 1999a) and in 2005 of 32,728 (Table 14) shown by gray dots.....	59
FIGURE 33. Flight path of aerial surveys along the Delaware Bay conducted by the NJDFW	60
FIGURE 34. Stable isotope signatures of primary coverts taken from 1,150 Red Knots on spring migration through Delaware Bay in May and June 2004 (P. W. Atkinson, unpubl. data). Boxes mark the 90% confidence intervals of birds of known wintering origin. The large dot represents the signature of a tertial taken from a bird nesting on Southampton Island, Nunavut, Canada. Dotted lines show the approximate separation between juvenile birds (with freshwater Arctic signature) and the northern and southern wintering populations	61
FIGURE 35. Predicted population trends and associated 95% confidence limits of adults (dashed lines), juveniles (lower gray line) and both combined (top gray line) for 10 yr from 2000, with (A) constant adult survival of 85% and juvenile survival being half that of adults ($\lambda = 1$) and (B) constant adult survival of 56% and juvenile survival being half that of adults ($\lambda = 0.66$). The small dots represent the aerial censuses of the over-wintering flock of adults in Tierra del Fuego during 2000–2002, and the large dots are the counts during 2003–2006. The 95% upper and lower confidence limits are based on 1,000 bootstrap iterations. Modified from Baker et al. (2004) and published in this form in Baker et al. (2005a).....	62
FIGURE 36. Density of the nests of Red Knots and American Golden Plovers in a 9.2 km ² study site on Southampton Island, Nunavut, Hudson Bay, Canada, during 2000–2004. American Golden Plovers were not included in the survey until 2001.....	63
FIGURE 37. Total counts from aerial surveys of Red Knots done in Bahía Lomas, Tierra del Fuego, Chile	66
FIGURE 38. Total population estimates (\pm 95% confidence interval) of Red Knots spending the austral summer south of San Antonio Oeste, Río Negro, Argentina, from capture–recapture methods, compared with aerial census numbers (Morrison and Ross 1989, Morrison et al. 2004,) and number of Red Knots at Río Grande, Tierra del Fuego (González et al. 2004).....	66
FIGURE 39. Maximum counts of Red Knots during northward migration at two stopover sites in Argentina: San Antonio Oeste and Playa Fracasso in Península Valdés (references in Table 16).....	67
FIGURE 40. Peak numbers of Red Knots recorded in Lagoa do Peixe National Park, Brazil, 1995–2003 (CEMAVE-IBAMA).....	72
FIGURE 41. Number of Red Knots counted at Altamaha Estuary, Georgia, 1996–1998 (B. Winn, unpubl. data) and 2000 (B. A. Harrington, unpubl. data).....	77
FIGURE 42. Red Knot surveys conducted between 2000 and 2004 in the Cape Romain region of Charleston, South Carolina (sources provided in Table 24).....	78

FIGURE 43. Number of Red Knots counted at Cape Romain National Wildlife Refuge, South Carolina, 2000–2004 (S. Dawsey, pers. comm.)	80
FIGURE 44. Number of Red Knots counted at Harbor Island, South Carolina, 2001–2003 (B. A. Harrington, unpubl. data)	80
FIGURE 45. Number of Red Knots counted at Hunting Island, South Carolina, 2001–2003 (B. A. Harrington, unpubl. data)	80
FIGURE 46. Number of Red Knots counted at (A) Wallops Island, (B) Chincoteague, and Metompkin Island, Virginia, 1976–1982 (B. A. Harrington, unpubl. data)	84
FIGURE 47. Maximum historic Red Knot counts from three regions of Massachusetts: (A) western Cape Cod, (B) North Shore, and (C) eastern Cape Code during southbound migration (B. A. Harrington, unpubl. data).....	92
FIGURE 48. Number of Red Knots counted at Scituate (upper) and Monomoy (lower), Massachusetts, 1965–1986 (B. A. Harrington, unpubl. data).....	93
FIGURE 49. Counts of Red Knots on four islands (Nue, Grande, Quarry, and Niapiskau) of the Mingan archipelago in the Gulf of St Lawrence, Quebec, Canada, during July–September 2006 (Y. Aubry, pers. comm.)	95
FIGURE 50. Aerial counts of Red Knots (<i>Calidris canutus rufa</i>) on major wintering areas in southern South America, January–February 2000–2005 – Bahía Lomas and Río Grande, Chile. All sites are in the main wintering area (Morrison et al. 2004).....	99
FIGURE 51. Important <i>Calidris canutus rufa</i> breeding, stopover, and wintering areas in the Western Hemisphere	103
FIGURE 52. Important <i>Calidris canutus rufa</i> stopover and wintering areas in the United States.....	104
FIGURE 53. Red Knot state legal status in each state in the U.S.....	104

APPENDICES

APPENDIX 1. REGIONAL MAPS OF RED KNOT CRITICAL HABITAT (MIGRATORY STOPOVER AND WINTERING-NONBREEDING AREAS).....	146
APPENDIX 2. PEAK COUNTS OF RED KNOTS DURING SPRING AND FALL MIGRATION AT VARIOUS SITES IN FLORIDA. INTERSTITIAL PERIODS NOT INCLUDED. ISS = INTERNATIONAL SHOREBIRD SURVEY	173
APPENDIX 3. PEAK COUNTS OF RED KNOTS DURING WINTER AT VARIOUS SITES IN FLORIDA. INTERSTITIAL PERIODS NOT INCLUDED. ISS = INTERNATIONAL SHOREBIRD SURVEY.....	177
APPENDIX 4. SUMMARY TABLE OF LAND OWNERSHIP AND EXISTING HABITAT PROTECTION FOR POPULATIONS OF RED KNOTS IN THE WESTERN HEMISPHERE	181