

INDEX TO VOLUME I.

- ACANTHIS**, 145.
 canescens, 152.
 canescens exilipes, 152.
 hornemannii, 152.
 hornemannii exilipes, 152.
 linaria, 152.
 linaria cabaret, 153.
 linaria holboellii, 153.
 linaria pallescens, 153.
Accipiter, 58.
 cooperi, 247.
 fuscus, 247, 295.
Acredula caudata, 342.
Acrocephalus dybowskii, 82.
Actiturus, 142.
 bartramius, 315.
Actodromas acuminata, 78.
 maculatus, 218.
Adney, E. T., the Cardinal Gros-
 beak breeding in Brooklyn, N. Y.,
 390.
Æchmophorus major, 316.
Ægialites albidipectus, 83.
 melodus circumcinctus, 397.
 mongolicus, 78.
Ægialitis collaris, 313.
 falklandica, 313.
Ægiothus, 145, 146.
 canescens canescens, 152.
 canescens exilipes, 152.
 exilipes, 152, 159.
 fuscescens, 152.
 hornemannii, 152.
 linarius, 152, 159.
 linarius exilipes, 152.
 linarius hoboellii, 153.
 rostratus, 153.
 rufescens, 152.
Agelæus phœniceus, 245.
Aguilla, 109, 110.
 rojizo, 109.
Aix, 46, 144.
 sponsa, 187, 249.
Alauda ludoviciana, 168.
 minor, 260.
 mongolica, 228.
 pensylvanicus, 167, 168.
 rubra, 168.
 rufa, 261.
Albicilla, 140.
Alcon, 30.
Alconcito, 110.
Alle nigricans, 35, 115.
Allen, J. A., Zoölogical Nomencla-
 ture, 338.
**Alouette aux joues brunes de Pen-
 sylvanie**, 167, 168.
Aluco flammeus, 29.
 flammeus americanus, 246.
Aluette de Pensylvanie, 167, 168.
American Ornithologists' Union,
 Second Meeting of, 369.
Ammodromus maritimus, 176.
Ampelis cedrorum, 176, 244.
 garrula, 158, 244.
Anas boscas, 248.
 boschas, 122.
 obscura, 248.
Anorthura hiemalis, 15.
 pallescens, 82.
 troglydotes hiemalis, 138.
Anotis, 144.
Anser, 143.
 albifrons gambeli, 248.
Anthus japonicus, 82.
 ludovicianus, 89, 121, 176,
 186.
 pensylvanicus, 168, 209.
 pratensis, 77.
 stejnegeri, 82.
Antrostomus parvulus, 24.
Apteryx australis, 81.
Aquila, 140.
 chrysaëtus canadensis, 79, 247.
Aramides ypecaha, 276.
Aramus pictus, 115.
 scolopaceus, 277.
Archibuteo lagopus sancti-johannis,
 79, 247, 397.
Ardea cocoi, 271.
 candidissima, 271.
 egretta, 271.
 herodias, 247.
 occidentalis, 162.
 sibilatix, 271.
 wardi, 161.
 würdemanni, 162.
Ardetta exilis, 248.

- Ardetta involucris*, 271.
Arenaria interpres, 229.
melanocephala, 229.
Arquatella, 142.
couesi, 77.
maritima, 76.
Asio accipitrinus, 29, 115.
brachyotus, 115.
Astrigalinus tristis, 244, 328.
Astur atricapillus, 247.
atricapillus henshawi, 166.
atricapillus striatulus, 252.
badius, 340.
brevipes, 340.
poliopsis, 340.
polyzonoides, 340.
sphenurus, 340.
Asturina pucherani, 30.
Atthis, 57.
Auk, Black-throated, 98.
Great, 81, 208.
Auriparus flaviceps, 354.
Avestruz, 318.
Aythia americana, 249.
 BAILEY, H. B., breeding habits of the Everglade Kite, 95.
 Baird, Brewer, and Ridgway, notice of their work on the Water Birds of North America, 382.
 Banks, J. W., nesting of the Broad-winged Hawk (*Buteo pennsylvanicus*), 95; the Great Horned Owl (*Bubo virginianus*) in confinement, 194.
 Barrows, W. B., birds of the Lower Uruguay, 20, 109, 270, 313; abnormal coloration in a caged Robin, 90.
 Batchelder, C. F., description of the first plumage of Clarke's Crow, 16; Buffon's Skua in Vermont, 97.
 Becasino, 314.
 Belding, L., notices of his papers on birds of Lower California, etc., 83, 284.
Bernicla, 143.
canadensis, 80, 248, 309.
canadensis hutchinsi, 80.
canadensis leucoparia, 80.
canadensis occidentalis, 80.
brenta, 248.
brenta nigricans, 96.
nigricans, 187.
 Bicknell, E. P., a Study of the Singing of our birds, 60, 126, 209, 322.
 Bird, Cardinal, 189.
Cedar, 6, 294.
Chipping, 190.
 Bird, Cow. See Cowbird.
Guachera, 307.
Indigo, 6, 67, 86.
Phæbe, 65.
Summer Red, 290.
Thistle, 294.
 Bittern, American, 247.
Least, 188, 248.
 Blackbird, Cow. See Cowbird.
Crow, 6, 86, 185.
Red-winged, 6, 245.
Yellow-headed, 245, 293.
 Blackhead, Ring-billed, 249.
 Bluebird, 6, 13, 86, 133, 244.
Californian, 121.
 Bobolink, 6, 8, 64, 245.
Bolborhynchus monachus, 28.
Bonasa umbellus, 186, 247.
Botaurus, 143.
lentiginosus, 87, 247.
mugitans, 115.
Brachyrhamphus brachypterus, 386.
hypoleucus, 396.
kittlitzii, 81.
 Brant, 248.
Black, 187.
 Brewer, T. M., see Baird, Brewer, and Ridgway.
 Brewster, W., notes on the Summer Birds of Berkshire County, Mass., 5; notice of his annotations to the 'Birds and Feathered Fowles' of Morton's 'New English Canaan,' 84; on a specimen of *Helminthophaga leucobronchialis* from Connecticut, 91; recent occurrence of the Black-backed Three-toed Woodpecker in Massachusetts, 93; a singular specimen of the Black-and-White Creeper, 190; note on the Lesser Glaucous-winged Gull in New York, 196; an ornithological swindler, 295; notice of his Birds observed in the Gulf of St. Lawrence, 379.
 Brown, N. C., a second season in Texas, 120; the Widgeon in Maine in February, 394.
 Browne, F. C., the Loggerhead Shrike again in Massachusetts, 291; *Geothlypis trichas* wintering in Massachusetts, 389.
 Bryant, W. E., Nest and eggs of *Myiadestes townsendi*, 91.
 Bubo, 57.
virginianus, 29, 194, 246.
virginianus subarcticus, 79.
Budytes flava, 77.

- Buffalo Naturalists' Field Club,
 notice of its 'Bulletin,' 184.
 Buho, 29.
 Bunting, Cow, 189.
 Lark, 286, 293.
 Snow, 159, 245, 294.
 Towhee, 13, 245.
 Buteo, 140.
 abbreviatus, 122.
 borealis, 123, 247.
 borealis krideri, 100, 123.
 cooperi, 253.
 erythronotus, 109.
 harlani, 253.
 pennsylvanicus, 95, 188.
 pteroles, 109.
 swainsoni, 247.
 Butorides cyanurus, 271.
 Butterball, 249.
 Buzzard, Turkey, 187, 247, 286, 293.
 CABURÉ, 29.
 Calamospiza bicolor, 121, 293.
 Calcarius, 118.
 nivalis, 118.
 Calendula, 56.
 Calidris, 142.
 arenaria, 397.
 Calodromas elegans, 318.
 Calypte costæ, 354.
 Calyptophilus, 3.
 frugivorus, 3.
 Campephilus boiæi, 25.
 Campylæmus labradorius, 207.
 Canace canadensis, 15, 247.
 Caprimulgus vociferus, 246.
 Cardinal, Texas, 122.
 Cardinalis ruber, 172.
 ruber carneus, 172.
 ruber coccineus, 172.
 ruber igneus, 172.
 virginianus, 172, 390.
 Carpenter, J. H., the nest of the
 Saw-whet Owl, 94.
 Carpintero, 25.
 Carpintero, 20, 25.
 blanco, 25.
 Carpodacus cassini, 288.
 frontalis, 288, 289, 355.
 purpureus, 288, 327.
 Carrancho, 111.
 Catbird, 6, 13, 132, 185, 244.
 Cathartes atratus, 113, 285.
 aura, 87, 113, 247, 293.
 Chætura pelagica, 161, 246.
 Chaja, 273.
 Certhia familiaris, 281.
 familiaris β americana, 281.
 familiaris rufa, 121, 136, 157.
 Ceryle alcyon, 179, 246.
 amazona, 26.
 americana, 26.
 torquata, 26.
 Chamberlain, M., nest and eggs of
 the Ruby-crowned Kinglet (*Regulus
 calendula*), 90; prehensile
 feet of the Crow (*Corvus frugivo-
 rus*), 92; a Woodpecker destroy-
 ing Cocoons, 93; Are trinomials
 necessary? 101, 200; New Brun-
 swick winter notes, 294; the
 Crow (*Corvus frugivorus*) as a
 fisherman, 391.
 Charadrius dominicus, 179, 248.
 fulvus, 78.
 virginicus, 313.
 Chat, Yellow-breasted, 67, 216, 389.
 Chauna chavaria, 272.
 Chen cærulescens, 240.
 hyperboreus, 34, 240, 386.
 Chickadee, Black-capped, 6, 8, 135,
 157, 244.
 Hudsonian, 185.
 Chimango, 111.
 Chloëphaga magellanica, 273.
 Chloris, 168.
 Chlorostilbon splendidus, 22.
 Chondestes grammicus, 121.
 Chordiles popetue, 246.
 virginianus, 24.
 Chorlito, 313.
 Chorlo, 313, 315.
 Chrysomitris, 361.
 pinus, 160, 176, 177, 328.
 Chrysoptilus cristatus, 25.
 Ciceronia pusilla, 81.
 Ciconia maguari, 271.
 Ciguëña, 271.
 Cinclus aquaticus, 345.
 Circus cinereus, 30.
 hudsonius, 19, 122, 185, 247.
 maculosus, 30.
 Cistothorus stellaris, 59, 139, 244.
 Cisne, 271.
 Clangula albeola, 249.
 glaucium americana, 177, 249.
 Clark, J. N., nesting of the Little
 Black Rail in Connecticut, 393.
 Coccyzus cinereus, 28.
 erythrophthalmus, 178, 179.
 melanocosyphus, 28.
 pumilus, 28.
 Colaptes agricola, 25.
 auratus, 246.
 Collins, J. W., notes on certain
 Laridæ and Procellariidæ of the
 New England Coast, 236; notice
 of his paper on Sea Birds used
 as fish-bait by fishermen, 380.

- Columba, 140.
 maculosa, 274.
 picazura, 274.
 Columbula *picui*, 274.
 Colymbus *adamsi*, 81.
 arcticus, 207.
 glacialis, 115.
 torquatus, 115, 185, 249.
 Compsothlypis *americana*, 169, 170, 212.
 gutturalis, 169.
 insularis, 169, 170.
 nigrilora, 169, 170.
 pitiayumi, 170.
 pitiayumi inornata, 170.
 Condor, California, 81.
 Contopus *borealis*, 15.
 Conurus *patagonus*, 28.
 Cooke, W. W., bird nomenclature of the Chippewa Indians, 242; the distribution and migration of *Zonotrichia querulea*, 332.
 Coot, American, 248.
 Cormorant, 189, 196.
 Double-crested, 249.
 Pallas's, 82.
 Violet-green, 163.
 Corvus *corax carnivorus*, 160, 245.
 cryptoleucus, 100.
 frugivorus, 115, 160, 245, 391.
 Cory, C. B., descriptions of several New Birds from Santo Domingo, 1, pl. i; notice of his 'Beautiful and Curious Birds,' 81; *Thryothorus ludovicianus* in Massachusetts, 91; *Bernicla brenta nigricans* in Massachusetts, 96; notice of his 'Birds of Haiti and San Domingo,' 285; the generic name *Ligea*, 290.
 Corythus *enucleator*, 115.
 Cosmonetta *histrionica*, 115.
 Coturniculus *passerinus*, 330.
 Coues, E., ornithophilologicalities, 49, 140; notice of his 'A hearing of Bird's Ears,' 182; trinomials are necessary, 197; notice of his 'Key to North American Birds,' second edition, 282; egg of Cowbird in nest of the Carolina Dove, 293; on some new terms recommended for use in Zoölogical Nomenclature, 320; Strickland as an advocate of 'Linnæus at '58,' 400.
 Coues, E., and D. W. Prentiss, notice of their Avifauna Columbi-ana, 386.
 Cowbird, 186, 193, 245, 293.
 Crane, Carolina, 399.
 Corn, 399.
 Alaskan, 80.
 Sand-hill, 85, 248.
 Whooping, 248, 288.
 Creeper, Black-and-white, 6, 87, 190.
 Crex, Brown, 136, 157, 294.
 pratensis, 399.
 Crispin, 27.
 Crossbill, Red, 88, 159, 292, 327.
 White-winged, 159.
 Crow, 6, 92, 160, 188, 189, 245, 294, 391.
 Carion, 285.
 Clarke's, 16.
 Cuckoo, 28.
 Cuculillo, 28.
 Cuervo, 112.
 del agua, 70.
 Cupidonia *cupido*, 247, 391.
 Curlew, Bristle-billed, 79.
 Eskimo, 393.
 Cyanocephalus *cyanocephalus*, 230.
 Cyanocitta *cristata*, 160, 245.
 Cyanospiza *parellina*, 88.
 Cygnus *americanus*, 89, 115.
 columbianus, 115.
 musicus, 173.
 nigricollis, 273.
 Cymochorea *leucorrhœa*, 98.
 Cypseloides niger borealis, 207.
 DAFILA *acuta*, 248.
 bahamensis, 274.
 spinicauda, 274.
 Dagleish, J. J., occurrence of the Royal Tern (*Sterna regia*) at Tangiers in Morocco, 97; a newly-discovered breeding-place of Leach's Petrel (*Cymochorea leucorrhœa*) in Scotland, 98.
 Davison, J. L., Wilson's Petrel in Western New York, 294.
 Dendrocopos *purus*, 35.
 Dendrocœa, 47.
 adelaidæ delicata, 83.
 æstiva, 175, 176, 212, 244.
 auduboni, 100.
 blackburniæ, 15, 121, 178.
 cærulescens, 15, 177, 179, 213, 244.
 castanea, 213.
 chrysoparia, 121.
 coronata, 121, 176, 177, 192, 213.
 discolor, 176.
 kirtlandi, 389.
 maculosa, 14, 15, 176, 213.
 palmarum hypochrysea, 213.
 pennsylvanica, 213.
 pinus, 14, 176, 178.

- Dendroeca striata*, 176, 177, 178, 179, 213.
 tigrina, 59, 213.
 virens, 178, 213.
Dicromanassa, 143.
Diplopterus galeritus, 27.
Dodo, 81.
Dolichonyx oryzivora, 176, 245.
Dormilon, 24.
Dove, Mourning, 247.
 Sea, 35.
Dovekie, 35.
Drew, F. M., notes on *Lagopus leucurus*, 392.
Drymornis bridgesi, 20.
Duck, Eider, 184, 289.
 King, 184.
 Labrador, 81.
 Pintail, 248.
 Red-head, 185, 188.
 Scaup, 249.
 Wood, 187, 249.
Dutcher, W., bird notes from Long Island, N. Y., 31, 174.
Dytes auritus, 249.
 nigricollis, 249.
 californicus, 18.
EAGLE, Bald, 247.
 Golden, 10, 247.
 Great Northern Sea, 140.
Ectopistes migratorius, 185, 247.
Egret, American, 32.
Eider, Spectacled, 80.
Elanus glaucus, 115.
 forficatus, 247.
Emberiza nivalis, 118.
Empidonax, 42.
 flaviventris, 15, 295.
Engyptila wellsi, 180.
Eremophila alpestris, 86, 160, 179, 245.
 alpestris chrysolæma, 121, 123.
Erismatura dominica, 274.
Erythacus, 117.
Espatula, 272.
Eudocimus albus, 239.
 ruber, 239.
Eudromias modesta, 313.
Eurynorhynchus pygmaeus, 78, 81.
FALCO columbarius, 399.
 peregrinus, 397.
 peregrinus nævius, 79, 295, 391.
 peregrinus pealii, 79.
 tinnunculus, 398.
Farlouzanne, 167, 168.
Fewkes, J. W., Do Crows carry objects in their claws? 92; Ducks transporting clams, 195.
Finch, Grass, 6, 8, 11, 31, 330.
 Harris's, 332.
 Lincoln's, 31.
 Pine, 160.
 Purple, 6, 13, 63, 67, 288, 327.
Flicker, Yellow-shafted, 246.
Flycatcher, Canada, 9, 11, 13.
 Couch's Tyrant, 93.
 Great-crested, 7, 65, 86, 391.
 Least, 6.
 Olive-sided, 7, 12.
 Townsend's, 286.
 Yellow-billed, 12.
Fox, W. H., *Dendroeca coronata* in Southern New Hampshire in Summer, 192.
Frazer, J. T., the Turkey Buzzard in Western New York, 293.
Fringilla borealis, 152.
 cabaret, 153.
 canescens, 153.
 lapponica, 118.
 linaria, 152.
 linaria alnorum, 153.
 linaria β , 154.
 linaria betularum, 152.
 linaria brevirostris, 152.
 linaria β *holböllii*, 153.
 linaria magnirostris, 153.
 linaria var. β , 153.
 minima, 153.
Fulica, 143.
 americana, 248.
 armillata, 277.
 leucoptera, 277.
Fulix affinis, 249.
 collaris, 249.
 marila, 178, 249.
Fulmar, Slender-billed, 80, 233.
Fulmarus glacialis, 233, 238.
 glacialis glupischa, 234.
 glacialis rogersi, 233.
 glacialoides, 233.
GADOW, H., notice of his Catalogue of the Paridæ and Laniidæ, 279.
Galeoscoptes carolinensis, 175, 176, 244.
Gallina del agua, 277.
Gallinago media wilsoni, 248.
 paraguaia, 314.
 wilsoni, 187.
Gallinita chica, 276.
 grande, 276, 277.
Gallinula galeata, 277.
Ganso de la sierra, 273.
Garza, 271.
 blanca, 271.
 blanca chica, 271.
Garzetta, 143.
 candidissima, 32.

- Gavilan, 30.
 Gaviota, 316.
 Gaviotila, 316.
 Geothlypis philadelphia, 14, 15.
 trichas, 175, 176, 177, 178, 179,
 215, 389.
 Gelochelidon nilotica, 365.
 Gill, T., notice of his 'Record of
 Scientific Progress for 1881,' 84.
 Golden-eye, American, 249.
 Goldfinch, American, 6, 64, 244,
 329.
 Goniaphea ludoviciana, 14.
 Goose, Blue-winged, 240.
 Canada, 80, 188, 248, 309.
 Emperor, 80.
 Snow, 34, 184, 240.
 White-fronted, 248.
 Goshawk, 247.
 Henshaw's, 166.
 Goss, N. S., notes on the breeding
 habits of the American Eared
 Grebes (*Dytes nigricollis californicus*), 18; birds new to the
 Fauna of Kansas and others rare
 in the State, captured at Wallace,
 Oct. 12 to 16, 1883, 100; notes
 on the nesting of the Yellow-
 throated Vireo (*Lanivireo flavifrons*), 124; notes on *Phalacrocorax violaceus*, and on *P. violaceus resplendens*, 163; capture of *Megalestris skua* off the coast of Cape Cod, Mass., 395; *Brachyrhamphus hypoleucus* off the coast of Southern California, 396.
 Grackle, Purple, 245.
 Graculus bairdi, 165.
 Grebe, American Eared, 18.
 Horned, 249.
 Thick-billed, 122, 249.
 Grosbeak, Black-headed, 367.
 Cardinal, 171, 289, 390.
 Evening, 244.
 Pine, 158, 244, 294, 327.
 Rose-breasted, 8, 9, 70, 245,
 367.
 Grouse, Canada, 247.
 Newfoundland Willow, 369.
 Ruffed, 9, 186, 247.
 Grus americana, 248.
 canadensis, 248.
 fraterculus, 80.
 Guillemot, Xantus's, 396.
 Guira piririgua, 26.
 Gull, Ivory, 80.
 Kittiwake, 380.
 Lesser Glaucous-winged, 196.
 Ring-billed, 19.
 Skua, 395.
 Gull, White-winged, 240, 287.
 Gurney, J. H., Jr., hybrid between
 Pedieccetes phasianellus and *Cupidonia cupido*, 391.
 Gybitz, 120.
 Gyrfalcon, 58, 94.
 HABIA, 366, 367.
 ludoviciana, 367.
 melanocephala, 367.
 Hæmatopus ostrilegus, 172.
 Hag, 237.
 Hagdon, 237.
 Haliaëtus albicilla, 140.
 hypoleucus, 82.
 leucocephalus, 247.
 Harelda, 144, 173.
 glacialis, 177.
 Harporhynchus lecontei, 355, 356.
 rufus, 13, 132, 175, 244.
 Hawk, Broad-winged, 95.
 Ccooper's 247.
 Duck, 391.
 Fish, 86, 247.
 Kriider's, 100.
 Marsh, 19, 122, 247.
 Red-shouldered, 186.
 Red-tailed, 123, 247.
 Rough-legged, 247.
 Sharp-shinned, 247.
 Sparrow, 247.
 Swainson's, 247.
 Swallow-tailed, 189.
 Hazard, R. G., 2d, the occurrence
 of the Golden Swamp Warbler
 (*Protonotaria citrea*) in Rhode
 Island, 290; a lay view of ornithophilologicalities, 300.
 Heliomaster furcifer, 21.
 Helmintherus, 49.
 Helminthophaga chrysoptera, 186.
 leucobronchialis, 91.
 Helminthophila celata, 121.
 peregrinus, 212.
 pinus, 210, 397.
 ruficapilla, 14, 211.
 Helminthotherus vermivorus, 210.
 Hemiprocne zonaris, 24.
 Helmitheus, 49.
 Hendrickson, W. F., capture of the
 Summer Redbird on Long Island,
 290.
 Henshaw, H. W., description of a
 new Song Sparrow from *the
 southern border of the United
 States, 223; on a new Gull from
 Alaska, 250; on the Shore Larks
 of the United States and adjacent
 territory, 254; Indian bird names,
 400.
 Herodias alba egretta, 32.

- Herodias egretta, 59.
 Heron, Great Blue, 247.
 Great White, 59.
 Night, 96, 122.
 Snowy, 32.
 Hesperiphona vespertina, 244.
 Heteractitis incanus, 236.
 Heteroscelus incanus, 79.
 Heterospizias meridionalis, 109.
 Hiaticula, 141.
 Hierofalco, 58.
 gyrfalco, 79.
 gyrfalco sacer, 79, 94.
 Hirundo bicolor, 325.
 erythrogastra, 325, 368.
 saturatus, 82.
 sclateri, 2, 285.
 Histrionicus minutus, 115.
 Holterhoff, G., *Calamospiza bicolor*
 in Southern California, 293;
 occurrence of the Least Tern at
 San Diego, Cal., 294; Eskimo
 Curlew at San Diego, Cal., 393.
 Hornemanni, 57.
 Hummingbird, Black-chinned, 123.
 Ruby-throated, 246.
 Hydranassa, 143.
 Hydrochelidon lariformis, 59.
 Hypopsalis fuscifera, 24.
 Hylocharis sapphirina, 22.
 Ilyocichla alicia, 178, 179.
 fuscescens, 178, 179, 243.
 mustelina, 284.
 unalascae pallasi, 121, 176, 178.
 ustulata swainsoni, 176, 178,
 179, 243.
 Hylotomus pileatus, 246.
 Ibis, Sacred, 81.
 Scarlet, 239.
 White, 239.
 Wood, 297.
 Icteria virens, 216, 389.
 Icterus galbula, 245.
 spurius, 390.
 Ictinia, 58.
 subcaerulea, 115, 122.
 Iliacus, 47, 56.
 JAEGER, Arctic, 237.
 Jay, Blue, 85, 86, 160, 245.
 Canada, 160, 245.
 Piñon, 230.
 Jeffries, J. A., notice of his paper
 on the 'Epidermal System of
 Birds,' 182; the occipital style of
 the Cormorant, 196.
 Jencks, F. T., another Gyrfalcon in
 Rhode Island, 94: Can we not
 have a simpler system of nomen-
 clature? 297.
 Jouy, P. L., notice of his paper on
 birds collected in Japan, 387.
 Junco bairdi, 284.
 hiemalis, 14, 15, 177, 178, 179,
 245.
 hiemalis connectens, 283.
 oregonus, 256.
 vulcani, 84.
 KIEBITZ, 120.
 Kildeer, 248.
 Kingbird, 6, 8, 87, 122, 245.
 Kingfisher, 246.
 Kinglet, Golden-crowned, 134, 294.
 Ruby-crowned, 90, 134.
 Kite, Everglade, 95.
 Swallow-tailed, 247, 288, 289.
 Kiwi, 81.
 Knowlton, F. II., the Orchard
 Oriole (*Icterus spurius*) in West-
 ern Vermont, 390.
 Knot, 307.
 Kybitz, 120.
 Kywitz, 120.
 LAGOPUS alba, 79, 82, 225.
 alba alleni, 369.
 alpinus, 80, 225, 227.
 leucurus, 392.
 muta, 226.
 mutus atkhensis, 79.
 rupestris, 79, 225.
 rupestris nelsoni, 226.
 rupestris occidentalis, 79, 225.
 rupestris atkhensis, 226.
 ridgwayi, 226.
 subalpinus, 227.
 Langille, J. H., Bicknell's Thrush,
 268.
 Lanius borealis, 158, 185, 186, 244,
 291, 344.
 bucephalus, 292.
 cristatus, 77, 81, 291.
 ludovicianus, 186, 193.
 major, 292.
 Lanivireo flavifrons, 124, 176.
 solitarius, 122.
 Laopteryx priscus, 84.
 Lark, Desert Horned, 264.
 Horned, 87, 123, 254, 285.
 Louisiana Horned, 167, 168.
 Meadow, 6, 245.
 Prairie Horned, 264.
 Red, 167, 168.
 Ruddy Horned, 267.
 Shore, 89, 160, 189, 245, 254,
 Sprague's, 286.
 Streaked Horned, 267.
 Texan Horned, 266.
 Tit, 89, 186. (See also Tit-
 lark.)

- Larus affinis*, 80, 359.
argentatus, 232.
cachinnans, 80, 232.
chalcopterus, 252.
cirrocephalus, 316.
delawarensis, 19.
dominicanus, 316.
fuscescens, 232.
glaucescens, 287, 360.
kumlieni, 196, 251.
leucopterus, 240, 287, 360.
marinus, 231, 232.
nelsoni, 250.
pelagicus, 232.
schistisagus, 231.
 Lawrence, G. N., characters of a new species of Pigeon of the genus *Egyptila*, from the Island of Grenada, W. I., 180; on the occurrence of the White-winged Gull (*Larus leucopterus* Faber) in the State of New York, 240; notice of his paper on new American Birds, 387.
Lechuza, 30.
Lechuzon, 29.
Lepidocolaptes atripes, 21.
Leptoptila chalcauchenia, 275.
Leucibis alba, 236.
 rubra, 236.
Leuconerpes candidus, 25.
Ligea, 1, 236.
 palustris, 1, 285.
Limosa lapponica novæ-zealandiæ, 79.
 uropygialis, 79.
Linaria, 145, 146.
 agrorum, 152.
 alnorum, 153.
 alnorum magnirostris, 153.
 betularum, 152.
 borealis, 152.
 brunnescens, 153.
 canescens, 153.
 lanceolata, 153.
 longirostris, 153.
 minor, 154.
 rufescens, 154.
Linnet, Pine, 328.
Linota canescens, 153.
 exilipes, 153.
 hornemannii, 152.
 rufescens, 154.
Lintner, G. A., Duck Hawks breeding in the Helderberg Mts., N. Y., 391.
Lobipes hyperboreus, 33.
 tricolor, 367.
Loomis, L. M., *Xanthocephalus icterocephalus* in Chester County, South Carolina, 293.
Loon, 186, 249.
Lophodytes cucullatus, 249.
Lophophanes bicolor, 134.
 inornatus cinerascens, 284.
Lophortyx californicus, 355.
 gambeli, 355.
Lorito, 28.
Loro, 28.
Loxia curvirostra, 88.
 curvirostra americana, 159, 292, 327.
 leucoptera, 159.
 rubra, 171.
MACHETES pugnax, 186.
Macrorhamphus griseus scolopaceus, 32.
Magpie, 245, 286.
Mallard, 122, 188, 248.
 Black, 248.
Marbleheader, 238.
Mareca americana, 394.
 penelope, 80.
 sibylatrix, 274.
Marling-spike, 237.
Martin pescador, 26.
 Purple, 6, 244.
Martinet, 318.
Mbatitú, 314.
Mbatitúí, 314.
McCormick, L. M., 'Avifauna Columbiana'—a Protest, 396.
McIlwraith, T., the Yellow-breasted Chat and Summer Redbird in Canada, 389; Pelicans on the move, 395.
Melanerpes, 41.
 erythrocephalus, 189, 246.
 formicivorus, 87.
Meleagris gallopavo americana, 247.
Melospiza fasciata, 176, 177, 178, 245.
 fasciata fallax, 223.
 fasciata montana, 224.
 lincolni, 31, 121, 175, 295, 397.
 palustris, 14, 176, 295.
Megalestris skua, 395.
Mergus alle, 115.
Mergus merganser americanus, 249.
 serrator, 177, 249.
Merle d'eau, 120.
Merriam, A. C., the Coues Lexicon of North American Birds, 36.
Merriam, C. H., third addendum to the Preliminary List of the Birds ascertained to occur in the Adirondack Region, Northeastern

- New York, 58; Bird Migration, 71; a plea for the metric system in ornithology, 203; *Vireo philadelphicus* in Northern New York, 291; second addendum to List of Birds ascertained to occur within ten miles from Point de Monts, Province of Quebec, Canada; based chiefly upon the notes of Napoleon A. Comeau, 295; breeding of *Passerculus princeps* on Sable Island, 390.
- Merula* confinis, 285.
 migratoria, 161, 178, 244.
 migratoria propinqua, 100.
- Metopiana peposaca*, 274.
- Microligea palustris*, 290.
- Micropus affinis*, 230.
 apus, 230.
 melanoleucus, 230.
 melba, 230.
 pallidus, 230.
 saxatilis, 230.
 unicolor, 230.
- Milvulus tyrannus*, 207.
- Mimocichla ardesiaca*, 285.
- Mimus carolinensis*, 132.
 polyglottus, 131, 192.
- Minor ornithological publications, notices of, 85, 185, 284.
- Mniotilta varia*, 176, 190, 210.
- Mockingbird, 69, 131, 192, 286.
- Molothrus*, 45.
 ater, 125, 186, 245.
- Morden, J. A., Cowbirds in a Black-and-white Creeper's Nest, 193.
- Morden, J. A., and W. E. Saunders, notice of their 'List of the Birds of Western Ontario,' 85.
- Motacilla*, 44, 56.
 amurensis, 387.
 blakistoni, 387.
 eques, 170.
 ludoviciana, 170.
 ocularis, 77, 81, 387.
 yarrelli, 398.
- Myiadestes*, 42, 56.
 leucotis, 182.
 montanus, 285.
 reevei, 182.
 townsendi, 91, 100.
- Myiarchus crinitus*, 391.
- Myiodioctes canadensis*, 15, 178, 217.
 mitratus, 216.
 pusillus, 217.
- NELSON, E. W., notice of his 'Birds of Bering Sea and the Arctic Ocean,' 76; brief diagnoses of two new races of North American birds, 165; the Ornithological Report of the 'Cruise of the Corwin,' 202; the breeding habits of the Pectoral Sandpiper (*Actodromas maculatus*), 218.
- Neocorys spraguui*, 121.
- Nettion carolinensis*, 248.
- Nighthawk, 246.
- Noddy, 238.
- Nothura maculosa*, 317.
- Numenius arquata*, 142.
 borealis, 316, 393.
 femoralis, 79.
 taitensis, 79.
- Nuthatch, Red-bellied, 13, 136, 157, 244, 294.
 White-bellied, 135, 157, 244.
- Nyctale acadica*, 94, 397.
 tengmalmi, 79.
 tengmalmi richardsoni, 79.
- Nyctea scandiaca*, 246.
- Nyctiardea grisea nævia*, 96, 122.
- Nycticorax gardeni*, 271.
- OCEANITES oceanica, 294, 380.
- Œdemia americana*, 178.
- Œdicnemus dominicensis*, 4.
- Œstrelata fisheri*, 83, 108.
 gularis, 83.
- Oil-bird, 238.
- Oporornis agilis*, 179, 192.
- Oreortyx picta plumifera*, 355.
- Oreothlypis gutturalis*, 169.
 superciliosa, 169.
- Oriole, 57.
 Baltimore, 6, 8, 65, 245.
 Orchard, 67, 68, 390.
- Ortyx, 141.
 virginianus, 6, 246.
- Ostrælegus*, 142.
- Ostrilegus*, 142.
- Otocorys alpestris*, 257, 263.
 alpestris arenicola, 259, 263, 265.
 alpestris chrysolæma, 260, 263.
 alpestris geraudi, 260, 263, 266.
 alpestris leucolæma, 258, 263.
 alpestris praticola, 258, 263, 264.
 alpestris rubeus, 260, 263, 266.
 alpestris strigata, 261, 264, 267.
 occidentalis, 259.
- Otomela cristata*, 292.
- Ovenbird, 6, 8, 9, 13, 213.
- Owl, American Hawk, 363.
 Barn, 246.

- Owl, Barred, 88, 246.
 Burrowing, 286.
 European Hawk, 363.
 Great Gray, 246, 286.
 Great Horned, 194, 246.
 Hawk, 246.
 Little Screech, 246.
 Long-eared, 246.
 Saw-whet, 94.
 Short-eared, 246.
 Snowy, 186, 246.
 White, 185.
- Oxyechus vociferus, 248.
- PAGOPHILA eburnea, 80.
- Pallasia, 228.
- Paloma, 275.
 del monte, 274.
- Palomita, 274.
- Pandion haliaëtus carolinensis, 79.
- Parisorum, 57.
- Park, A. F., the Lesser Glaucous-winged Gull in New York, 196.
- Parra jacana, 277.
- Parula americana, 170, 175, 176, 178, 179.
 brasiliana, 170.
 inornata, 170.
 insularis, 170.
 nigrilora, 170.
 pitiaiyumi, 170.
 pitiaiyumi inornata, 170.
 pitiaiyumi insularis, 170.
 pitiaiyumi nigrilora, 170.
- Parus americanus, 170.
 atricapillus, 135, 157, 244, 397.
 carolinensis, 397.
 cinctus, 77.
 cinctus griseescens, 78.
 hudsonius, 15, 88, 157, 185.
 hudsonius evura, 78, 283.
- Passerculus anthinus, 106.
 guttatus, 83.
 princeps, 31, 292, 390.
 rostratus, 83.
 sanctorum, 83.
 sandwichensis savanna, 176, 177, 178, 179, 329.
- Passer, 57.
 domesticus, 87, 159, 208, 295.
- Passerella iliaca, 177, 178, 295.
- Passerina cyanea, 295.
- Patito, 273, 274.
- Pato, 273.
- Pavo del monte, 274.
- Pediæcetes phasianellus, 391.
- Pedincho, 26.
- Pelican, White, 395.
- Pelidna subarquata, 32.
- Penelope obscura, 275.
- Perdiz, 317.
 grande, 317.
- Perisoreus, 57.
 canadensis, 15, 160, 245.
- Perspicillata, 144, 173.
- Petrel, Leach's, 98.
 Wilson's, 294.
- Petrochelidon pyrrhonota, 368.
- Pewee, Bridge, 6.
 Say's, 122.
 Wood, 6, 8.
- Phalacrocorax, 144.
 bicristatus, 189.
 brasilianus, 270.
 dilophus, 249.
 perspicillatus, 82, 144.
 violaceus, 163.
 violaceus resplendens, 163.
- Phalarope, Northern, 33.
 Red, 33.
 Wilson's, 33.
- Phalaropus fulicarius, 33, 142.
 wilsoni, 367.
- Philohela minor, 248.
- Phœnicopterus ignipalliatu, 272.
- Phoneus brachyurus, 292.
- Phylacte canagica, 80.
- Phyllopneste borealis, 227.
 evermanni, 227.
 trochilus, 227.
- Phylloscopus borealis, 77.
- Pica caudata, 115.
 rustica, 115.
 rustica hudsonica, 245.
- Picafior, 22.
 mayor, 21.
- Picicorvus columbianus, 17.
- Picoïdes arcticus, 161, 246, 288.
 trydactylus alascensis, 165.
 trydactylus americanus, 161, 295.
- Picus cactorum, 25.
 mixtus, 25.
 nuttalli, 207.
 pubescens, 122, 161, 246.
 villosus, 161, 246.
- Pigeon, Passenger, 185, 186, 247.
- Pilincho, 26.
- Pipit, American, 186.
- Pinicola enucleator, 158, 244, 327.
 enucleator flammula, 149.
 flammula, 149.
- Pipilo erythrophthalmus, 14, 175, 245.
- Platalea ajaja, 272.
- Plectrophanes, 118.
 nivalis, 159, 177, 178, 245.
- Plectrophenax, 119.
- Plegadis falcinellus, 272.

- Plotus anHINGA, 121.
 Plover, Golden, 248.
 Killdeer, 248.
 Mongolian, 78.
 Mountain, 183.
 Podager nacunda, 24.
 Podasocys montanus, 121, 183.
 Podiceps holboelli, 177.
 rollandi, 317.
 Podilymbus podiceps, 122, 249.
 Polioptila cærulea, 83.
 cærulea obscura, 83.
 Polysticta stelleri, 80.
 Poœcetes gramineus, 31, 330.
 Porzana carolina, 97, 176, 248.
 jamaicensis, 393, 397.
 maruetta, 143, 173.
 noveboracensis, 397.
 Prentiss, D. W., see Coues, E.
 Priocella glacialis, 233.
 tenuirostris, 80, 233.
 Procellaria curilica, 234.
 pacifica, 234.
 tenuirostris, 233, 234.
 Progne concolor, 368.
 elegans, 368.
 furcata, 368.
 purpurea, 368.
 subis, 244, 368.
 Protonotaria citrea, 290.
 Psaltriparus grindæ, 284.
 Pseudotantalus, 83.
 Ptarmigan, Alpine, 227.
 Atkhan, 79, 225.
 Rock, 79, 225.
 Subalpine, 227.
 Pterocorys, 228.
 Publications received, lists of, 89,
 190, 289.
 Puffinus fuliginosus, 237, 380.
 major, 237, 380, 381.
 tenuirostris, 108, 223.
 Pyrrhula æstiva, 290, 390.
 rubra, 14, 178, 179, 244, 326.
 Pyrrhula, 48, 56.
 cassini, 77.
 Pyrrhuloxia sinuata, 122.
 QUAIL, 186, 188, 247, 287.
 Migratory, 186.
 Querquedula, 144.
 brasiliensis, 273.
 cyanoptera, 273.
 discors, 248.
 flavirostris, 273.
 versicolor, 274.
 Quiscalus, 57.
 æneus, 86.
 major, 185.
 purpureus, 245.
 RAIL, Little Black, 393.
 Rail, Red-breasted, 97.
 Sora, 248.
 Virginian, 399.
 Rallus antarcticus, 276.
 elegans, 97.
 longirostris crepitans, 175.
 maculatus, 276.
 rythrhynchus, 276.
 virginianus, 178, 399.
 Raven, 10, 160, 245.
 White-necked, 100.
 Redbird, Summer, 390.
 Redhead, 249.
 Redpoll, American Mealy, 159.
 Lesser, 159.
 Redstart, 6, 9, 11, 69, 217.
 Regulus calendula, 90, 134, 185.
 satrapa, 134, 177.
 Rhea americana, 318.
 Rhodostethia rosea, 80.
 Rhyacophilus solitarius, 315.
 Rhyncæa semicollaris, 314.
 Rhynchotus rufescens, 317.
 Rhyncophanes maccowni, 121.
 Ridgway, R., notice of his paper on
 supposed undescribed birds from
 the Commander Islands, etc., 82;
 do. on a new Warbler from the
 Island of Santa Lucia, W. I., 83;
 do. on a new Plover from Chili,
 83; do. on a new Petrel from
 Alaska, 83; do. on the genus
 Tantalus, 83; do. on a collection of
 Birds from Costa Rica, 84;
 note on *Zenaidura yucatanensis*
 Lawr., 96; note on *Phalacrocorax*
 violaceus resplendens, 165;
 remarks upon the close relation-
 ship between the White and
 Scarlet Ibises (*Eudocimus albus*
 and *E. ruber*), 239; note on
 Astur atricapillus striatulus,
 252; on the possible specific
 identity of *Buteo cooperi* Cass.
 with *B. harlani* (Aud.), 253;
 notice of his paper on new birds
 from Lower California, 284;
 probable breeding of the Red
 Crossbill (*Loxia curvirostra*
 americana) in Central Maryland,
 292; probable breeding-place of
 Passerculus princeps, 292; notice
 of his paper on rare species of
 Neotropical birds, 386; do. on
 the Pied Wagtails of Eastern
 Asia, 387; another Kirtland's
 Warbler in Michigan, 389.
 Rissa, 144.
 brevirostris, 80, 358.

- Rissa tridactyla*, 380.
tridactyla kotzebuii, 80.
 Robin, 6, 8, 9, 70, 87, 90, 126, 160, 188, 189, 244.
 Western, 100, 123.
Rostrhamus sociabilis plumbeus, 95.
Rupornis ridgwayi, 4.
 SAGE, J. H., another example of *Helminthophaga leucobronchialis* from Connecticut, 91.
Salpinctes obsoletus, 121.
 Sandpiper, Curlew, 32.
 Pectoral, 218, 248.
 Sharp-tailed, 78.
 Saunders, W. E., see Morden, J. A.
 Savanna, 57.
Sayornis, 38.
Sayornis fuscus, 179.
 sayi, 122.
Saxicola oenanthe, 295, 378, 398.
Scolopax, 142.
 rusticula, 173.
Scops asio, 246.
 brasilianus, 29.
 giu, 340.
 Scoter, Velvet, 80, 184.
 Scott, W. L., the winter Passeres and Picariæ of Ottawa, 156; the generic name *Troglodytes*, 400.
 Sea-hen, 238.
 Sennett, G. B., nest and eggs of Couch's Tyrant Flycatcher (*Tyrannus melancholicus couchi*), 93; Black-throated Auk (*Synthliboramphus antiquus* in Wisconsin, 98.
 Seton, Ernest E. T., nest and habits of the Connecticut Warbler (*Oporornis agilis*), 192.
Setophaga ruticilla, 175, 176, 217.
 Shalow, H., notes on '*Lanius cristatus*' and '*L. borealis*' of Nelson's 'Birds of Bering Sea and the Arctic Ocean,' 291.
 Sharpe, R. B., notice of his Catalogue of the Timeliidæ, 278; a note on the genus *Progne*, 367.
 Shearwater, Greater, 237, 380, 381.
 Slender-billed, 234.
 Sheldrake, American, 249.
 Hooded, 249.
 Red-headed, 249.
 Shoveller, 248.
 Shrike, Great Northern, 158, 244, 324.
 Loggerhead, 186, 193, 291.
 Shufeldt, R. W., notice of his paper on the Osteology of *Podiceps montanus*, 183.
Sialia arctica, 121.
 mexicana, 121.
 sialis, 133, 244.
Sitta europæa, 343.
 canadensis, 15, 89, 136, 157, 244.
 carolinensis, 135, 157, 244.
 carolinensis aculeata, 281.
 villosa, 281.
Siurus auricapillus, 13, 175, 176, 178, 179, 213, 295.
 motacilla, 121, 215.
 nævius, 175, 176, 178, 214, 286.
 nævius notabilis, 83.
 Skua, Buffon's, 97.
 Great, 238.
 Slade, E., a new element of diagnosis, 402.
 Snipe, Red-bellied, 32.
 Wilson's, 187, 248.
 Snowbird, Black, 9, 11, 13, 245.
 Eastern, 86.
 Solitaire, Townsend's, 100.
 Sparrow, Baird's, 286.
 Chipping, 122.
 English, 85, 87, 159, 188, 189, 287.
 Field, 6, 8, 70.
 Henslow's, 7.
 House, 87, 208, 373.
 Intermediate White-crowned, 100.
 Ipswich, 31, 292.
 Leconte's, 286.
 Mountain Song, 224.
 Savanna, 6, 11, 329.
 Song, 6, 8, 65, 70, 245.
 Swamp, 6, 7, 67.
 Titlark, 106.
 White-crowned, 330.
 White-throated, 11, 12, 122, 331.
 Yellow-winged, 7, 330.
Spatula platalea, 274.
Speotyto cunicularia, 30.
Sphyrapicus varius, 15, 179, 246, 295.
 varius nuchalis, 100.
Spinus lawrencii, 362.
 notatus, 362.
 pinus, 362.
 psaltria, 362.
 psaltria arizonæ, 362.
 psaltria mexicanus, 362.
 tristis, 362.
Spiza townsendi, 191.
Spizella breweri, 121.
 domestica, 175.
 montana, 177.

- Spizella monticola, 295.
 socialis, 122.
 Starling, Red-winged, 85.
 Stearns, W. A., notice of his 'Notes on the Natural History of Labrador,' 284.
 Steatornis caripensis, 307.
 Steganopus wilsoni, 33.
 tricolor, 367.
 Stejneger, L., *Dendrocopos purus*, a New Species of Woodpecker from Kamtschatka, 35; notice of his 'Contributions to the History of the Commander Islands. No. 1,' 81; on changes in ornithological nomenclature—a reply to critics, 114; notes on the genus *Acanthis*, 145; *Analecta ornithologica*: I, the occurrence of *Turdus aliciae* in the Palæarctic Region, 166; II, on the earliest available name of the American Titlark, 167; III, a brief review of the synonymy of the genus *Compsothlypis* (= *Parula*), 168; IV, on the earliest available name of the Cardinal Grosbeak, 171; V, more 'Ornithophilologicalities,' 172; notice of his 'Systematic Arrangement of the Turdidæ,' 181; *Analecta ornithologica* (Second Series): VI, on the Ptarmigans of Nelson's 'Birds of Bering's Sea,' etc., especially those belonging to the group *Attagen* Kaup, 225; VII, on some changes necessary in North American and European ornithological nomenclature, if generic appellations previously applied in botany be not rejected, 228; VIII, *Larus schistisagus*, a new species of Gull from the North Pacific, 231; IX, *Priocella tenuirostris* (Aud.) not a bird of Bering Sea or the Arctic Ocean, 233; X, on old and new generic names, 234; *Analecta Ornithologica* (Third Series): XI, note on Arctic *Lari*, 358; XII, *Chrysomitris* or *Spinus*? 360; XIII, on the systematic name of the American Hawk Owl, 362; XIV, on *Sterna nilotica* of Hasselquist, 364; XV, *Habia* against *Zamelodia*, 366; XVI, on the oldest available name of Wilson's Phalarope, 367; a new subspecies of Willow Grouse from Newfoundland, 369; notice of his paper on 'Trinomials in American Ornithology,' 381.
 Stelgidopteryx serripennis, 121.
 Stephens, F., collecting in the Colorado Desert—Leconte's Thrasher, 353.
 Stercorarius buffoni, 97, 237.
 pomarinus, 173.
 pomatorhinus, 173.
 skua, 238.
 Sterna, 144.
 anglica, 34, 364.
 antillarum, 294.
 cantiaca, 186.
 caspia, 34.
 fuliginosa, 59.
 magnirostris, 316.
 nilotica, 364.
 regia, 97.
 superciliaris, 316.
 Sturnus cinclus, 120
 Sthenelides melancorypha, 235.
 Strepsilas interpres, 173, 399.
 Strix canadensis, 362.
 caparoch, 362.
 funerea, 362.
 hudsonica, 363.
 Subarquatus, 142.
 Sula, 144, 173.
 Surnia funerea, 79.
 funerea ulula, 79.
 ulula, 363.
 ulula caparoch, 363.
 Swallow, Barn, 6, 325.
 Cliff, 6.
 White-bellied, 185, 325.
 Swan, Trumpeter, 184.
 Swift, 24.
 Chimney, 13, 161, 188, 246.
 Sylvia brasiliana, 170.
 eversmanni, 78, 227.
 pitiayumi, 170.
 plumbea, 170.
 pusilla, 170.
 torquata, 170.
 venusta, 170.
 Synthliborhamphus antiquus, 98.
 Syrnum cinereum, 286.
 nebulosum, 88.
 TALBOT, D. H., Night Herons and Rails in Dakota, 96.
 Tanager, Scarlet, 8, 9, 13, 64, 190, 244, 326.
 Tantalus, 83.
 loculator, 272.
 Tattler, Wandering, 79.
 Teal, Blue-winged, 248.
 Green-winged, 248.
 Telmatodytes palustris, 244.

- Tern, Black, 59.
 Caspian, 34.
 Gull-billed, 34, 365.
 Least, 294.
 Royal, 97.
 Sooty, 59.
- Tero-tero, 278.
- Thalassaëtus pelagicus, 82.
- Thalassoica glacialoides, 80.
- Thamnophilus argentinus, 21.
 cerulescens, 21.
- Theristicus melanopsis, 272.
- Thrasher, Brown, 6.
 Leconte's, 353, 355, 356.
- Thrasyaëtus, 41.
- Thrush, Bicknell's 58, 268.
 Brown, 132, 244.
 Golden-crowned, 67.
 Gray-cheeked, 130.
 Hermit, 1, 10, 11, 121, 131.
 Large-billed Water, 215.
 Olive-backed, 12, 13, 129, 243.
 Small-billed Water, 215.
 Wilson's, 6, 8, 9, 11, 129, 243.
 Wood, 6, 8, 9, 11, 128, 190.
- Thryothorus ludovicianus, 91, 137.
- Tijerita-Dormilon, 24.
- Tinnunculus sparverius, 247.
- Tit, Hudson's Bay, 157.
- Titlark, American, 209.
- Titmouse, Tufted, 134.
- Torcaz, 274.
- Totanus cinerascens, 115.
 flavipes, 315.
 glottis, 115.
 melanoleucus, 178, 315, 380.
- Towhee, Arctic, 286.
- Townsend, C. H., notice of his 'Notes on the Birds of Westmorland County, Penn.,' 184.
- Townsend, C. W., breeding of the Mockingbird near Boston, Mass., 192.
- Tringa canutus, 295, 307.
 fuscicollis 314.
 maculata, 314.
 merula, 120.
 vanellus, 120.
- Tringoides macularius, 175.
- Trochilus colubris, 123, 246.
- Troglodytes aëdon, 137, 244.
- Turdus aliciae, 130, 166.
 aliciae bicknelli, 58, 268.
 fuscescens, 129.
 migratorius, 87, 90, 127.
 mustelinus, 14, 128, 190.
 nanus, 131.
 pallasi, 15.
 pinicola, 181.
- Turdus swainsoni, 15.
 ustulatus swainsoni, 129.
- Turkey, Domestic, 187, 188.
 Wild, 187.
- Turner, M. H., *Vireo philadelphicus* in the Adirondack Region, 291.
- Turnstone, 229.
 Black, 229.
- Tyrannus carolinensis, 122, 245.
 intrepidus, 87.
- ULULA cinerea, 79, 246.
 cinerea lapponica, 79.
 flammeata, 362.
- Urinator, 117.
- Urophasianus, 141.
- Urubitinga, 140.
- VANELLUS, 141.
 capella, 120.
 cayennensis, 278.
- Vanneau, 120.
- Vireo, Blue-headed, 332.
 Philadelphia, 291.
 Red-eyed, 6, 8, 9, 11, 65, 217.
 Solitary, 122.
 Warbling, 6, 217.
 White-eyed, 6, 332.
 Yellow-throated, 6, 124, 332.
- Vireo atricapillus, 121.
 flavifrons, 332.
 gilvus, 218.
 noveboracensis, 179, 333.
 olivaceus, 217.
 philadelphicus, 291.
 solitarius, 333.
- Vireosylva olivacea, 176, 177, 178, 179.
- Vulture, Black, 286.
- WARBLER, Audubon's, 100.
 Bay-breasted, 213.
 Black-and-white Creeping, 210.
 Black-and-yellow, 11.
 Blackburnian, 13, 213.
 Black-capped Flycatching, 217.
 Black-poll, 213, 269.
 Black-throated Blue, 12, 213, 244.
 Black-throated Green, 9, 11, 13, 213.
 Blue-winged Yellow, 210, 213.
 Canadian Flycatching, 217.
 Cape May, 59, 213.
 Chestnut-sided, 6, 9, 213.
 Connecticut, 192.
 Golden Swamp, 290.
 Hooded, 216.
 Kirtland's, 389.

- Warbler, Mourning, 9, 12, 13.
 Nashville, 6, 7, 211.
 Prairie, 189.
 Summer, 212.
 Tennessee, 212.
 Worm-eating, 210.
 Yellow, 6, 8.
 Yellow Redpoll, 213.
 Yellow-rumped, 213.
 Yellow-throated, 189.
- Ward, C. W., notes on *Ardea wardi* Ridgw., 161.
- Wasseramsel, 120.
- Water-Thrush. See Thrush.
- Waxwing, Bohemian, 10, 244, 288, 294.
 Cedar, 244.
 Northern, 158.
- Welsh, F. R., odd nesting-site of the Great-crested Flycatcher, 391.
- Wheatear, 378, 398.
- Whippoorwill, 246.
- Widgeon, American, 394.
 European, 80.
- Wife, Old, 184.
- Willard, S. W., the migration of our winter birds, 221.
- Wilsonia bonapartii, 231.
 minuta, 231.
 mitrata, 231.
- Wilsonia pusilla, 231.
- Wintle, E. D., the Loggerhead Shrike (*Lanius ludovicianus*), 193.
- Woodcock, 87, 248.
- Woodpecker, Alaskan Three-toed, 164.
 Banded-backed Three-toed, 161.
 Black-backed Three-toed, 93, 161.
 Downy, 9, 93, 122, 161, 246.
 Golden-winged, 6.
 Hairy, 9, 12, 87, 161, 246.
 Ivory-billed, 286.
 Pileated, 9, 12, 246.
 Red-headed, 189, 246.
 Red-naped, 100.
 Yellow-bellied, 12, 86, 286.
- Wren, Carolina, 137, 286.
 House, 6, 85, 137, 244.
 Long-billed Marsh, 138.
 Short-billed Marsh, 59, 139.
 Winter, 12, 13, 138.
- YELLOW-THROAT, Maryland, 389.
- Young, C. J., notes on certain birds observed on a voyage from Liverpool to Quebec in September, 1883, 398.
- ZAMELODIA, 366.

ERRATA.

- Page 31, line 16, transpose the words *Extent* and *Length*.
 " 32, " 14, for 1881 read 1882.
 " 81, " 38, for Smidt read Smit.
 " 96, last word, for *grævia* read *naevia*.
 " 105, line 35, dele the words 'publication of the.'
 " 106, " 3, for of read on.
 " 185, " 6, add —C. H. M.
 " 208, " 4, for Picking read Picking.
 " 223, " 33, for collection read Museum.
 " " " 34, for Museum read collection.
 " 260, " 40, for 8 read 7.
 " 261, " 20, for 9 read 8.
 " 290, " 25, for mein read mien.
 " 303, " 36, for naturalist read naturalists.
 " 308, " 28, for Blackiston read Blakiston.
 " " " 46, for 1877 read 1887.
 " 406, " 9, (first column) for Astrigalinus read Astragalinus.