

yellow, heavily streaked with rufous brown; wings dark brown edged with yellow; tail-feathers brown, heavily marked with yellow on the inner webs and faintly edged with it on the outer.

Length, 4.75; wing, 2.35; tail, 2; tarsus, .80; bill, .30.

Habitat. St. Andrews Island.

A LIST OF THE BIRDS TAKEN BY MR. ROBERT HENDERSON, IN THE ISLANDS OF OLD PROVIDENCE AND ST ANDREWS, CARIBBEAN SEA, DURING THE WINTER OF 1886-87.*

BY CHARLES B. CORY.

Old Providence.

- Dendroica palmarum (*Gmel.*).
- Dendroica coronata (*Linn.*).
- Compsothlypis americana (*Linn.*).
- Seiurus noveboracensis (*Gmel.*).
- Seiurus noveboracensis notabilis (*Grinn.*).
- Seiurus aurocapillus (*Linn.*).
- Seiurus motacilla (*Vieill.*).
- Certhiola tricolor *Ridgw.*
- Vireo approximans *Ridgw.*
- Vireosylvia grandior *Ridgw.*
- Spiza americana (*Gmel.*).
- Euethia bicolor (*Linn.*).
- Tyrannus tyrannus (*Linn.*).
- Elainea cinerescens *Ridgw.*
- Lampornis hendersoni *nobilis.*
- Coccycus minor (*Gmel.*).
- Melopelia leucoptera (*Linn.*).
- Columba leucocephala *Linn.*
- Actitis macularia (*Linn.*).
- Ardea virescens (*Linn.*).
- Ardea cærulea (*Linn.*).
- Ardea tricolor ruficollis (*Gosse*).
- Fregata aquila (*Linn.*).

* [An author's edition of 250 copies of this paper was published May 28, 1887.—EDD.]

Sula piscator (*Linn.*).

Puffinus auduboni *Finsch.*

St. Andrews.

Mimus magnirostris nobis.

Galeoscoptes carolinensis (*Linn.*).

Mniotilla varia (*Linn.*).

Dendroica flava *nobis.*

Seiurus noveboracensis (*Gmel.*).

?**Certhiola tricolor** *Ridgw.*

Vireosylvia canescens nobis.

Vireo noveboracensis (*Gmel.*).

Euethia bicolor (*Linn.*).

Icterus lawrencii nobis.

?**Elainea martinica** (*Linn.*)

Sphyrapicus varius (*Linn.*).

Ceryle alcyon (*Linn.*).

Engyptila neoxena nobis.

Actitis macularia (*Linn.*).

Ardea virescens (*Linn.*).

Ardea tricolor ruficollis (*Gosse*).

Fregata aquila (*Linn.*).

Sula piscator (*Linn.*).



BIRDS OF TOM GREEN AND CONCHO COUNTIES,
TEXAS.

BY WILLIAM LLOYD.

THE present paper deals principally with the avi-fauna of the valleys of the Concho River and its tributaries east to the Colorado River. It also includes the birds of the plains west of the Pecos River, and north to the Texas and Pacific Railroad, and some few noted incidentally south in Crockett and Edwards Counties, and in Nueces Cañon. The district has a general and nearly equal elevation of nearly 2000 feet above the sea-level, and