

those from neighboring islands. The wings and tail are very dark, almost black, but my cabinet also contains specimens from Porto Rico which are difficult to distinguish from birds taken in Cuba and San Domingo.

Mimus polyglottos portoricensis BRYANT.

Mimus polyglottus var. *portoricensis* BRYANT, Proc. Boston Soc. Nat. Hist. XI, p. 68 (1866).

Mimus polyglottus GUNDLACH, Anales de la Soc. Esp. de Hist. Nat. VII, p. 173 (1878).

HABITAT.—Porto Rico.

A COLLECTION OF BIRDS TAKEN BY CYRUS S
WINCH IN THE ISLANDS OF ANGUILLA, AN-
TIGUA, AND ST. EUSTATIUS, WEST INDIES,
DURING APRIL, MAY, JUNE, AND A
PART OF JULY, 1890.

BY CHARLES B. CORY.

ANGUILLA.

Podilymbus podiceps (<i>Linn.</i>).	Elænea martinicana (<i>Linn.</i>).
Gallinula galeata (<i>Licht.</i>).	Euetheia bicolor (<i>Linn.</i>).
Fulica caribæa (<i>Ridgw.</i>).	Loxigilla noctis (<i>Linn.</i>).
Himantopus mexicanus <i>Müll.</i>	Cœreba bartholemica? (<i>Sparrm.</i>).
Falco caribbæarum <i>Gmel.</i>	Cichlherminia fuscata densirostris (<i>Vieill.</i>).
Ceryle alcyon (<i>Linn.</i>).	
Tyrannus dominicensis (<i>Gmel.</i>).	Galeoscoptes carolinensis (<i>Linn.</i>).

ANTIGUA.

Larus atricilla (<i>Linn.</i>).	Pelecanus fuscus <i>Linn.</i>
Sterna sandvicensis acuflavida <i>Cabot.</i>	Gallinula galeata (<i>Licht.</i>).
Sterna dougalli <i>Mout.*</i>	Fulica caribæa <i>Ridgw.</i>
Sterna antillarum (<i>Less.</i>).	Himantopus mexicanus <i>Müll.</i>

* Specimens of this species taken in June have the basal half of the bill red, and in three specimens the red color extends more than half the length of the bill. In all these birds the black cap is complete. In specimens taken in September, which have the black cap imperfect or mixed with white and gray, the bill is entirely black.

<i>Symphemia semipalmata</i> (<i>Gmel.</i>).	<i>Bellona exilis</i> (<i>Gmel.</i>).
<i>Totanus melanoleucus</i> (<i>Gmel.</i>).	<i>Eulampis holosericeus</i> (<i>Linn.</i>).
<i>Colinus virginianus</i> (<i>Linn.</i>).	<i>Tyrannus dominicensis</i> (<i>Gmel.</i>).
<i>Columba leucocephala</i> <i>Linn.</i>	<i>Elænea martinica</i> (<i>Linn.</i>).
<i>Zenaida zenaida</i> (<i>Bonap.</i>).	<i>Euetheia bicolor</i> (<i>Linn.</i>).
<i>Geotrygon mystacea</i> (<i>Temm.</i>).	<i>Loxigilla noctis</i> (<i>Linn.</i>).
<i>Columbigallina passerina</i> (<i>Linn.</i>).	<i>Piranga erythromelas</i> <i>Vieill.</i>
<i>Buteo latissimus</i> (<i>Wils.</i>).*	<i>Vireo calidris</i> (<i>Linn.</i>).
<i>Falco caribbæarum</i> (<i>Gmel.</i>).	<i>Cœreba dominicana</i> (<i>Taylor</i>).
<i>Speotyto amaura</i> <i>Lawr.</i>	<i>Dendroica ruficapilla</i> (<i>Gmel.</i>).
<i>Coccyzus minor</i> (<i>Gmel.</i>).	<i>Allenia montana</i> (<i>Lafr.</i>).
<i>Coccyzus americanus</i> (<i>Linn.</i>).	<i>Cichlherminia fuscata densirostris</i> (<i>Vieill.</i>).

ST. EUSTATIUS.

<i>Phaëthon flavirostris</i> <i>Brandt.</i>	<i>Vireo calidris</i> (<i>Linn.</i>).
<i>Ardea cærulea</i> <i>Linn.</i>	<i>Progne dominicensis</i> (<i>Gmel.</i>).
<i>Columba corensis</i> <i>Gmel.</i>	<i>Cœreba dominicana</i> (<i>Taylor</i>).
<i>Zenaida zenaida</i> (<i>Bonap.</i>).	<i>Mniotilta varia</i> (<i>Linn.</i>).
<i>Falco caribbæarum</i> <i>Gmel.</i>	<i>Dendroica ruficapilla</i> (<i>Gmel.</i>).
<i>Bellona exilis</i> (<i>Gmel.</i>).	<i>Compsothlypis americana</i> (<i>Linn.</i>).
<i>Eulampis holosericeus</i> (<i>Linn.</i>).	<i>Dendroica discolor</i> (<i>Vieill.</i>).
<i>Tyrannus dominicensis</i> (<i>Gmel.</i>).	<i>Setophaga ruticilla</i> (<i>Linn.</i>).
<i>Euetheia bicolor</i> (<i>Linn.</i>).	<i>Allenia montana</i> (<i>Lafr.</i>).
<i>Loxigilla noctis</i> (<i>Linn.</i>).	<i>Cichlherminia fuscata densirostris</i> (<i>Vieill.</i>).

A LIST OF THE BIRDS COLLECTED IN THE
ISLANDS OF ST. CROIX AND ST. KITTS, WEST
INDIES, DURING MARCH AND APRIL, AND
IN GUADELOUPE DURING AUGUST,
SEPTEMBER, AND OCTOBER, 1890.

BY CHARLES B. CORY.

ST. CROIX.

<i>Gallinula galeata</i> (<i>Licht.</i>).	<i>Totanus flavipes</i> (<i>Gmel.</i>).
<i>Ardea virescens</i> <i>Linn.</i>	<i>Columba leucocephala</i> <i>Linn.</i>
<i>Nycticorax violaceus</i> (<i>Linn.</i>).	<i>Columba corensis</i> <i>Gmel.</i>

* A rather small pale form, resident in Antigua, and quite abundant.