

Stream, and having no harbor that vessels drawing more than seven feet of water can enter. We stopped here to meet Mr. Charles Washburn, one of my collectors, who had been staying on the Islands for about three weeks. Very few resident species were obtained, but it is more than probable that a number of Bahama forms occur there which were not observed by us, as the time spent on these Islands was much too short to make anything like a thorough investigation.

<i>Sterna maxima</i> <i>Bodd.</i>	<i>Vireo crassirostris</i> <i>Bryant.</i>
<i>Pelecanus fuscus</i> <i>Linn.</i>	<i>Vireo altiloquus barbatulus</i> (<i>Cab.</i>). <i>Cæreba bahamensis</i> (<i>Reich.</i>). <i>Mniotilla varia</i> (<i>Linn.</i>). <i>Helinaia swainsoni</i> <i>Aud.</i>
<i>Fregata aquila</i> (<i>Linn.</i>). <i>Ardea herodias</i> <i>Linn.</i>	<i>Compsothlypis americana</i> (<i>Linn.</i>). <i>Dendroica tigrina</i> (<i>Gmel.</i>). <i>Dendroica coronata</i> (<i>Linn.</i>). <i>Dendroica dominica</i> (<i>Linn.</i>). <i>Dendroica palmarum</i> (<i>Gmel.</i>). <i>Dendroica discolor</i> (<i>Vieill.</i>). <i>Geothlypis trichas</i> (<i>Linn.</i>). <i>Setophaga ruticilla</i> (<i>Linn.</i>). <i>Seiurus motacilla</i> (<i>Vieill.</i>). <i>Seiurus aurocapillus</i> (<i>Linn.</i>). <i>Polioptila cærulea</i> (<i>Linn.</i>). <i>Galeoscoptes carolinensis</i> (<i>Linn.</i>). <i>Mimus polyglottos orpheus</i> (<i>Linn.</i>).
<i>Ardea virescens bahamensis</i> (<i>Brewster</i>). <i>Ægialitis vocifera</i> (<i>Linn.</i>). <i>Columbigallina passerina</i> (<i>Linn.</i>). <i>Columba leucocephala</i> <i>Linn.</i> . <i>Crotophaga ani</i> <i>Linn.</i> . <i>Ceryle alcyon</i> (<i>Linn.</i>). <i>Sphyrapicus varius</i> (<i>Linn.</i>). <i>Doricha evelynæ</i> (<i>Bourc.</i>). <i>Tyrannus dominicensis</i> (<i>Gmel.</i>). <i>Agelaius phœniceus bryanti</i> <i>Ridgw.</i> <i>Euethia bicolor</i> (<i>Linn.</i>). <i>Ammodramus savannarum passerinus</i> (<i>Wils.</i>).	

LIST OF THE BIRDS COLLECTED BY C. L. WINCH
IN THE CAICOS ISLANDS AND INAGUA, BA-
HAMAS, DURING JANUARY AND FEBRUARY,
AND IN ABACO, IN MARCH, 1891.

BY CHARLES B. CORY.

CAICOS ISLANDS.

Dafila bahamensis (*Linn.*). — Not uncommon; breeds on the Caicos Islands.

Aythya affinis (*Eyton*).

Ardea rufescens *Bodd.*—Several specimens of both the white and colored plumage of this species.

Ardea tricolor ruficollis (*Gosse*).
Sympemia semipalmata (*Gmel.*).
Hæmatopus palliatus (*Temm*).
Columbigallina passerina (*Linn*).
Columba leucocephala *Linn.*.
Zenaida zenaida (*Linn.*).
Pandion haliaëtus carolinensis (*Gmel.*).
Coccyzus minor maynardi *Ridgw.*
Crotophaga ani *Linn.*.
Sphyrapicus varius (*Linn.*).
Doricha evelynæ (*Bourc.*).

Tyrannus magnirostris D'Orb.—Three specimens taken. It has not previously been recorded north of Inagua.

Loxigilla violacea (*Linn.*).
Euetheia bicolor (*Linn.*).
Vireo crassirostris (*Bryant*).
Cæreba bahamensis (*Reich.*).
Compsothlypis americana (*Linn.*).
Dendroica petechia gundlachi *Baird*.
Dendroica coronata (*Linn.*).

Dendroica kirtlandii *Baird*—Two specimens taken. I believe it has no previous record so far south. The naturalists of the 'Albatros' expedition found it not uncommon in Rum Cay.

Dendroica palmarum (*Gmel.*).
Dendroica discolor (*Vieill.*).
Sciurus aurocapillus (*Linn.*).
Polioptila cærulea (*Linn.*).
Mimus gundlachi *Cab.*
Margarops fuscatus (*Vivill.*).

INAGUA.

Ardea herodias *Linn.*.
Sphyrapicus varius (*Linn.*).
Doricha lyrura *Gould*.
Tyrannus magnirostris D'Orb.
Myiarchus sagræ *Gundl.*

Loxigilla violacea (*Linn.*).—Mr. Winch writes me he believes the adult female of this species assumes a black plumage similar to the male. A number of black specimens taken by him proved on dissection to be females.

Euetheia bicolor (*Linn.*).
Vireo crassirostris (*Bryant*).
Cæreba bahamensis (*Reich.*).—Birds from Inagua have larger bills than those which occur on the northern Bahama Islands.
Dendroica tigrina (*Gmel.*).
Dendroica petechia gundlachi *Baird*.
Dendroica coronata (*Linn.*).

Dendroica dominica (*Linn.*).

Dendroica discolor (*Vieill.*).

Seiurus aurocapillus (*Linn.*).

Polioptila cærulea (*Linn.*).

Mimus gundlachi *Cab.*

Mimus polyglottos elegans (*Sharpe*). — Since writing the paper on *Mimus polyglottos* and *M. orpheus* which appeared in the January 'Auk' (p. 45) I have received some fifty specimens of the small *Mimus* from Inagua. With this additional material for comparison I find that a large majority of the Inagua specimens are somewhat smaller than *orpheus* or *polyglottos* and have the primary coverts nearly and often completely covering the white on the quills. These differences are not constant but are probably sufficient to justify subspecific separation.

Margarops fuscatus (*Vieill.*).

A box of birds lately received from Inagua contained two species not observed by Mr. Winch. These are :—

Dendroica striata (*Forst.*). — Six specimens, April 23 to May 2.

Dendroica cærulescens (*Gmel.*). — Nine specimens, April 22 to May 1.

ABACO.

Ægialitis semipalmata *Bonap.*

Columbigallina passerina (*Linn.*).

Cathartes aura (*Linn.*).

Crotophaga ani *Linn.*

Ceryle alcyon (*Linn.*).

Dryobates villosus maynardi

Ridgw.

Doricha evelynæ (*Bourc.*).

Sporadinus ricordi (*Gerv.*).

Myiarchus sagræ *Gundl.*

Pitangus bahamensis *Bryant.*

Loxigilla violacea (*Linn.*).

Euetheia bicolor (*Linn.*).

Spindalis zena townsendi *Ridgw.*

Vireo crassirostris (*Bryant*).

Vireo altiloquus barbatulus (*Cab.*).

Cæreba bahamensis (*Reich*).

Mniotilla varia (*Linn.*).

Dendroica kirtlandi *Baird.*

Dendroica petechia gundlachi
Baird.

Dendroica discolor (*Vieill.*).

Dendroica coronata (*Linn.*).

Dendroica dominica (*Linn.*).

Geothlypis trichas (*Linn.*).

Geothlypis tanneri *Ridgw.*

Seiurus noveboracensis (*Gmel.*).

Seiurus aurocapillus (*Linn.*).

Polioptila cærulea *Linn.*

Galeoscoptes carolinensis (*Linn.*).

Mimus polyglottos orpheus (*L.*).

Mimocichla plumbea (*Linn.*).