

LIST OF THE BIRDS OBTAINED BY MR. C. S. WINCH
ON THE ISLANDS OF GREAT BAHAMA AND
ABACO, BAHAMA ISLANDS, DURING JUNE,
AND ON ELEUTHERA IN JULY, 1891.

BY CHARLES B. CORY.

GREAT BAHAMA.

<i>Ægialitis wilsonia</i> (<i>Ord.</i>).	<i>Geothlypis rostratus tanneri</i> (<i>Ridg.</i>)
<i>Geotrygon martinica</i> (<i>Gmel.</i>).	<i>Dendroica petechia gundlachi</i> <i>Bd.</i>
<i>Chordeiles minor</i> <i>Cab.</i>	<i>Dendroica vigorsii</i> (<i>Aud.</i>).
<i>Dryobates villosus maynardi</i> <i>Ridg.</i>	<i>Dendroica pityophila bahamensis</i> <i>Cory.</i>
<i>Sporadinus ricordi</i> (<i>Gerv.</i>).	<i>Cæreba bahamensis</i> (<i>Reich.</i>).
<i>Pitangus bahamensis</i> <i>Bryant.</i>	<i>Sitta pusilla</i> <i>Lath.</i> —First record of this species in the Bahamas.
<i>Myiarchus sagræ</i> <i>Gundl.</i>	<i>Vireo crassirostris</i> <i>Bryant.</i>
<i>Tyrannus dominicensis</i> (<i>Gmel.</i>).	<i>Mimus polyglottos orpheus</i> (<i>Linn.</i>). <i>Mimocichla plumbea</i> (<i>Linn.</i>).
<i>Agelaius phœniceus bryanti</i> <i>Ridg.</i>	
<i>Loxigilla violacea</i> (<i>Linn.</i>).	
<i>Spindalis zena townsendi</i> <i>Ridgw.</i>	
<i>Callichelidon cyaneoviridis</i> <i>Bryant.</i>	

ABACO.

<i>Buteo borealis</i> (<i>Gmel.</i>).	<i>Loxigilla violacea</i> (<i>Linn.</i>).
<i>Zenaidura macroura</i> (<i>Linn.</i>).	<i>Spindalis zena townsendi</i> <i>Ridgw.</i>
<i>Chordeiles minor</i> <i>Cab.</i>	<i>Vireo crassirostris</i> <i>Bryant.</i>
<i>Sporadinus ricordi</i> (<i>Gerv.</i>).	<i>Callichelidon cyaneoviridis</i> <i>Bryant.</i>
<i>Doricha evelynae</i> (<i>Bourc.</i>).	<i>Geothlypis rostratus tanneri</i> (<i>Ridg.</i>)
<i>Centurus blakei</i> <i>Ridgw.</i>	<i>Dendroica vigorsii</i> (<i>Aud.</i>).
<i>Dryobates villosus maynardi</i> <i>Ridg.</i>	<i>Dendroica pityophila bahamensis</i> <i>Cory.</i>
<i>Myiarchus sagræ</i> <i>Gundl.</i>	<i>Dendroica dominica</i> (<i>Linn.</i>).
<i>Contopus bahamensis</i> <i>Bryant.</i>	<i>Cæreba bahamensis</i> (<i>Reich.</i>).
<i>Pitangus bahamensis</i> <i>Bryant.</i>	<i>Polioptila cærulea</i> (<i>Linn.</i>).
<i>Tyrannus dominicensis</i> (<i>Gmel.</i>).	<i>Mimus polyglottos orpheus</i> (<i>Linn.</i>).
<i>Icterus northropi</i> <i>Allen.*</i>	<i>Mimocichla plumbea</i> (<i>Linn.</i>).
<i>Agelaius phœniceus bryanti</i> <i>Ridg.</i>	

ELEUTHERA.

<i>Colymbus dominicus</i> <i>Linn.</i>	<i>Speotyto cunicularia floridana</i>
<i>Ardea virescens</i> <i>Linn.</i>	<i>Ridgw.</i>
<i>Totanus flavipes</i> (<i>Linn.</i>).	<i>Geotrygon martinica</i> (<i>Gmel.</i>).

* Several specimens taken, all being slightly darker yellow than the type. Not previously known except from Andros Island.

Saurothera bahamensis <i>Bryant.</i>	Loxigilla violacea (<i>Linn.</i>).
Doricha evelynæ (<i>Bourc.</i>).	Euetheia bicolor (<i>Linn.</i>).
Coccyzus minor maynardi <i>Ridgw.</i>	Vireo crassirostris <i>Bryant.</i>
Myiarchus sagræ <i>Gundl.</i>	Geothlypis coryi <i>Ridgw.</i>
Tyrannus dominicensis (<i>Gmel.</i>).	Cœreba bahamensis (<i>Reich.</i>).
Contopus bahamensis <i>Bryant.</i>	Mimus gundlachi <i>Cab.</i>
Spindalis zena stejnegeri <i>Cory.</i>	Mimocichla plumbea (<i>Linn.</i>).

LIST OF BIRDS COLLECTED ON THE ISLAND OF
INAGUA, BAHAMA ISLANDS, FROM MAY 1 TO
JULY 10, 1891.

BY CHARLES B. CORY.

Phaethon flavirostris <i>Brandt.</i>	Speotyto cunicularia bahamensis
Sterna antillarum (<i>Less.</i>).	<i>Cory.</i>
Sterna dougalli (<i>Montag.</i>).	Chrysotis leucocephala (<i>Linn.</i>).
Sterna sandvicensis acuflavida	Doricha lyrura <i>Gould.</i>
(<i>Cabot.</i>)	Coccyzus americanus (<i>Linn.</i>).
Sterna maxima <i>Bodd.</i>	Coccyzus minor maynardi <i>Ridgw.</i>
Nycticorax violaceus (<i>Linn.</i>).	Myiarchus sagræ <i>Gundl.</i>
Ardea virescens <i>Linn.</i>	Tyrannus dominicensis (<i>Gmel.</i>).
Ardea candidissima <i>Gmel.</i>	Dolichonyx oryzivorus (<i>Linn.</i>).
Ajaja ajaja (<i>Linn.</i>).	Loxigilla violacea (<i>Linn.</i>).
Tringa minutilla <i>Vieill.</i>	Euetheia bicolor (<i>Linn.</i>).
Ereunetes pusillus (<i>Linn.</i>).	Vireo altiloquus barbatulus (<i>Cab.</i>).
Ægialitis semipalmata <i>Bonap.</i>	Vireo crassirostris <i>Bryant.</i>
Ægialitis wilsonia (<i>Ord.</i>).	Cœreba bahamensis (<i>Reich.</i>).
Ægialitis vocifera (<i>Linn.</i>).	Seiurus aurocapillus (<i>Linn.</i>).
Ægialitis meloda (<i>Ord.</i>).	Seiurus noveboracensis (<i>Gmel.</i>).
Symphearia semipalmata (<i>Gmel.</i>).	Setophaga ruticilla (<i>Linn.</i>).
Arenaria interpres (<i>Linn.</i>).	Geothlypis trichas (<i>Linn.</i>).
Hæmatopus palliatus <i>Temm.</i>	Dendroica striata (<i>Forst.</i>).
Himantopus mexicanus (<i>Müll.</i>).	Dendroica palmarum (<i>Gmel.</i>).
Zenaida zenaïda (<i>Bonap.</i>).	Dendroica petechia gundlachi <i>Bd.</i>
Melopelia leucoptera (<i>Linn.</i>).	Polioptila cærulea (<i>Linn.</i>).
Pandion haliaetus carolinensis	Mimus polyglottos elegans (<i>Sharpe</i>)
(<i>Gmel.</i>).	Mimus gundlachi <i>Cab.</i>
	Margarops fuscatus (<i>Vieill.</i>).