

liar lark song, but was unable to investigate the matter. Later I wrote Mr. Vernon Bailey and Mr. B. H. Dutcher, both of whom had been observing the intergradations of plumage and habits in Kansas of these two birds, and I asked of them if they had observed any birds with a blended voice. Neither had; and Mr. Bailey was kind enough to ask other ornithologists at the Smithsonian Institution concerning the matter. None had noticed any intergrading.

But recently (March 9, 1896) I was in the same region of my former duck hunt and heard distinctly a Western Lark's song. It was fairly typical but too highly pitched. A mile away I heard another that was deeply liquid and gurgling—in fact quite typical of *S. m. neglecta*, but only a fourth of a mile away I heard one whose song was that of *S. magna* except that it had near the middle a rather highly pitched gurgle. It did not have, in its many repetitions, any downward ending as the other two birds had. Neither was the entire song so highly pitched as that of the typical *S. magna*—scores of which were singing around it.

To my mind here was a distinct intergradation—perhaps the result of hybridism. The region was in Audrain County, Mo., about fifteen miles southwest of Mexico, Mo., and about one fourth of a mile north of the divide between the water-sheds of the Missouri and Mississippi Rivers.

I had shot the *S. m. neglecta* once before in this county further eastward.

This is the only region near me where *S. m. neglecta* is found so far as I know, though it comes into Southwest Missouri. I might add that I am well acquainted with the song of *S. m. neglecta*, having heard it in Kansas, Colorado and S. Dakota.—JAMES NEWTON BASKETT, *Mexico, Mo.*

Pinicola enucleator at Worcester, Mass.—A flock of sixteen Pine Grosbeaks, containing a few bright males, was reported in the northern part of this city Jan. 15, 1896. For a little over a month the Grosbeaks were seen in different parts of the city, one or two or a half dozen at a time, feeding upon mountain-ash berries and seeds of maple and Scotch elm. The last seen was on Feb. 21. But few bright males were seen, the largest number being reported on Feb. 19, when a flock of fifteen contained "several brilliant specimens."—HELEN A. BALL, *Worcester, Mass.*

Evening Grosbeak in Southern Wisconsin.—On January 21, 1896, while passing an old, deserted cemetery on the outskirts of Delavan (Wis.), I heard lively chattering among the large 'evergreen' trees within the graveyard and knew at once that it was the note of some bird new to me. On watching closely I soon saw a beautiful male Grosbeak (*Coccothraustes vespertinus*) appear on the outside of the tree, and later a couple more and two or three of the plainer clothed females. There

must have been at least a dozen in the tree. They were seen every day in good numbers up to about the middle of March, and throughout the remaining part of the winter a small flock of ten or a dozen were constant residents of the immediate vicinity of the above mentioned cemetery. This flock was last seen on March 30. It is their first occurrence in this locality to my knowledge, and I find no one who ever remembers seeing the bird here before. Pine Grosbeaks (*Pinicola enucleator*) were also observed in the county during December last.—N. HOLLISTER, *Delavan, Wisc.*

Zonotrichia albicollis and Mniotilta varia at Pasadena, Cal.—On Nov. 21, 1894, while collecting sparrows in a large blackberry patch just inside the western limits of Pasadena I shot an immature female White-throated Sparrow from a flock of *Z. coronata*, thus adding another record for this species from California.

Early in the morning of Oct. 8, 1895, I shot an immature female Black-and-white Warbler in the Arroyo Seco just west of Pasadena. The bird was at the time alone, and apparently as much at home in Southern California as she would have been east of the Rocky Mountains. So far as I can ascertain this is the second record of this species from California.—HORACE A. GAYLORD, *Pasadena, Cal.*

The Wintering of the Towhee at Longwood, Massachusetts.—I am glad to be able to report the following information in regard to the Towhee (*Pipilo erythrophthalmus*) noted December 25, 1895, at Longwood (see Auk, Vol. XIII, p. 178).

Mr. Henry Vose Greenough, who saw the Towhee with me on Christmas Day, reported to me having seen on March 23, 1896, a male Towhee about a brush pile, some one hundred and fifty yards from the spot where we had noted the one in December. On March 24 I went with him to this place and in a neighboring hemlock hedge we found Pipilo.

The brush pile is on the edge of an estate, only a few hundred feet from a stable, pig-sty and hen yard, where food and protection from the winter weather were easily accessible. When we started the Towhee on the 24th he flew straight for the hen yard and then being pursued, to another hemlock hedge leading us in a circuit back to the brush pile.

I believe there can be little doubt that this is our Christmas Towhee, which had wintered here, for the following reasons, viz.: Protection and food supply at hand; a male bird, as was the former one; in practically the same locality, and because it is exceedingly unlikely that a single bird would migrate northward fully a month in advance of its fellows.

We have not noted this bird during January and February, though we both have covered the neighboring ground almost daily, because the place is just on the edge of this little patch of woodland and the Towhee evidently never wandered far from his brush pile and the farm yard.