

Another Species Added to the Avifauna of South Carolina.— On the afternoon of December 15, I heard that a strange bird had been seen, since the morning of the 9th, around Chisolm's Mill on the Ashley River at the foot of Tradd Street, Charleston, and on going down on the morning of the 16th, I shot the bird, after a chase of ten minutes, as it lit on a side fence. It proved to be an Arkansas Kingbird (*Tyrannus verticalis*) a new species for this state. The bird is a young male in somewhat frayed plumage. The flame-patch on the crown is very indistinct.

On dissection I was very much surprised to find that the stomach contained, besides insect remains, a fiddler crab and about a dozen seeds. The winter dearth of insect life probably forced the bird to resort to this unusual food.

In coming to South Carolina this Kingbird had wandered one thousand miles east of its regular route. This record is the seventh east of the Mississippi and the southernmost, by over 400 miles, of the four Atlantic state records. The skin is now in the Charleston Museum.— E. BURNHAM CHAMBERLAIN, *Charleston, S. C.*

The Arkansas Kingbird (*Tyrannus verticalis*) in Rhode Island.— Last fall I spent at Watch Hill, R. I. The migration reached its height on September 24. I was working towards a small swamp, trying to count the birds, as the trees, bushes and grass were literally swarming with them. A large flycatcher suddenly flew out of an oak tree in front of me, and settled on a dead branch near the ground about 75 feet away. While on the wing it looked exactly like a Kingbird, and as this would have been an uncommonly late date I hurried up to get a closer view with my binoculars. What was my surprise when I discovered that my Kingbird had bright yellow underparts and no white band on the tip of its tail. There was no doubt but that it was an Arkansas Kingbird. The slate gray back with a very faint tinge of olive and the absence of any rufous on the tail distinguished it from the Crested Flycatcher, not to mention that its shape and flight and habits were totally different. I watched it for half an hour, several times getting within very close range. This is the first record for the state of Rhode Island.— LUDLOW GRISCOM, *New York City.*

Capture of *Myiarchus crinitus* (Linn.) in Eastern Cuba.— A specimen of the Crested Flycatcher, in fine condition, was shot on February 18, 1913, in the woods at the San Carlos Plantation at Guantánamo, Oriente Province. This is the fourth record for the Island, and the first for eastern Cuba, as Gundlach's three records are from the western end.— CHAS. T. RAMSDEN, *Guantanamo, Cuba.*

An Unusual Observation on the Crow (*Corvus brachyrhynchos*) at Lubec, Maine.— I have seen non-raptorial birds of various species attack other birds in self defense, or in defense of their nests and young, but

today (December 31, 1913) while on a tour of the local places where birds are to be found in winter, I made an observation new to me, and while actually interesting and thrilling, it was exasperating and disappointing. While walking along a stretch of sandy beach with a companion we suddenly flushed a belated shore bird, without having time to bring our glasses to bear upon it; from the shape of its wing, mode of flight and general coloration we were quite sure that it was a Sanderling. The bird took a course directly ahead of us, and along the water's edge; so we followed hoping to positively identify it.

Three hundred yards in advance of us, sitting on the beach was a flock of about two hundred Crows quietly basking in the sun. As soon as our bird drew near them it swerved out over the water, when one of the Crows rose and flew directly at it.

The smaller bird dodged repeated attacks and was drawing ahead, when another Crow at the farther end of the flock, seeing the escape flew out to intercept it and turned the Sanderling (?) back, giving the first assailant an opportunity to swoop down and deliver a telling blow with its wing. The smaller bird fell, but recovered itself before touching the water. Then to avoid its two assailants it was obliged to fly towards the land and directly over the large flock of Crows, when about twenty of their number rose in a bunch and bore the unfortunate to the beach, all endeavoring to get at it at the same time. As soon as we discovered the little fellow's real danger, we ran and shouted, but they did not heed us until close upon them, when they flew off leaving nothing but a small piece of breast with feathers attached, the body being carried away.

We followed them up from one place of alighting to another hoping to find a wing or some part of the body but without success.—CLARENCE H. CLARK, *Lubec, Maine.*

Starlings (*Sturnus vulgaris*) at Providence, R. I.—It may be of interest to the readers of 'The Auk,' in following the movements of the Starling to know they have appeared in Providence, R. I. My business requiring my presence in this city for a few weeks, on the morning of November 23, I started out with my field glass to see what could be seen here in the line of bird life. In passing through Swan Point Cemetery I was not a little surprised to encounter a flock of about forty Starlings, many of them in full summer plumage; they were constantly in song, many times singing notes quite like our Bluebird. I have had other excellent observations of them covering a date as late as December 6.

In the same section on December 1, 1912, I found two Mockingbirds but have failed to record them this season.—E. E. CADUC, *Boston, Mass.*

Starlings (*Sturnus vulgaris*) in Cambridge, Mass.—A flock of Starlings numbering about sixty birds appeared at Fresh Pond in Cambridge, in late November, 1913. They were first seen by me on the 27th day and have been successively observed up to the middle of February, the time of