

leaving the other undisturbed. The bird collected on November 16, while lacking the spotting on the abdomen and ventral region, has a much lengthened appendage on one secondary feather while the other is profusely spotted on the abdomen.

My belief is that the albinistic birds are descended from an albinistic parent for in all these years that I have been observing these abnormal birds I have always left undisturbed one or more that plainly showed albinism and each season from one to four birds would appear with the greatest regularity at the *same* place showing that this strain is descended from the birds left undisturbed by me from year to year. This makes 24 uninterrupted years that I have observed these birds.—ARTHUR T. WAYNE, *Mount Pleasant, S. C.*

**Bohemian Waxwing at Jackson, Mich.**—On March 11, 1923, in company with Randal McCain I saw two flocks of Bohemian Waxwings, one of thirty, the other of sixty individuals. These birds stayed in the vicinity of Jackson until early in May in broken companies, sometimes feeding on the American holly placed in winter porch boxes, sometimes on the berries of vines and shrubs throughout the city.—EDITH K. FREY, *Jackson, Mich.*

**White-eyed Vireo at Madison, Wis.**—A White-eyed Vireo (*Vireo griseus griseus*) associated with Golden-crowned Kinglets, was observed in a densely leaved seedling apple tree on October 21, 1923. On shooting, the bird appeared to have vanished completely. While attempting to be philosophical, a fluttering object was by chance observed to strike the ground at a considerable distance on my right. The potential leaf proved to be an immature male Vireo of the above species.

While it is useless to speculate on the presence of this always rare Wisconsin species at so late a season, its presence probably represents a case, recognized in recent years as not uncommon, of immature birds wandering north of the breeding grounds in autumn.—A. W. SCHORGER, *Madison, Wis.*

**Philadelphia Vireo in the New York City Region.**—In my recently published 'Handbook' to the birds of this region I discoursed at some length on the extreme local rarity of this Vireo. Two days after the appearance of this book or to be exact, on September 16, 1923, I discovered one on the "Point" of the Ramble in Central Park, New York City, in some low bushes where I had found one two years previously. It was very leisurely in its movements, and was only 25 feet away at about the level of my eyes when I spied it. It objected strongly to my presence, and scolded me harshly, gradually working nearer as it did so. A big wave of migrants had arrived overnight, and the scolding attracted a crowd of Warblers. At one time an adult male Tennessee Warbler was less than two feet from the Vireo, affording a faultless opportunity for comparison.