

Dendragapus fuliginosus sierrae (Grinnell).

Dendragapus fuliginosus sitkensis (Swarth).

Dendragapus fuliginosus howardi (Dickey and van Rossem).

Much additional work is required before the ranges of the seven forms can be exactly defined. The separation of the bird inhabiting Vancouver Island may be warrantable, but first the question of dichromatism in the females, giving a gray and a rufous phase, must be settled.—ALLAN BROOKS, *Okanagan Landing, B. C.*

Rare Michigan Records. *Numenius hudsonicus*.—HUDSONIAN CURLEW.—A fine adult male, Hudsonian Curlew, was taken at Whitefish Point, Chippewa County, on September 24, 1928, by Miss Curren Hawkins, and sent to the Museum in the flesh. This species has never been common in Michigan, and authentic records are few. The last spring record was on May 30, 1925, when an adult bird was taken at Newberry, Luce County, and sent to the Museum. This species occurs as a rare migrant in Michigan.

Pelecanus erythrorhynchos. WHITE PELICAN.—An immature male was taken at Oscoda by fishermen on September 25, 1928. It was in a very poor flesh when sent to the Museum, where it was made into a study skin.

An adult bird was taken alive near Muskegon, Muskegon County, on September 22, 1928, and is now in the John Ball Park Zoo at Grand Rapids. Mr. R. L. McGrady was casting for muskellunge at Mona Lake, when this bird flew in front of and near him, and was caught "on the wing" by the hooks of his bait. While there are a number of state records for this species, it only occurs as a straggler in Michigan.

Falco columbarius. PIGEON HAWK.—On October 17, 1928, an adult female was taken near Muskegon (in that County) by Frank Antisdale, who sent it in the flesh to the Museum of Zoölogy. On September 16, the writer collected a fine male at "Point Lookout," Arenac County. This species seems to follow the Lake Shore beaches, usually during the fall migration of small birds in September and October. In a large series in our collection there is but one Spring record, that of April 30, 1918, in Berrien County, when the writer collected a beautiful adult male on top of a high Sand Dune. Since that date the species has not been seen by the writer until October 17, 1928, but it no doubt occurs rarely in the fall migration. The only record of its occurrence in any numbers was in 1915 on Charity Island, Saginaw Bay, when the writer saw a dozen from September 9 to October 11, and collected nine of them, five females and four males.—NORMAN A. WOOD, *Museum of Zoology, University of Michigan, Ann Arbor, Michigan.*

Lewis' Woodpecker in Rhode Island.—On November 16, Mr. Jeremiah Triggs, superintendent of Roger Williams Park, brought in to me an adult Lewis's Woodpecker (*Asyndesmus lewisi*) which he took away