

Dendroica dominica dominica. YELLOW-THROATED WARBLER.—One was noted at Fairhope by H. Flagg, on September 18, 1932, apparently the first fall record for the state.

Hedymeles ludoviciana. ROSE-BREADED GROSBEAK.—An immature male was found dead by Kenneth Edwards at Fairhope on October 9, 1932. It is an uncommon migrant in Alabama.

Passerina cyanea. INDIGO BUNTING.—An immature male was picked up dead by S. W. Alexander at Fairhope on October 6, 1932, the first autumn record for the southern counties. I personally have noted it as fairly common in the spring.

Carpodacus purpureus purpureus. EASTERN PURPLE FINCH.—One of this species was seen at Fairhope on January 18, 1933, by D. McIntosh, apparently the first record for the coastal counties.

Junco hyemalis hyemalis. SLATE-COLORED JUNCO.—One was noted at Fairhope on November 13, 1932, by D. McIntosh. H. Flagg reports ten seen on January 3, 1933, fifteen on January 8, and again fifty on February 4, 1933. These are the first records for the coast region.—HELEN M. EDWARDS, *School of Organic Education, Fairhope, Alabama*.

Notes from Mountain Lake, Florida.—The following notes have been made on my winter sojourns at Mountain Lake, Fla., about thirty miles east of Lake Kissimmee.

Ardea wurdemanni. WÜRDEMANN'S HERON.—On March 14, 1931, while motoring from Lake Wales to Lake Kissimmee, in the ditch along the roadside about five miles from Lake Kissimmee, I was fortunate enough to see one of these herons. It allowed me to approach so close that there was no question of its identity.

Aramus pictus pictus. LIMP KIN.—There are still many Limpkins in the Kissimmee district and a particularly large colony on the stream between Lake Rosalie and Tiger Lake which is heavily wooded and very wild.

Xanthocephalus xanthocephalus. YELLOW-HEADED BLACKBIRD. On March 5, 1932, I secured, near the public playground, near Lake Wales, Fla., a Yellow-headed Blackbird which I believe is the fifth record of the species for the state.—HENRY O. HAVEMEYER, 25 Broadway, New York.

Notes from Tippecanoe Co., Indiana.—The following notes have been made during the past year by the writers, in Tippecanoe County, Indiana, which is located on the Wabash River in the west-central part of the state.

Casmerodius albus egretta. AMERICAN EGRET.—On the unusually late date of October 12, 1932, one bird, apparently normal, was seen at Headley's Lake, at which one or more birds had been frequently noted since late July. Usually all the Egrets have left Indiana by the middle of September.

Querquedula discors. BLUE-WINGED TEAL.—On July 8, 1932, a single individual was seen on Headley's Lake, and a trip there on July 18 dis-