

present on May 6. Two males collected on May first had almost completed the pre-nuptial molt. Perhaps the strain of molting induced them to linger on a good feeding ground until this late date. Dr. Eaton gave April 18 as the latest state record available.—DEAN AMADON, *Franklinville, New York*.

Notes from Central Pennsylvania.—May 15—Forty-eight Common Terns (*Sterna h. hirundo*) and two Black Terns (*Chlidonias n. surinamensis*) in company with eleven Ring-billed Gulls (*Larus delawarensis*) were observed on the Susquehanna River about one mile below Sunbury (Shindle and May).

May 23—Four Red-backed Sandpipers (*Pelidna a. sakhalina*) were observed on Thompson's Pond in the borough of State College. The observer was within thirty feet of the birds and used Zeiss 8 x 40 binoculars. On the same day approximately thirty Semipalmated Plover, six Semipalmated Sandpipers and about forty Least Sandpipers. The last three species mentioned were at the Red Mill and near Potter's Mills, Centre County (Stokes, Melenger, Meyer and May). The Red-backed Sandpipers remained in the locality for three days.

May 26—A Broad-winged Hawks' (*Buteo platypterus*) nest was found near Shingletown Gap in Centre County. The nest contained three eggs. Nesting material—twigs and leaves in the crotch of a white oak about twenty-five feet from the ground. (June 12th—Hawk still sitting on nest, no close examination made.) About two hundred feet from the Hawks' nest a nest of the Northern Pileated Woodpecker (*Ceophloeus p. abieticola*) was found. One of the adults was seen leaving the nest. The next day when visited the female was on the nest about forty-five minutes later she was relieved by the male. (June 12—after watching the nest for about an hour the female was seen entering the cavity where she remained for about forty minutes when she was relieved by the male.)

May 29—One male White-winged Scoter (*Melanitta deglandi*) in company with two Red-breasted Mergansers (*Mergus serrator*) on the Susquehanna river about two miles below Sunbury (Shindle and May).—R. M. MAY, *Dept. of Forests and Waters, Harrisburg, Pa.*

Notes from the Madison, Wisconsin, Region.—*Florida c. caerulea*. LITTLE BLUE HERON.—On July 28, 1934, two Little Blue Herons were found at Fish Lake, Dane County. Present also were two American Egrets (*Casmerodius a. egretta*). The following day there were five Little Blue Herons and three Egrets. All of the former were in juvenal plumage. One of them, a male, was collected on July 31. There seem to be only two previous references to their occurrence in the state.

Falco c. columbarius. EASTERN PIGEON HAWK.—On April 8, five miles north of Prairie du Sac, a male was taken: weight 172.5 grams.

Limosa haemastica. HUDSONIAN GODWIT.—Three of this species were found in a pond near Verona on May 12. The following morning one was