collection in themselves, representing various founders of the A. O. U. and extending in dates back to the 80's. These are offered to any interested institution or individual and I will send them all to one address or divided them among any applicants there may be, say in a month subsequent to the distribution of this issue of The Auk.—W. L. MCATER, Fish and Wildlife Service, Chicago 54, Illinois.

RESEARCH REQUEST

I am interested in obtaining reprints, copies of unpublished manuscripts and results or observations made in connection with the biological control of mosquitoes. I want to bring this information together in the form of annotated bibliographies.—
J. B. Gerberich, Dept. of Biological Science, Michigan State College, East Lansing, Michigan.

CORRESPONDENCE

Under-wing Fishing of the Black Heron

EDITOR OF THE AUK:—The remarkable under-wing fishing of Melanophoyx ardesiaca so excellently described and illustrated by Capt. Delacour in The Auk (63: 441-442, pl. 11, July, 1946) may have a significance which is not suggested in the note. It is well known that some fish (e. g., the black bass) congregate in shadows and avoid sunlit water. Possibly the Black Heron is utilizing this habit in some fish of the region. Would that we knew whether such a habit is instinctive (inherited), discovered de novo by each individual, or learned imitatively!—Charles H. Blake, Massachusetts Institute of Technology, Cambridge, Massachusetts.