

White-tailed Kite in Nicaragua.—An immature male White-tailed Kite (*Elanus leucurus majusculus*) was taken by Brigadier General Nicholas Biddle of Bethayres, Pennsylvania, at San Isidro, Nicaragua, on 23 February 1961. The specimen is now in the collection of the Academy of Natural Sciences of Philadelphia (A.N.S.P. 168357). The comparatively long tail, which measures 171 mm from the bases of the middle rectrices, serves to identify it as the northern race, which has not hitherto been recorded south of British Honduras and Guatemala. The nominate form is apparently confined to South America.—JAMES BOND, *Academy of Natural Sciences of Philadelphia, Philadelphia 3, Pennsylvania.*

Nest and eggs of the Cabanis Tiger Heron in Chiapas, Mexico.—On a trip to southeastern Chiapas 8–20 July 1962, accompanied by Miguel Alvarez del Toro, I found two isolated nests of *Heterocnus mexicanus*. The nests, 10 miles apart, were in mangroves (*Rhizophora mangle*) some 45 miles northwest of Puerto Madero on the Pacific coast. One nest was 3 feet above water and was photographed on 14 July (see Figure 1). The other nest was 15 feet above the water and was at the periphery of a colony of Agami Herons (*Agamia agami*).



Figure 1. Nest and eggs of Cabanis Tiger Heron, Chiapas, Mexico, 14 July 1962. From a color transparency.