

# INDEX TO VOLUME 96

Compiled by Jean Ferner

This index is primarily one to new information. Previously published information referred to in this volume has not been indexed unless extensive comparisons or reinterpretations are involved. For some species, especially ones not in the A.O.U. Check-list, scientific names only have been entered, and in some instances entries under scientific names are more extensive than those under vernacular names. New generic, specific, and subspecific names are printed in **boldface** type.

- Abnormality of *Melanerpes erythrocephalus*, 206–207; *Oxyura* sp., 479  
*Accipiter cooperii*, 782  
*A. superciliosus*, 345–349  
*Acestrura bombus*, 346, 350  
*Aechmophorus occidentalis*, 573–586  
*Agelaius cyanopus*, 181  
*A. forbesi*, 179–181  
*A. phoeniceus*, 190–192, 208–211, 353–363, 623, 789  
*A. thilius*, 478  
*Agriornis albicauda*, 477  
*A. montana*, 477, 480  
*Aix sponsa*, 74, 296–305  
Alabama, 116–130, 206–207, 533  
Alaska, 56–67, 131–142, 189–190, 234–252, 306–318, 670, 792–794  
Albinism in *Melanerpes erythrocephalus*, 206–207  
*Alectoris chukar*, 499–505  
Ali, Sálim, Field guide to the birds of the eastern Himalayas (rev.), 438–439  
Allan, Thomas A., Parental behavior of a replacement male Dark-eyed Junco, 630–631  
*Allocotopterus deliciosus*, 347, 350  
Altruism (see Communal breeding)  
Amadon, Dean, review by, 832–833  
*Amazilia amabilis*, 346  
*A. franciae*, 346  
*Amazona albifrons*, 184  
*A. farinosa*, 345  
*A. festiva*, 345, 350  
*Amazonetta brasiliensis*, 74  
American Ornithologists' Union, Awards, 212; Proceedings of the Ninety-sixth Stated Meeting, 1AA–19AA  
*Ammodramus henslowii*, 623  
*A. savannarum*, 621–625  
*Amphispiza belli*, 499–505, 799  
*A. bilineata*, 157–158, 161–162, 165  
*Anabacerthia variegaticeps*, 379  
*Anas cyanoptera*, 476  
*A. discors*, 655–656, 659  
*A. flavirostris*, 476, 479–480  
*A. georgica*, 476, 480  
*A. platyrhynchos*, 74, 655–656, 706  
*A. rubripes*, 737–745  
*A. versicolor*, 476–480  
Anatomy and morphology of *Aechmophorus occidentalis*, 580–582; *Agelaius cyanopus*, 181–182; *A. phoeniceus*, 355–356; *Anas rubripes*, 742–744; *Bonasa umbellus*, 308–309, 316; *Canachites canadensis*, 309–310, 316; *Cerorhinca monocerata*, 150; *Certhidea olivacea*, 800–802; *Chen caerulescens*, 70–72, 545–546; *C. rossii*, 545–546; *Chlorochrysa calliparaea*, 808–809; *C. nitidissima*, 809; *C. phoenicotis*, 809; *Conopias inornata*, 768, 773; *Coturnix coturnix*, 12–30; *Curaeus curaeus*, 181–182; *C. forbesi*, 181–182; *Dendragapus obscurus*, 308, 316; *Dendroica caerulescens*, 757, 762; *D. castanea*, 757, 760, 762; *D. cerulea*, 757; *D. coronata*, 757; *D. discolor*, 757; *D. dominica*, 757; *D. fusca*, 757; *D. kirtlandii*, 757; *D. magnolia*, 757, 769; *D. palmarum*, 757; *D. pensylvanica*, 757, 760; *D. petechia*, 757; *D. pinus*, 757; *D. striata*, 757; *D. tigrina*, 757; *D. virens*, 757; *Empidonax minimus*, 265–266, 270; *Florida caerulea*, 121, 123; *Fringilla coelebs*, 91; *Geospiza conirostris*, 800–802; *G. difficilis*, 800–802; *G. fortis*, 800–802; *G. fuliginosa*, 800–802; *G. magnirostris*, 800–802; *G. scandens*, 800–802; *Gnorimopsar chopi*, 181–182; *Hemiriccus cinnamomeipictus*, 443–446; *H. kaempferi*, 443, 446; *H. mirandae*, 443, 446; *Hesperiphona vespertina*, 90–91; hybrids of *Chen caerulescens* and *C. rossii*, 545–547; hybrids of *Tyrannus couchii* and *T. melancholicus*, 227–229; hybrids of *Zonotrichia albicollis* and *Z. atricapilla*, 595–597; hybrids of *Zonotrichia leucophrys* and *Z. querula*, 596–599; *Junco hyemalis*, 78–88, 94–99; *Lagopus lagopus*, 106–115, 313–316; *L. leucurus*, 310–311, 316; *L. mutus*, 311–316; *Lunda cirrhata*, 150; *Pedioecetes phasianellus*, 308, 316; *Pipilo erythrorthalus*, 688–693; *Pitylus fuliginosus*, 612; *P. grossus*, 612; *Ptychoramphus aleuticus*, 150; *Saltator albicollis*, 612; *S. atriceps*, 612; *S. atricollis*, 612; *S. atripennis*, 612; *S. aurantiirostris*, 612; *S. cinctus*, 610–612; *S. coerulescens*, 612; *S. maximus*, 612; *S. nigriceps*, 612; *S. orenocensis*, 612; *S. rufiventris*, 612; *S. similis*, 612; *Setophaga ruticilla*, 265–266, 270, 279; *Sterna albifrons*, 31–39; *S. hirundo*,

- 12–30; *Sturnus vulgaris*, 12–30; *Tangara arthus*, 809; *T. chilensis*, 809; *T. cyanocephala*, 809; *T. fastuosa*, 809; *T. parzudakii*, 809; *T. ruficeps*, 809; *T. xanthocephala*, 809; *Tyrannus couchii*, 222–230; *T. melancholicus*, 222–230; *Zonotrichia albicollis*, 91, 595–597; *Z. atricapilla*, 595–597; *Z. leucophrys*, 596–599; *Z. querula*, 88–90, 596–599
- Ankney, C. Davison, Does the wing molt cause nutritional stress in Lesser Snow Geese?, 68–72; see Scott, D. M., and ———
- Anous stolidus*, 74
- A. tenuirostris*, 74
- Anser fabalis*, 74
- Antarctica, 675–681
- Anthus correndera*, 477
- A. furcatus*, 477
- Antvireo, Plain, 379, 380
- Antwren, Slaty, 379
- Anumbius annumbi*, 607
- Aphelocoma coerulescens*, 158, 166
- Aquila chrysaetos*, 548, 669
- Ara ambiguus*, 345, 348
- A. severa*, 345
- Aramides wolfi*, 345, 348
- Aratinga astec*, 184
- Ardea herodias*, 364, 628–630
- Arenaria interpres*, 189
- Argentina, 606–608
- Arizona, 413, 776–788, 796–798
- Armstrong, Edward A., The life and lore of the bird. In nature, art, myth and literature (rev.), 640–641
- Arnall, L., and I. F. Keymer, Bird diseases (rev.), 432–433
- Arremon aurantiirostris*, 348
- Ashkenazie, Shoshana, and Uriel N. Safriel, Breeding cycle and behavior of the Semipalmated Sandpiper at Barrow, Alaska, 56–67
- Asio flammeus*, 725
- Asthenes dorbignyi*, 477
- A. wyatti*, 477
- Atlapetes brunneinucha*, 378, 380–386
- Attila, Bright-rumped, 378
- Attila spadiceus*, 347, 378
- A. torridus*, 347, 350
- Auk, Great, 790–792
- Auklet, Cassin's, 143–151; Rhinoceros, 145
- Aulacorhynchus haematopygus*, 346, 350
- Auriparus flaviceps*, 157–158, 161–162
- Australia, 489–498
- Automolus ochrolaemus*, 346
- A. rubiginosus*, 346
- Aythya americana*, 706
- Baja California, 152–167
- Baker, R. Robin, The evolutionary ecology of animal migration (rev.), 818–819
- Balph, David F., see Balph, Martha Hatch, et al.
- Balph, Martha Hatch, Flock stability in relation to social dominance and agonistic behavior in wintering Dark-eyed Juncos, 714–722; David F. Balph, and H. Charles Romesburg, Social status signaling in winter flocking birds: an examination of a current hypothesis, 78–93
- Bananaquit, 378
- Baptista, Luis F., see Bitterbaum, Erik, and ———
- Barbet, Prong-billed, 378
- Barbtail, Spotted, 378, 380–387
- Baryphthengus ruficapillus*, 346
- Basileuterus culicivorus*, 378, 380–386
- B. rivularis*, 347
- B. tristriatus*, 378, 380, 383–388
- Becard, Barred, 379
- Beehler, Bruce McP., Upland birds of northeastern New Guinea (rev.), 644
- Behavior, activity patterns, 704–713; age-related, 117–118, 806–807; agonistic, 78–93, 94–99, 186–187, 274–278, 333, 353–363, 384, 388, 414, 415–417, 513, 529–530, 557, 591, 593, 613–614, 677–681, 706–711, 714–722, 733, 789; altruistic (see Communal breeding); antipredator, 375–390, 491–492, 496, 782–783; breeding, 1–5, 6–9, 32, 38, 42–55, 56–67, 104, 117–128, 333, 364–369, 393–400, 421–422, 472–473, 484–488, 489–498, 562–572, 587–594, 604–608, 613–614, 630–631, 677–681, 704–713, 729–730, 732–734, 771–774, 776–788, 797; copulatory, 417–418, 489, 589, 630, 777; courtship, 37–38, 58–59, 134, 356–357, 466, 472–473, 489, 493–497, 587, 591, 593, 706–709, 777–778; distraction, 62; dominance, 78–93, 94–99, 126, 274–278, 353–363, 415–417, 506, 509–514, 590, 613–614, 677–681, 714–722, 734, 777, 779, 781; effect of environment on, 61–63, 386, 421–422, 528–529, 630, 780, 797; escape, 122–127; feeding, 61–64, 118–127, 186–187, 195, 270–274, 277, 284–295, 320–325, 375–390, 395–400, 413–414, 415–417, 418–420, 421–422, 448–461, 487, 548, 557, 590–592, 607, 608–609, 614, 630, 706–711, 729–730, 732–734, 773–774, 777–786, 806–807; feeding aggregations, 375–390; food-begging, 286–288, 291–294, 415–417; foraging (see Behavior, feeding); homosexual, 1–5, 6–9; incubation, 59–66, 104, 118, 491–492, 589–590, 604–606, 614, 677–681, 711, 771–772, 778–779; individual distance, 98, 720; interspecific, 64, 265–283, 333, 375–390, 413–414, 449, 453, 455, 591, 593, 608–609, 677–681, 706, 782–783, 789, 806–807; learning, 471, 506–508, 609, 688–693, 806–807; maintenance, 525–531, 706–711; mate replacement, 630–631; mating preference, 1–5, 6–9; movements in home range, 63, 382,

- 386–390; nesting, 56–67, 119, 134, 139, 417–418, 491–492, 587–589, 592, 604, 607–608, 677–681, 706–709, 778–780; nest-site tenacity, 58, 665–666, 668, 671; ontogeny of, 34–38; pair, 1–5, 6–9, 34–38, 58–66, 417–418, 472–473, 492–497, 528–529, 587–594, 613–614, 630–631, 706–709; parental care, 62–64, 118–127, 284–295, 366–367, 398–400, 415–417, 489–498, 590–592, 604–606, 614, 630–631, 708–711, 729–730, 732–733, 771–774, 781–783, 797; site fidelity, 40–55, 359–361, 565–566, 570; social, 64, 78–93, 94–99, 375–390, 471, 492–494, 509–514, 714–722; status-signaling, 78–93, 94–99; territorial, 40–55, 63–64, 122–124, 134, 139–140, 186–187, 275, 278, 382, 386–390, 418, 421–422, 448–461, 489–498, 512–513, 557, 631, 706–711, 733–734; thermoregulatory, 604–606, 797; vocal, 31–39, 62–63, 287–295, 356–357, 385, 388, 413–414, 472–473, 489, 491–492, 496, 506–515, 587–588, 590–591, 688–693, 710, 733, 771–772, 777
- Behavior of *Agelaius phoeniceus*, 353–363, 789; *Anabacerthia variegaticeps*, 379; *Anas flavirostris*, 479; *Anumbius annumbi*, 607; *Atlapetes brunneinucha*, 378, 380–386; *Attila spadiceus*, 378; *Basileuterus culicivorus*, 378, 380–386; *B. tristriatus*, 378, 380, 383–388; *Calcarius lapponicus*, 134, 139–140; *Calidris alba*, 557; *C. pusilla*, 56–67; *Campylorhamphus pusillus*, 379, 380; *Carpodacus mexicanus*, 466, 471–473; *Catharus frantzii*, 379; *C. fuscater*, 378; *C. mexicanus*, 378; *Charadrius vociferus*, 604–606; *Chlorophonia callophrys*, 378; *Chlorospingus ophthalmicus*, 378, 380–386; *Cinclodes fuscus*, 480; *Cistothorus palustris*, 507–515; *Coereba flaveola*, 378; *Conopias inornata*, 771–774; *Coturnix coturnix*, 10–30; *Cranioleuca erythrops*, 378, 380–386; *Cyanolyca cucullata*, 378; *Dendroica virens*, 379, 383; *Dryocopus pileatus*, 587–594; *Dysithamnus mentalis*, 379–380; *Empidonax flavescens*, 378; *E. minimus*, 265–283; *Eudyptes chrysocome*, 104; *Florida caerulea*, 117–128; *Fringilla coelebs*, 91; *Fulica americana*, 479, 704–713; *Furnarius rufus*, 607; *Glyphorynchus spirurus*, 379; *Gymnorhinus cyanocephalus*, 796–797; *Henicorhina leucophrys*, 378, 380–386; *Hesperiphona vespertina*, 90–91; *Hylophilus decurtatus*, 379; *Junco hyemalis*, 78–88, 94–99, 630–631, 714–722; *Larus californicus*, 6–9; *L. delawarensis*, 1–5, 6–9, 284–295, 806–808; *Lysurus crassirostris*, 379; *Menura superba*, 489–498; *Metriopelia ceciliae*, 479; *Mimus polyglottos*, 421–422; *M. saturninus*, 607–608; *Mionectes olivaceus*, 378; *Mniotilla varia*, 378, 383; *Molothrus ater*, 484–488; *Myadestes melanops*, 378; *Myioobius miniatus*, 378, 380, 382–387; *Myrmotherula schisticolor*, 379; *Nectarinia famosa*, 449; *N. kilimensis*, 449; *N. reichenowi*, 449, 453–460; *N. venusta*, 449; *Nesomimus parvulus*, 730–735; *Nycticorax nycticorax*, 364–369; *Oporornis formosus*, 379, 380, 383; *Otus asio*, 320–325; *Oxyura* sp., 479; *Pachyramphus versicolor*, 379; *Pandion haliaetus*, 413–414, 415–417; *Parabuteo unicinctus*, 776–788; *Parus atricapillus*, 40–55; *Passerulus sandwichensis*, 393–400; *Passerina ciris*, 417–418; *Phaeoptilus tapera*, 607; *Phoenicoparrus andinus*, 333; *P. jamesi*, 333; *Phoenicopterus chilensis*, 333; *Picoides nuttallii*, 418–420; *Piculus simplex*, 378; *Piranga rubra*, 379; *Platyrinchus mystaceus*, 378, 380; *Polioptila dumicola*, 608; *Premnoplex brunnescens*, 378, 380–387; *Pygoscelis adeliae*, 677–681; *P. antarctica*, 677–681; *Quiscalus mexicanus*, 789; *Q. quiscula*, 608–609; *Rhynchococcyx brevirostris*, 378; *Semnornis frantzii*, 378; *Setopaga ruticilla*, 265–283; *Sialia currucoides*, 613–615; *S. sialis*, 562–572; *Sittasomus griseicapillus*, 378, 380–387; *Sphyrapicus varius*, 195; *Sterna hirundo*, 10–30; *Strix occidentalis*, 526–531; *Sturnus vulgaris*, 10–30; *Syndactyla subalaris*, 378, 380–387; *Tangara dowii*, 379; *T. icterocephala*, 378; *Thripadectes rufobrunneus*, 379; *Troglodytes ochraceus*, 378; *Verminivora chrysoptera*, 379, 380, 383; *V. peregrina*, 186–187; *Wilsonia pusilla*, 379, 383; *Xiphorhynchus erythropygius*, 378, 380–387; *Zonotrichia albicollis*, 91; *Z. querula*, 88–90
- Beissinger, Steven R., see Osborne, David R., and
- 
- Belize, 227–229
- Billings, Raymond R., see Vermeer, Kees, et al.
- Bitterbaum, Erik, and Luis F. Baptista, Geographical variation in songs of California House Finches (*Carpodacus mexicanus*), 462–474
- Blackbird, Brewer's, 499–505; Forbes', 179–183; Red-winged, 190–192, 208–211, 353–363, 623, 789; Rusty, 392; Yellow-headed, 192–195
- Bluebird, Eastern, 562–572; Mountain, 613–615
- Boag, P. T., see Grant, P. R., et al.
- Bock, Walter J., editor of Reviews, 213–217, 426–440, 632–648, 818–834; reviews by, 637–638, 642–643, 646–648, 821–823
- Bolborhynchus aurifrons, 477
- B. orbygnesius, 477, 480
- Bolivia, 328–342
- Bombycilla cedrorum, 196–198
- Bonasa umbellus, 308–309, 316
- Botswana, 407–408
- Bovino, R. Russell, and Edward H. Burtt, Jr., Weather-dependent foraging of Great Blue Herons (*Ardea herodias*), 628–630

- Branta canadensis*, 234–252  
 Brauner, Joseph, In memoriam: Aretas Andrews Saunders, 173–178  
 Brazil, 179–183, 443, 446  
 Breeding of *Aechmophorus occidentalis*, 573–586; *Agelaius phoeniceus*, 353–363; *Amphispiza belli*, 798–799; *Aquila chrysaetos*, 669; *Bombycilla cedrorum*, 196–197; *Branta canadensis*, 234–252; *Buteo lagopus*, 669, 672–673; *Calcarius lapponicus*, 133, 138; *Calidris pusilla*, 56–67; *Cerorhinca monocerata*, 147–150; *Charadrius vociferus*, 604–606; *Circus cyaneus*, 370–374; *Conopias inornata*, 769–774; *Dendroica castanea*, 600–603; *Dryocopus pileatus*, 587–594; *Empidonax minimus*, 274, 277–279; *Eudyptes chrysocome*, 100–105; *Falco peregrinus*, 662–674; *F. rusticolus*, 669–672; *Florida caerulea*, 116–130; *Fulica americana*, 479, 704–713; *F. gigantea*, 479; *Gallinula chloropus*, 479; *Gymnorhinus cyanocephalus*, 796–797; *Hemimicrurus cinnamomeipectus*, 444–445; *Juncos hyemalis*, 630–631; *Lanius ludovicianus*, 798–799; *Larus argentatus*, 694–703; *L. californicus*, 6–9; *L. delawarensis*, 1–5, 6–9, 284–295; *Lunda cirrhata*, 150, 792–794; *Melanerpes erythrocephalus*, 206–207; *Menura superba*, 489–498; *Metrostomelina melanoptera*, 480; *Mimus longicaudatus*, 723–727, 733; *M. polyglottos*, 421–422; *Molothrus ater*, 483–488; *Nesomimus parvulus*, 723–736; *Nycticorax nycticorax*, 364–369; *Oreoscoptes montanus*, 798–799; *Pandion haliaetus*, 415–417; *Parabuteo unicinctus*, 413, 776–788; *Parus atricapillus*, 42–55; *Passerulus sandwichensis*, 391–401; *Passerina ciris*, 417–418; *Phalacrocorax olivaceus*, 478–479; *Phoenicoparrus jamesi*, 332, 334; *Phoenicopterus ruber*, 615–619; *Phrygilus plebejus*, 480; *Plegadis ridgwayi*, 479, 481; *Podiceps rolland*, 478, 481; *Ptychoramphus aleuticus*, 143–151; *Pygoscelis adeliae*, 675–681; *P. antarctica*, 675–681; *Quelea quelea*, 520–521; *Rissa tridactyla*, 537–543; *Setophaga ruticilla*, 274, 277–279; *Sialia sialis*, 562–572; *Sicalis uropygialis*, 480; *Spizella breweri*, 798–799; *Steatornis caripensis*, 188; *Tyto alba*, 204  
 Breeding season of *Aechmophorus occidentalis*, 577; *Bombycilla cedrorum*, 196–197; *Calidris pusilla*, 57–58; *Conopias inornata*, 769; *Hemimicrurus cinnamomeipectus*, 444–445; *Larus argentatus*, 696; *Menura superba*, 489; *Mimus longicaudatus*, 724; *Nesomimus parvulus*, 724; *Parabuteo unicinctus*, 413; *Passerulus sandwichensis*, 394; *Pygoscelis adeliae*, 676–677, 679–680; *P. antarctica*, 676–677, 679–680; *Quelea quelea*, 520; *Saltator cinctus*, 610; *Sialia sialis*, 563; *Steatornis caripensis*, 188  
 Breeding success, effects of interference on, 1–5, 190–191, 364–369, 618, 665–667, 672, 797–798, 815–817; estimation of, 651–661  
 Breeding success of *Aechmophorus occidentalis*, 578; *Agelaius phoeniceus*, 190–191, 353–363, 789; *Amphispiza belli*, 798–799; *Anas discors*, 655–656, 659; *A. platyrhynchos*, 655–656; *Bombycilla cedrorum*, 197; *Buteo lagopus*, 669, 672–673; *Calcarius lapponicus*, 135, 138; *Circus cyaneus*, 370–374; *Conopias inornata*, 774; *Coturnix coturnix*, 28–29; *Dendroica kirtlandii*, 651–653, 655–658; *Falco peregrinus*, 662–674; *F. rusticolus*, 669, 672; *Florida caerulea*, 116–130; *Fulica americana*, 710–711; *Gelochelidon nilotica*, 203; *Gymnorhinus cyanocephalus*, 797–798; *Lanius ludovicianus*, 798; *Larus argentatus*, 694–703; *L. californicus*, 6–9; *L. delawarensis*, 1–5, 6–9; *Lunda cirrhata*, 793–794; *Melanerpes erythrocephalus*, 206–207; *Mimus longicaudatus*, 725–727; *M. polyglottos*, 421–422; *Nesomimus parvulus*, 725–727, 731–732; *Nycticorax nycticorax*, 364–369; *Oreoscoptes montanus*, 798–799; *Pandion haliaetus*, 415–417; *Passerulus sandwichensis*, 391–401; *Passerina ciris*, 417–418; *Phoenicopterus ruber*, 618; *Pooecetes gramineus*, 802–805; *Pygoscelis adeliae*, 677–681; *P. antarctica*, 677–681; *Rissa tridactyla*, 537–543; *Sialia sialis*, 562–572; *Spizella breweri*, 798–799; *Sterna hirundo*, 28–29; *Sturnus vulgaris*, 28–29  
 Brisbin, I. Lehr, Jr., see Clay, Deborah L., et al.  
 British Columbia, 143–151, 196, 306–318  
 Brodkorb, Pierce, Brewster Award, 212  
 Brush-Finch, Chestnut-capped, 378, 380–386  
*Bubo virginianus*, 202, 477  
 Bunting, Lazuli, 499–505; Painted 417–418  
 Burger, A. E., and A. J. Williams, Egg temperatures of the Rockhopper Penguin and some other penguins, 100–105  
 Burger, Joanna, Colony size: a test for breeding synchrony in Herring Gull (*Larus argentatus*) colonies, 694–703; and Michael Gochfeld, Age differences in Ring-billed Gull kleptoparasitism on Starlings, 806–808  
 Burtt, Edward H., Jr. (Ed.), The behavioral significance of color (rev.), 640; see Bovino, R. Russell, and ——  
 Bush-Tanager, Common, 378, 380–386  
*Buteo jamaicensis*, 74, 782  
*B. lagopus*, 669, 672–673  
*B. poecilochrous*, 476, 479  
*Butthraupis eximia*, 348, 350  
 Butler, Ronald G., and Peter Lukasiewicz, A field study of the effect of crude oil on Herring Gull (*Larus argentatus*) chick growth, 809–812  
 Butler, T. Y., The birds of Ecuador and the Galapagos Archipelago (rev.), 645–646

- Cacicus uropygialis*, 347  
*Calcarius lapponicus*, 131–142, 392  
 Calef, George W., and Douglas C. Heard, Reproductive success of Peregrine Falcons and other raptors at Wager Bay and Melville Peninsula, Northwest Territories, 662–674  
*Calidris alba*, 190, 551–561  
*C. bairdii*, 476  
*C. melanotos*, 476  
*C. pusilla*, 56–67  
 California, 196, 234–252, 410, 418–420, 462–474, 544–550, 551–561, 574–575, 577, 580  
 Calls (see Vocalizations)  
*Calypte costae*, 157–158, 161–162, 165  
*Campylorhamphus pusillus*, 379, 380  
*C. trochilirostris*, 346  
*Campylorhynchus brunneicapillus*, 159, 162, 166  
*Canachites canadensis*, 309–310, 316  
 Cantor, Frederic L., see Leck, Charles F., and
- 
- Caprimulgus indicus*, 189  
 Cardinal, 159  
*Cardinalis cardinalis*, 159  
*C. sinuatus*, 159  
*Carduelis flammea*, 392  
*C. hornemannii*, 392  
*Carpodacus mexicanus*, 159, 162, 462–474  
*Carpodectes hopkei*, 347, 349  
*Casmerodius albus*, 476  
*Catamenia inornata*, 478  
*Cathartes aura*, 476  
*Catharus frantzii*, 379  
*C. fuscater*, 378  
*C. mexicanus*, 378  
*C. ustulatus*, 347, 351  
*Catherpes mexicanus*, 159  
*Celeus loricatus*, 346  
*Centrocercus urophasianus*, 499–505  
*Centurus aurifrons*, 184  
*C. pygmaeus*, 184  
*Cephalopterus penduliger*, 347, 349  
*Cercomacra tyrannia*, 347  
*Cerorhinca monocerata*, 145–150  
*Certhidea olivacea*, 726, 729, 800–802  
 Chaffinch, 91, 253–264  
 Chandler, Randall S., see Rutledge, James T., and
- 
- Charadrius mongolus*, 189  
*C. vociferus*, 604–606  
 Chat, Yellow-breasted, 499–505  
 Check List Committee of the South Dakota Ornithologists' Union, Byron E. Harrell (Ed.), The birds of South Dakota (rev.), 639–640  
*Chen caerulescens*, 68–72, 544–550  
*C. rossii*, 544–550  
 Chickadee, Black-capped, 40–55  
 Chile, 328–342, 619–621  
*Chloephaga melanoptera*, 476, 480  
*Chlorochrysa calliparaea*, 808–809  
*C. nitidissima*, 809  
*C. phoenicotis*, 809  
*Chlorophanes spiza*, 348  
*Chlorophaenia*, Golden-browed, 378  
*Chlorophaenia callophrys*, 378  
*Chlorospingus ophthalmicus*, 378, 380–386  
*Chondestes grammacus*, 160  
*Chrysolophus amherstiae*, 74  
*C. pictus*, 74  
 Chukar, 499–505  
*Cinclodes*, Bar-winged, 477, 480  
*Cinclodes atacamensis*, 477  
*C. fuscus*, 477, 480  
*Circus cyaneus*, 370–374, 476  
*Cissolophya yucatanica*, 184  
*Cistothorus palustris*, 507–515  
 Clark, L., and Diana J. Gabaldon, Nest desertion by the Piñon Jay, 796–798  
 Clark, Richard J., Dwight G. Smith, and Leon H. Kelso, Working bibliography of owls of the world (rev.), 643–644  
 Clarke, C. H. D., In memoriam: Hoyes Lloyd, 402–406  
 Clay, Deborah L., I. Lehr Brisbin, Jr., and Kenneth A. Youngstrom, Age-specific changes in the major body components and caloric values of growing Wood Ducks, 296–305  
 Cleptoparasitism (see Kleptoparasitism)  
 Clutch size of *Aechmophorus occidentalis*, 577–578; *Bombycilla cedrorum*, 197; *Branta canadensis*, 236–237, 247–250; *Conopias inornata*, 770; *Dendroica castanea*, 601–602; *Fulica americana*, 710–711; *Larus argentatus*, 697–698; *L. californicus*, 6–9; *L. delawarensis*, 1–5, 6–9; *Menura superba*, 489; *Mimus longicaudatus*, 725–726, 733; *Nesomimus parvulus*, 725–726, 733; *Nycticorax nycticorax*, 366, 368; *Parabuteo unicinctus*, 776; *Passerculus sandwichensis*, 394, 396; *Phrygilus plebejus*, 480; *Sialia sialis*, 564  
*Cnemarchus (=Xolmis?) rufipennis*, 477  
 Coates, Brian J., Birds in Papua New Guinea (rev.), 644  
*Coereba flaveola*, 378  
 Cohen, Joel E., Food webs and niche space (rev.), 430–432  
*Colaptes auratus*, 158, 166  
*C. rupicola*, 477  
*Colibri coruscans*, 477  
*Colonia colonus*, 347, 350  
 Colorado, 192–195  
*Columba cayennensis*, 345  
*C. goodsoni*, 345  
*C. subvinacea*, 345  
 Commentaries, The amateur in ornithology, see Mayfield, Harold F.; Author's response to Commentary by David C. Duffy, see Ellison,

- Laurence N.; Human disturbance and breeding birds, see Duffy, David C.; Priorities in bird migration studies, see Gauthreaux, Sidney A., Jr.; R. W. Schreiber's review of *The Sulidae: gannets and boobies*, see J. B. Nelson
- Communal breeding of *Anumbius annumbi*, 607; *Conopias inornata*, 768, 771–774; *Furnarius rufus*, 607; *Larus californicus*, 8–9; *L. delawarensis*, 7–9; *Mimus saturninus*, 607–608; *Nesomimus parvulus*, 730–735; *Parabuteo unicinctus*, 776–788; *Phaeoptile tapera*, 607; *Polioptila dumicola*, 608
- Conner, Richard N., and James C. Kroll, Food-storing by Yellow-bellied Sapsuckers, 195
- Connors, P. G., see Myers, J. P., et al.
- Conopias inornata*, 767–775
- Conover, Michael R., see Miller, Don E., and —; Don E. Miller, and George L. Hunt, Jr., Female-female pairs and other unusual reproductive associations in Ring-billed and California gulls, 6–9
- Cooperative breeding (see Communal breeding)
- Coot, American, 476, 479–480, 704–713; Giant, 476, 479
- Cormorant, Double-crested, 204–206, 364; Neotropic, 476, 478; Olivaceous, 204–206
- Corvus brachyrhynchos*, 367, 804
- C. corax*, 159
- C. cryptoleucus*, 782
- Costa Rica, 186–187, 198–199, 375–390
- Cotinga amabilis*, 184
- Coturnicops noveboracensis*, 422–423
- Coturnix coturnix*, 10–30, 300–303
- Cowbird, Brown-headed, 209–210, 483–488, 499–505, 600–602; Shiny, 774
- Crake, Paint-billed, 425
- Cranioleuca erythrops*, 378, 380–386
- Crow, Common, 367, 804
- Crypturellus cinereus*, 345
- C. soui*, 345
- Curaeus curaeus*, 181–182
- C. forbesi*, 179–183
- Cyanerpes caeruleus*, 348
- C. cyaneus*, 348, 350
- Cyanocitta cristata*, 423–425
- Cyanocorax dickeyi*, 200–201
- C. morio*, 200–201
- C. sanblasiana*, 200–201
- C. yncas*, 184
- Cyanolyca cucullata*, 378
- Cymbilaimus lineatus*, 347
- Cyphorhinus arada*, 347
- Dacnis venusta*, 348, 350
- Damophila julie*, 346
- Daptrius americanus*, 345, 348
- Day, Robert H., Eric P. Knudtson, Dennis W. Wollington, and Robert P. Schulmeister, *Caprimulgus indicus*, *Eurynorhynchus pygmeus*, *Otus scops*, and *Limicola falcinellus* in the Aleutian Islands, Alaska, 189–190
- Dendragapus obscurus*, 308, 316
- Dendrocincla fuliginosa*, 346
- Dendroica caerulescens*, 757–758, 760, 762
- D. castanea*, 600–603, 757, 764
- D. cerulea*, 757
- D. coronata*, 499–505, 757–764
- D. discolor*, 756–757, 759–760
- D. dominica*, 757–763
- D. fusca*, 757–760
- D. kirtlandii*, 651–653, 655–658, 756–760, 763
- D. magnolia*, 757–760, 763
- D. palmarum*, 757–764
- D. pensylvanica*, 757–760, 763
- D. petechia*, 159, 392, 757, 759–761
- D. pinus*, 757, 759, 761–763
- D. striata*, 392, 757–761
- D. tigrina*, 757–764
- D. virens*, 379, 383, 757–760, 763
- Dinsmore, James J., see Ryan, Mark R., and —
- Distribution, spatial, 40–55, 134, 146–150, 267–269, 333–341, 393, 413, 494–496, 516–524, 554–555, 578–580, 583, 665, 669, 770
- Distribution of *Accipiter superciliosus*, 345, 349; *Acestrura bombus*, 346, 350; *Agelaius thilius*, 478; *Agriornis albicauda*, 477; *A. montana*, 477; *Allocotopterus deliciosus*, 347, 350; *Amazilia amabilis*, 346; *A. franciae*, 346; *Amazona farinosa*, 345; *A. festiva*, 345, 350; *Amphisiza bilineata*, 157–158, 165; *Anas cyanoptera*, 476; *A. flavirostris*, 476; *A. georgica*, 476; *A. versicolor*, 476; *Anthus correndera*, 477; *A. furcatus*, 477; *Aphelocoma coerulescens*, 158; *Ara ambiguus*, 345, 348; *A. severa*, 345; *Aramides wolfi*, 345, 348; *Arremon aurantiirostris*, 348; *Asthenes dorbignyi*, 477; *A. wyatti*, 477; *Attila spadiceus*, 347; *A. torridus*, 347, 350; *Aulacorhynchus haematocephalus*, 346, 350; *Auriparus flaviceps*, 157–158; *Automolus ochrolaemus*, 346; *A. rubiginosus*, 346; *Baryphthengus ruficapillus*, 346; *Basileuterus rufularis*, 347; *Bolborhynchus aurifrons*, 477; *B. orbygnysius*, 477; *Bombycilla cedrorum*, 196; *Bubo virginianus*, 477; *Buteo poecilochrous*, 476; *Buthraupis eximia*, 348, 350; *Cacicus uropygialis*, 347; *Calidris bairdii*, 476; *C. melanotos*, 476; *Calypte costae*, 157–158, 165; *Campylorhamphus trochilirostris*, 346; *Campylorhynchus brunneicapillus*, 159; *Caprimulgus indicus*, 189; *Cardinalis cardinalis*, 159; *C. sinuatus*, 159; *Carpodacus mexicanus*, 159; *Carpodectes hopkei*, 347, 349; *Casmerodium albus*, 476; *Catamenia inornata*, 478; *Cathartes aura*, 476; *Catharus ustulatus*, 347, 351; *Catherpes mexicanus*, 159; *Celeus lori*

- catus*, 346; *Cephalopterus penduliger*, 347, 349; *Cercomacra tyrannia*, 347; *Chen caerulescens*, 547–548; *C. rossii*, 547–548; *Chloephaga melanoptera*, 476; *Chlorophanes spiza*, 348; *Chondestes grammacus*, 160; *Cinclodes atacamensis*, 477; *C. fuscus*, 477; *Circus cyaneus*, 476; *Cnemarchus (=Xolmis?) rufipennis*, 477; *Colaptes auratus*, 158; *C. rufigula*, 477; *Colibri coruscans*, 477; *Colonia colonus*, 347, 350; *Columba cayennensis*, 345; *C. goodsoni*, 345; *C. subvinacea*, 345; *Conopias ornata*, 768; *Corvus corax*, 159; *Crypturellus cinereus*, 345; *C. soui*, 345; *Curaeus forbesi*, 180; *Cyanerpes caeruleus*, 348; *C. cyaneus*, 348, 350; *Cymbilaimus lineatus*, 347; *Cyphorhinus arada*, 347; *Dacnis venusta*, 348, 350; *Damophila julie*, 346; *Daptrius americanus*, 345, 348; *Dendrocina fuliginosa*, 346; *Dendroica petechia*, 159; *Diuca speculifera*, 478; *Dryoscopus lineatus*, 346; *Dysithamnus mentalis*, 347; *Electron platyrhynchum*, 346; *Empidonax* spp., 160; *Euphonia minuta*, 348, 350; *E. saturata*, 348, 350; *Eurynorhynchus pygmeus*, 180; *Eutoxeres aquila*, 346; *Falco femoralis*, 476; *F. peregrinus*, 670–672; *F. sparverius*, 476; *Formicarius nigricapillus*, 347; *Fulica americana*, 476; *F. ardesiaca*, 476; *F. gigantea*, 476; *Gallinago (Capella) gallinago*, 476; *Gallinula chloropus*, 476; *Geococcyx californianus*, 158; *Geositta cunicularia*, 477; *G. tenuirostris*, 477; *Geranospiza caerulescens*, 345, 349; *Glaucis aenea*, 346; *Glyptohynchus spirurus*, 346; *Grallaria perspicillata*, 347; *Gymnopithys leucaspis*, 347; *Harpia harpyja*, 345, 348–349; *Heliodoxa jacula*, 346; *Heliothryx barroti*, 346; *Hemithraupis guira*, 348; *Hemitriccus cinnamomeipectus*, 443–444; *H. kaempferi*, 443, 446; *H. mirandae*, 443, 446; *Heterosphingus xanthopygius*, 348; hybrids of *Chen caerulescens* and *C. rossii*, 547–548; hybrids of *Tyrannus couchii* and *T. melancholicus*, 223, 226; *Hyloctistes subulatus*, 346; *Hylophilus minor*, 347; *Hylophylax naeviooides*, 347; *Icterus cucullatus*, 159; *I. mesomelas*, 347; *Iridophanes pulcherrima*, 348, 350; *Lanius ludovicianus*, 159; *Larus serranus*, 477; *Leptasthenura andicola*, 477; *Leptotodon cayanensis*, 345; *Leptopogon superciliosus*, 347; *Leptotila pallida*, 345; *Lessonia rufa*, 477; *Leucopternis semiplumbea*, 345; *Limicola falcinellus*, 189–190; *Lipaugus unirufus*, 347, 350; *Lophortyx californicus*, 158, 499–505; *Lophotriccus pileatus*, 347; *Malaconotila panamensis*, 346, 350; *Manacus manacus*, 347; *Melanerpes pucherani*, 346; *M. uropygialis*, 158; *Metriopelia aymara*, 477; *M. ceciliae*, 477; *M. melanoptera*, 477; *Micrastur ruficollis*, 345, 349; *Microbates cinereiventris*, 347; *Microcerclus marginatus*, 347; *Microrhopias quixensis*, 347; *Mimus polyglottos*, 159; *Mionectes olivaceus*, 347, 349; *Mitrospingus cassini*, 348; *Momotus momota*, 346, 350; *Muscisaxicola alpina*, 477; *M. rufivertex*, 477; *Myiarchus cinerascens*, 157–158; *Myiobius atricaudus*, 347; *M. barbatus*, 347; *Myiodynastes maculatus*, 347; *Myiophobus fasciatus*, 347; *Myiotriccus ornatus*, 347; *Myrmeciza exsul*, 347; *M. immaculata*, 347; *Myrmotherula axillaris*, 347; *M. fulvitarsis*, 347; *M. schisticolor*, 347; *M. surinamensis*, 347; *Neocrex erythrops*, 425; *Notharchus macrorhynchus*, 346, 350; *N. tectus*, 346, 350; *Nothoprocta ornata*, 476; *N. pentlandii*, 476; *Nothura maculosa*, 476; *Notiochelidon murina*, 477; *Nyctibius griseus*, 346, 349; *Nycticorax nycticorax*, 476; *Nystalus radiatus*, 346, 348; *Ochthoeca oenanthesoides*, 477; *Oreocreatus underwoodii*, 346, 350; *Odontophorus erythrops*, 345; *Oreopholus ruficollis*, 476; *Oreotrochilus estella*, 477; *Ornithion semiflavum*, 347; *Ortalis erythroptera*, 345; *Otus guatemalae*, 345; *O. scops*, 189; *Oxyura jamaicensis*, 476; *O. sp.*, 476; *Pachyramphus albogriseus*, 347; *P. cinnamomeus*, 347; *Parula pitiajumi*, 347; *Patagona gigas*, 477; *Penelope purpurascens*, 345, 348–349; *Petrochelidon andecola*, 477; *Phaethornis longuemareus*, 346; *P. superciliosus*, 346; *P. yaruqui*, 346; *Phainopepla nitens*, 160; *Phalacrocorax auritus*, 204–206; *P. olivaceus*, 204–206, 476; *Phalcobaenus megalopterus*, 476; *Pheucticus melanocephalus*, 160; *Philydor erythrocercus*, 346; *Phleocryptes melanops*, 477; *Phloeocetes guayaquilensis*, 346; *P. melanocheirus*, 346, 348; *Phoenicoparrus andinus*, 328–342; *P. jamesi*, 328–342; *Phoenicopterus chilensis*, 328–342, 476; *P. ruber*, 615–619; *Phrygilus alaudinus*, 478; *P. fruticeti*, 478; *P. gayi*, 478; *P. plebejus*, 478; *Piaya cayana*, 345; *Picoides scalaris*, 157–158; *Piculus leucolaemus*, 346; *P. rubiginosus*, 346; *Picumnus olivaceus*, 346; *Pionopsitta haematotis*, 345; *Pionus chalcopterus*, 345; *P. menstruus*, 345; *P. sordidus*, 345, 350; *Pipilo chlorurus*, 160; *P. fuscus*, 159; *Pipra coronata*, 347, 350; *P. mentalis*, 347; *Pipromorpha oleaginea*, 347; *Pittasoma rufopileatum*, 346, 350; *Pitylus grossus*, 348; *Platycichla leucops*, 347; *Platyrinchus coronatus*, 347, 349; *P. mystaceus*, 347; *Plegadis ridgwayi*, 476; *Podiceps occipitalis*, 476; *P. rolland*, 476; *Poecilotriccus ruficeps*, 347, 350; *Pogonotriccus ophthalmicus*, 347, 350; *Polioptila caerulea*, 157–158; *P. melanura*, 157–158; *P. schistaceigula*, 347; *Popelairia versii*, 346; *Premnoplex brunnescens*, 346, 349; *Pteroglossus sanguineus*, 346; *Ptychoram-*

- phus aleuticus*, 147–148; *Pulsatrix perspicillata*, 345, 348; *Pyriglena leuconota*, 347; *Querula purpurata*, 347; *Rallus sanguinolentus*, 476; *Ramphastos ambiguus*, 346, 348; *R. brevis*, 346, 348; *Ramphocaenus melanurus*, 347; *Rhynchocyclus brevirostris*, 347; *R. fulvipectus*, 347, 350; *Rissa tridactyla*, 537–543; *Saltator cinctus*, 610, 612; *Schiffornis turdinus*, 347, 349; *Sclerurus guatemalensis*, 346; *S. mexicanus*, 346, 349; *Sicalis uropygialis*, 478; *Sirystes sibilator*, 347, 350; *Speotyto (Athene) cunicularia*, 477; *Spinus atratus*, 478; *S. crassirostris*, 478; *Spizella breweri*, 160; *Steatornis caripensis*, 187–188; *Tachuris rubrigastra*, 477; *Tachyphonus delatrii*, 348; *T. luctuosus*, 348; *Tangara gyrola*, 348; *T. icterocephala*, 348, 350; *T. lavinia*, 348, 350; *T. nigroviridis*, 348, 350; *Taraba major*, 347; *Terenotriccus erythrurus*, 347; *Thalurania furcata*, 346; *Thamnophilus punctatus*, 347, 351; *Theristicus caudatus*, 476; *Thinocorus orbignyanus*, 476; *Threnetes ruckeri*, 346; *Tinamus major*, 345, 348–349; *Todirostrum nigriceps*, 347; *Tolmomyias assimilis*, 347; *Touit dilectissima*, 345, 350; *Toxostoma cinereum*, 159; *Tringa flavipes*, 476; *T. melanoleuca*, 476; *Troglodytes musculus* (=aedon?), 477; *Trogon collaris*, 346; *T. melanurus*, 346; *T. viridis*, 346; *Turdus chiguanco*, 477; *T. nudigenis*, 347; *T. obsOLEtus*, 347; *Tyranniscus nigrocapillus*, 347, 350; *T. viridiflavus*, 347, 350; *Tryannulus elatus*, 347; *Tyrannus couchii*, 221–233; *T. melancholicus*, 221–233; *Tyto alba*, 480; *Upucerthia serrana*, 477; *U. validirostris*, 477; *Uropsalis segmentata*, 477; *Vanellus resplendens*, 476; *Veniliornis kirkii*, 346; *Vermivora celata*, 160; *Vireo olivaceus*, 347; *Vultur gryphus*, 476; *Xenops minutus*, 346; *X. rutilans*, 346; *Xiphorhynchus erythropygius*, 346; *Zenaida asiatica*, 158; *Z. auriculata*, 477; *Z. macroura*, 158; *Zonotrichia capensis*, 478; *Z. leucophrys*, 160
- Diuca speculifera*, 478
- Dives dives*, 184
- Dolnik, Victor R., and Valery M. Gavrilov, Bioenergetics of molt in the Chaffinch (*Fringilla coelebs*), 253–264
- Dominican Republic, 615–619
- Dove, Mourning, 158, 499–505; White-winged, 158
- Dromaius novaehollandiae*, 74
- Dryocopus lineatus*, 346
- D. pileatus*, 587–594
- Duck, Black, 737–745; Ruddy, 476, 479; Wood, 296–305
- Duffy, David C., Commentary: Human disturbance and breeding birds, 815–816
- Dysithamnus mentalis*, 347, 379–380
- Eagle, Bald, 413–414, 548; Golden, 548, 669; Harpy, 345, 348–349
- Ecology, breeding communities, 37–38, 143–151, 202–203, 675–681; commensal feeding, 375–390; community structure, 280–281, 499–505, 623, 675–681; competition, 265–283, 413–414, 448–461, 521–523, 532–536, 557–560, 584, 675–681, 710–711, 719–721, 789, 806–807; demography, 152–167, 518–523, 532–536, 537–543, 562–572, 573–586, 651–661; dispersal, 40–55, 163–165, 521–523; frugivory, 183–186; habitat partitioning, 147–150, 161–162, 265–283, 334–341, 394–396, 418–420, 516–524, 532–536, 537–543, 551–561, 680–681, 719–721, 756–766, 789; habitat selection, 131–142, 147–150, 152–167, 244–245, 265–283, 394–396, 421–422, 601, 621–625, 680–681, 756–766, 789, 803–804; population structure, 537–543, 544–550, 562–572, 573–586; predator-prey relationships, 202–203, 319–327, 370–374, 408–412, 548, 551–561, 600–603, 620, 680, 798–799; seasonal variation, 499–505, 518–523, 532–536; territoriality, 131–142, 710–711
- Ecology of *Accipiter superciliosus*, 345, 349; *Aescitrura bombus*, 346, 350; *Aechmophorus occidentalis*, 573–586; *Agelaius phoeniceus*, 623, 789; *Alectoris chukar*, 499–505; *Allocotopterus deliciosus*, 347, 350; *Amazilia amabilis*, 346; *A. franciae*, 346; *Amazona farinosa*, 345; *A. festiva*, 345, 350; *Ammodramus henslowii*, 623; *A. savannarum*, 621–625; *Amphispiza belli*, 499–505, 798–799; *A. bilineata*, 161–162, 165; *Anabacerthia variegaticeps*, 379; *Aphelocoma coerulescens*, 166; *Ara ambiguus*, 345, 348; *A. severa*, 345; *Aramides wolfi*, 345, 348; *Arremon aurantiirostris*, 348; *Atlapetes brunneinucha*, 378, 380–386; *Attila spadiceus*, 347, 378; *A. torridus*, 347, 350; *Aulacorhynchus haematopygus*, 346, 350; *Auriparus flaviceps*, 161–162; *Automolus ochrolaemus*, 346; *A. rubiginosus*, 346; *Barythrhengus ruficapillus*, 346; *Basileuterus culicivorus*, 378, 380–386; *B. rivularis*, 347; *B. tristriatus*, 378, 380, 383–388; *Branta canadensis*, 244–245; *Buthraupis eximia*, 348, 350; *Cacicus uropygialis*, 347; *Calcarius lapponicus*, 131–142; *Calidris alba*, 551–561; *C. pusilla*, 56–67; *Calypte costae*, 161–162, 165; *Campylorhamphus pusillus*, 379, 380; *C. trochilirostris*, 346; *Campylorhynchus brunneicapillus*, 162, 166; *Carpodacus mexicanus*, 162; *Carpodectes hopkei*, 347, 349; *Cathartes frantzii*, 379; *C. fuscater*, 378; *C. mexicanus*, 378; *C. ustulatus*, 347, 351; *Celeus loricatus*, 346; *Centrocercus urophasianus*, 499–505; *Cephalopterus penduliger*, 347, 349; *Cercomacra tyrannia*, 347; *Cerorhinca monocerata*, 147;

- Chlorophanes spiza*, 348; *Chlorophonia callophrys*, 378; *Chlorospingus ophthalmicus*, 378, 380–386; *Circus cyaneus*, 370–374; *Coereba flaveola*, 378; *Colaptes auratus*, 166; *Colonia colonus*, 347, 350; *Columba cayennensis*, 345; *C. goodsoni*, 345; *C. subvinacea*, 345; *Cranioleuca erythropsis*, 378, 380–386; *Crypturellus cinereus*, 345; *C. soui*, 345; *Cyanerpes caeruleus*, 348; *C. cyaneus*, 348, 350; *Cyanolyca cucullata*, 378; *Cymbilaimus lineatus*, 347; *Cyphorhinus arada*, 347; *Dacnis venusta*, 348, 350; *Damophila julie*, 346; *Daptrius americanus*, 345, 348; *Dendrocincia fuliginosa*, 346; *Dendroica caerulescens*, 757–758, 760, 762; *D. castanea*, 600–603, 757–758, 760–764; *D. cerulea*, 757; *D. coronata*, 499–505, 757–758, 760–764; *D. discolor*, 757, 760; *D. dominica*, 757–758, 760, 762–763; *D. fusca*, 757–758, 760, 763; *D. kirtlandii*, 757, 760–762; *D. magnolia*, 757–758, 760, 763; *D. palmarum*, 757–758, 760–764; *D. pensylvanica*, 757–758, 760, 763; *D. petechia*, 757, 760–761; *D. pinus*, 757, 761–763; *D. striata*, 757–758, 760–761; *D. tigrina*, 757–758, 760–764; *D. virens*, 379, 383, 757–758, 760, 763; *Dryoscopus lineatus*, 346; *Dysithamnus mentalis*, 347, 379, 380; *Electron platyrhynchum*, 346; *Empidonax flavescens*, 378; *E. minimus*, 265–283; *Eremophila alpestris*, 499–505, 623; *Euphagus cyanocephalus*, 499–505; *Euphonia minuta*, 348, 350; *E. saturata*, 348, 350; *Eutoxeres aquila*, 346; *Falco sparverius*, 499–505; *Formicarius nigricapillus*, 347; *Fulica americana*, 710–711; *Gelochelidon nilotica*, 202–203; *Geococcyx californianus*, 166; *Geranospiza caerulescens*, 345, 349; *Glaucis aenea*, 346; *Glyphorynchus spirurus*, 346, 378; *Grallaria perspicillata*, 347; *Gymnopithys leucaspis*, 347; *Haliaeetus leucocephalus*, 413–414; *Harpia harpyja*, 345, 348–349; *Heliodoxa jacula*, 346; *Heliothryx barroti*, 346; *Hemithraupis guira*, 348; *Henicorhina leucophrys*, 378, 380–386; *Heterospingus xanthopygius*, 348; *Hirundo rustica*, 499–505; *Hylotistes subulatus*, 346; *Hylophilus decurtatus*, 379; *H. minor*, 347; *Hylophylax naevioides*, 347; *Icteria virens*, 499–505; *Icterus galbula*, 499–505; *I. mesomelas*, 347; *Iridophanes pulcherrima*, 348, 350; *Junco hyemalis*, 499–505, 532–536, 719–721; *Lanius ludovicianus*, 161–162, 798–799; *Larus argentatus*, 694–703; *L. delawarensis*, 806–807; *Leptodon cayanensis*, 345; *Leptopogon superciliaris*, 347; *Leptotila pallida*, 345; *Leucopternis semiplumbea*, 345; *Lipaugus unirufus*, 347, 350; *Lophortyx californicus*, 161–162, 166; *Lophotriccus pileatus*, 347; *Lunda cirrhata*, 147; *Lysurus crassirostris*, 379; *Malacoptila panamensis*, 346, 350; *Manacus manacus*, 347; *Melanerpes pucherani*, 346; *Melospiza melodia*, 499–505; *Micrastur ruficollis*, 345, 349; *Microbates cinereiventris*, 347; *Microcerulus marginatus*, 347; *Microrhopias quixensis*, 347; *Mimus polyglottos*, 161–162, 421–422; *Mionectes olivaceus*, 347, 349, 378; *Mitrospingus cassini*, 348; *Mniotilla varia*, 378, 383; *Molothrus ater*, 499–505; *Momotus momota*, 346, 350; *Myadestes melanops*, 378; *Myiarchus cinerascens*, 161; *Myiobius atricaudus*, 347; *M. barbatus*, 347; *Myioborus miniatus*, 378, 380, 382–387; *Myiodynastes maculatus*, 347; *Myiophobus fasciatus*, 347; *Myiotriccus ornatus*, 347; *Myrmeciza exsul*, 347; *M. immaculata*, 347; *Myrmotherula axillaris*, 347; *M. fulviventris*, 347; *M. schisticolor*, 347, 379; *M. surinamensis*, 347; *Nectarinia famosa*, 449, 453; *N. kilimensis*, 449, 453; *N. reichenowi*, 448–461; *N. venusta*, 449, 453; *Notharchus macrorhynchos*, 346, 350; *N. tectus*, 346, 350; *Nyctibius griseus*, 346, 349; *Nystalus radiatus*, 346, 348; *Ocreatus underwoodii*, 346, 350; *Odontophorus erythropsis*, 345; *Oporornis formosus*, 379, 380, 383; *Oreoscoptes montanus*, 798–799; *Ornithon semifalvum*, 347; *Ortalis erythroptera*, 345; *Otus asio*, 319–327; *O. guatemalae*, 345; *Pachyramphus albogriseus*, 347; *P. cinnamomeus*, 347; *P. versicolor*, 379; *Pandion haliaetus*, 408–412, 413–414; *Parabuteo unicinctus*, 785–786; *Parula pitiayumi*, 347; *Parus atricapillus*, 40–55; *Passerulus sandwichensis*, 391–401, 623–624; *Penelope purpurascens*, 345, 348–349; *Perdix perdix*, 499–505; *Phaethornis longuemareus*, 346; *P. superciliosus*, 346; *P. yaruqui*, 346; *Phasianus colchicus*, 499–505; *Philydor erythrocercus*, 346; *Phloeoceastes guayaquilensis*, 346; *P. melanoleucus*, 346, 348; *Piaya cayana*, 345; *Pico pica*, 499–505; *Picoides nuttallii*, 418–420; *P. scalaris*, 161–162; *Piculus leucolaemus*, 346; *P. rubiginosus*, 346; *P. simplex*, 378; *Picumnus olivaceus*, 346; *Pionopsitta haematonotis*, 345; *Pionus chalcopterus*, 345; *P. menstruus*, 345; *P. sordidus*, 345, 350; *Pipra coronata*, 347, 350; *P. mentalis*, 347; *Pipromorpha oleaginea*, 347; *Piranga ludoviciana*, 499–505; *P. rubra*, 379; *Pittasoma rufopileatum*, 346, 350; *Pitylus grossus*, 348; *Platycichla leucops*, 347; *Platyrinchus coronatus*, 347, 349; *P. mystaceus*, 347, 378, 380; *Poeocilotrichus ruficeps*, 347, 350; *Pogonotriccus ophthalmicus*, 347, 350; *Polioptila caerulea*, 161; *P. melanura*, 161; *P. schistaceigula*, 347; *Pooecetes gramineus*, 499–505, 622–624, 803–804; *Popelairia conversii*, 346; *Premnoplex brunneus*, 346, 349, 378, 380–387; *Pteroglossus saquineus*, 346; *Ptychoramphus aleu-*

- ticus*, 143–151; *Pulsatrix perspicillata*, 345, 348; *Pygoscelis adeliae*, 675–681; *P. antarctica*, 675–681; *Pyriglena leuconota*, 347; *Querula quelea*, 516–524; *Querula purpurata*, 347; *Quiscalus mexicanus*, 789; *Ramphastos ambiguus*, 346, 348; *R. brevis*, 346, 348; *Ramphocænus melanurus*, 347; *Regulus calendula*, 499–505; *Rhynchoscyrus brevirostris*, 347, 378; *R. fulvipectus*, 347, 350; *Salpinctes obsoletus*, 499–505; *Schiffornis turdinus*, 347, 349; *Sclerurus guatemalensis*, 346; *S. mexicanus*, 346, 349; *Semnornis frantzii*, 378; *Setophaga ruticilla*, 265–283; *Sialia sialis*, 562–572; *Sirystes sibilator*, 347, 350; *Sittasomus griseicapillus*, 378, 380–387; *Spizella breweri*, 798–799; *S. pusilla*, 623; *Sturna albifrons*, 37–38; *Sturnella magna*, 622–623; *S. neglecta*, 499–505; *Sturnus vulgaris*, 499–505, 806–807; *Syndactyla subalaris*, 378, 380–387; *Tachyphonus delatrii*, 348; *T. luctuosus*, 348; *Tangara dowii*, 379; *T. gyrola*, 348; *T. icterocephala*, 348, 350, 378; *T. lavinia*, 348, 350; *T. nigroviridis*, 348, 350; *Taraba major*, 347; *Teronotriccus erythrurus*, 347; *Thalurania furcata*, 346; *Thamnophilus punctatus*, 347, 351; *Threnetes ruckeri*, 346; *Thripadectes rufobrunneus*, 379; *Tinamus major*, 345, 348–349; *Todirostrum nigriceps*, 347; *Tolmomyias assimilis*, 347; *Touit dilectissima*, 345, 350; *Troglodytes ochraceus*, 378; *Trogon collaris*, 346; *T. melanurus*, 346; *T. viridis*, 346; *Turdus migratorius*, 499–505; *T. nudigenis*, 347; *T. obsoletus*, 347; *Tryanniscus nigrocipillus*, 347, 350; *T. viridiflavus*, 347, 350; *Tyrannulus elatus*, 347; *Tyrannus tyrannus*, 499–505; *Tyto alba*, 620; *Veniliornis kirkii*, 346; *Verminivora chrysoptera*, 379, 380, 383; *Vireo olivaceus*, 347; *Wilsonia pusilla*, 379, 383, 499–505; *Xenops minutus*, 346; *X. rutilans*, 346; *Xiphorhynchus erythropygius*, 346, 378, 380–387; *Zenaida asiatica*, 161–162, 166; *Z. macroura*, 499–505; *Zonotrichia leucophrys*, 499–505
- Ecuador, 187–189, 343–352
- Egger, Mark, Varied Thrushes feeding on talitrid amphipods, 805–806
- Eggs, collections, 746–755; composition, 407–408; density, specific gravity, and volume, 207–211; laying, 57–59, 245, 247–250, 365–368, 413, 483–488, 570–571, 589–590, 676–677, 694–703, 710, 724, 776; temperature, 100–105, 626–627
- Eggs, descriptions and measurements, *Aechmophorus occidentalis*, 578; *Aegelaius phoeniceus*, 208–211; *Aix sponsa*, 74; *Amazonetta brasiliensis*, 74; *Anas platyrhynchos*, 74; *Anous stolidus*, 74; *A. tenuirostris*, 74; *Anser fabalis*, 74; *Buteo jamaicensis*, 74; *Calidris pu-*
- silla*, 59; *Chrysolophus amherstiae*, 74; *C. pictus*, 74; *Conopias inornata*, 770–771; *Dromaius novaehollandiae*, 74; *Eudyptes chrysocome*, 101–104; *E. chrysolophus*, 101; *Fulica americana*, 74; *Gygis alba*, 74; *Hirundo rustica*, 208–211; *Larus argentatus*, 697–698; *L. delawarensis*, 3, 7; *L. heermanni*, 74; *L. occidentalis*, 74; *Lophophorus impejanus*, 74; *Metriopelia aymara*, 480; *Molothrus ater*, 209–210; *Numida meleagris*, 74; *Oceanodroma leucorhoa*, 626–627; *Pavo muticus*, 74; *Pelecanus occidentalis*, 74, 407–408; *P. onocrotalus*, 407–408; *Phaethon rubricauda*, 74; *Phainopepla nitens*, 199; *Phainoptila melanoxantha*, 198–199; *Phasianus colchicus*, 74; *Pica pica*, 192–195; *Pterocincus pennata*, 74; *Ptilogonyx caudatus*, 199; *P. cinereus*, 199; *Pygoscelis adeliae*, 101; *P. papua*, 101; *Quiscalus quiscula*, 209–210; *Rissa tridactyla*, 74; *Sayornis phoebe*, 209–210; *Spheniscus demersus*, 101; *Struthio camelus*, 74; *Sturnus vulgaris*, 209–210, 407–408; *Sula leucogaster*, 74; *Syrmaticus soemmerringii*, 74; *Turdus migratorius*, 74, 208–211; *Xanthocephalus xanthocephalus*, 192–195
- Eisenmann, E., reviews by, 645–646
- Electron platyrhynchum*, 346
- Elliot, R. D., and R. I. G. Morrison, The incubation period of the Yellow Rail, 422–423
- Ellis, David H., and Wayne H. Whaley, Two winter breeding records for the Harris' Hawk, 413
- Ellison, Laurence N., Author's response to Commentary by David C. Duffy, 816–817; see Tremblay, Jacqueline, and ———
- Emlen, John T., Land bird densities on Baja California islands, 152–167
- Empidonax flavescens*, 378
- E. minimus*, 265–283
- E. spp.*, 160
- England, 537–543
- Environment, influence on *Aegelaius phoeniceus*, 360–361; *Alectoris chukar*, 499–505; *Amphispiza belli*, 499–505; *Ardea herodias*, 628–630; *Bonasa umbellus*, 315; *Branta canadensis*, 243–250; *Calidris pusilla*, 58, 61–63, 66; *Canachites canadensis*, 315; *Centrocercus urophasianus*, 499–505; *Cerorhinca monocerata*, 147–150; *Charadrius vociferus*, 604–606; *Dendroica coronata*, 499–505; *Eremophila alpestris*, 499–505; *Euphagus cyanocephalus*, 499–505; *Falco sparverius*, 499–505; *Gymnorhinus cyanocephalus*, 797; *Hirundo rustica*, 499–505; *Icteria virens*, 499–505; *Icterus galbula*, 499–505; *Junco hyemalis*, 499–505, 532–536, 720–721; *Lagopus lagopus*, 106–115, 314–315, 682–687; *L. leucurus*, 314–315; *L. mutus*, 314–315; *Lophortyx californicus*, 499–505; *Lunda cirrhata*, 150, 793–794; *Melospiza*

- melodia*, 499–505; *Mimus longicaudatus*, 724; *Molothrus ater*, 499–505; *Nesomimus parvulus*, 724; *Otus asio*, 795–796; *Parabuteo unicinctus*, 780; *Perdix perdix*, 499–505; *Phasianus colchicus*, 499–505; *Phoenicoparrus andinus*, 333; *P. jamesi*, 333; *Phoenicopterus chilensis*, 333; *Pica pica*, 499–505; *Piranga ludoviciana*, 499–505; *Pooecetes gramineus*, 499–505; *Ptychoramphus aleuticus*, 147–150; *Quelea quelea*, 516–524; *Regulus calendula*, 499–505; *Salpinctes obsoletus*, 499–505; *Sturnella neglecta*, 499–505; *Sturnus vulgaris*, 499–505; *Turdus migratorius*, 499–505; *Tyrannus tyrannus*, 499–505; *Wilsonia pusilla*, 499–505; *Zenaida macroura*, 499–505; *Zonotrichia leucophrys*, 499–505
- Eremophila alpestris*, 392, 499–505, 623
- Erickson, W. A., see Jaeger, M. M., et al.
- Ethiopia, 516–524
- Eudyptes chrysocome*, 100–105  
*E. chrysolophus*, 101
- Euphagus carolinus*, 392
- E. cyanocephalus*, 499–505
- Euphonia affinis*, 184
- E. hirundinacea*, 184
- E. minuta*, 348, 350
- E. saturata*, 348, 350
- Eurynorhynchus pygmeus*, 189
- Eutoxeres aquila*, 346
- Evolution of behavior, 764–765; breeding characteristics, 733–734, 764–765, 783–787; clutch size, 416; competitive abilities, 127–128; dispersal, 54–55; heart-body size ratios, 316–317; large size, 127–128; locomotor skills, 127–128; plumage variability, 86–92, 94–99, 548–549; rapid development, 127–128; status-signaling, 86–92, 94–99; territoriality, 559–560
- Evolution of *Aechmophorus occidentalis*, 584; *Chen rossii*, 548–549; *Cyanocorax sanblasiiana*, 200–201; *Dendroica castanea*, 764; *D. coronata*, 764; *D. palmarum*, 764; *D. pensylvanica*, 764; *D. pinus*, 764; *D. striata*, 764; *D. tigrina*, 764; **Hemitriccus cinnamomeipectus**, 446–447; *H. kaempferi*, 446–447; *H. mirandae*, 446–447; hybrids of *Chen caerulescens* and *C. rossii*, 548–549; hybrids of *Tyrannus couchii* and *T. melancholicus*, 231; *Junco hyemalis*, 86–88, 94–99; *Pandion haliaetus*, 416; *Tyrannus couchii*, 231; *T. melancholicus*, 231; *Zonotrichia querula*, 88–89
- Falco femoralis*, 476
- F. peregrinus*, 662–674
- F. rusticolus*, 669–672
- F. sparverius*, 476, 479, 499–505
- Falcon, Peregrine, 662–674
- Feathers (see Molts and plumages)
- Finch, House, 159, 462–474; Sooty-faced, 379; Warbler, 800–802
- Firewood-gatherer, 607
- Fitzner, R. E., see Rotenberry, J. T., et al.
- Fitzpatrick, John W., and John P. O'Neill, A new tody tyrant from northern Peru, 443–447
- Fjeldså, Jon, Guide to the young of European pre-social birds (rev.), 214–215
- Flamingo, American, 615–619
- Flatbill, Eye-ringed, 378
- Fledging, success of (see Breeding success)
- Flicker, Gilded, 158
- Florida, 410–411, 415–417
- Florida caerulea*, 116–130
- Flycatcher, Ash-throated, 158; Least, 265–283; Olive-striped, 378; White-bearded, 767–775; Yellowish, 378
- Foliage-gleaner, Lineated, 378, 380–387; Spectacled, 379
- Food and feeding habits of *Accipiter superciliosus*, 349; *Agriornis montana*, 480; *Amazona albifrons*, 183–186; *Amphispiza bilineata*, 161–162; *Anabacerthia variegaticeps*, 379; *Anas rubripes*, 737–741; *Aratinga astec*, 183–186; *Ardea herodias*, 628–630; *Atlapetes brunneinucha*, 378; *Attila spadiceus*, 378; *Auriparus flaviceps*, 161–162; *Basileuterus culicivorus*, 378; *B. tristriatus*, 378; *Branta canadensis*, 245–246; *Buteo poecilochrous*, 479; *Calcarius lapponicus*, 132, 136–139; *Calidris alba*, 551–561; *C. pusilla*, 62–64; *Calypte costae*, 161–162; *Campylorhynchus pusillus*, 379; *Campylorhynchus brunneicapillus*, 162; *Carpodacus mexicanus*, 162; *Carpodectes hopkei*, 349; *Catharus frantzii*, 379; *C. fuscater*, 378; *C. mexicanus*, 378; *Centurus aurifrons*, 183–186; *C. pygmaeus*, 183–186; *Cephalopterus penduliger*, 349; *Chen caerulescens*, 70–72; *Chlorophonia callophrys*, 378; *Chlorospingus ophthalmicus*, 378; *Circus cyaneus*, 370–374; *Cissiophaga yucatanica*, 183–186; *Coereba flaveola*, 378; *Conopias inornata*, 768; *Cotinga amabilis*, 183–186; *Coturnix coturnix*, 28–29; *Cranioleuca erythrops*, 378; *Cranolyca cucullata*, 378; *Cyanocorax yncas*, 183–186; *Dendroica caerulescens*, 759; *D. castanea*, 600–603, 759; *D. coronata*, 759, 761; *D. discolor*, 759; *D. dominica*, 759, 762; *D. fusca*, 759; *D. kirtlandii*, 759, 762; *D. magnolia*, 759; *D. palmarum*, 759, 761–762; *D. pensylvanica*, 759; *D. petechia*, 759; *D. pinus*, 759, 762; *D. striata*, 759; *D. tigrina*, 759, 761; *D. virens*, 379, 759; *Dives dives*, 183–186; *Dryocopus pileatus*, 592; *Dysithamnus mentalis*, 379; *Empidonax flavescens*, 378; *E. minimus*, 269–281; *Eudyptes chrysocome*, 104; *Euphonia affinis*, 183–186; *E. hirundinacea*, 183–186; *Florida*

- caerulea*, 118–127; *Fulica americana*, 710–711; *Geospiza difficilis*, 730; *G. magnirostris*, 730; *Geranospiza caerulescens*, 349; *Glyphorynchus spirurus*, 378; *Haliaeetus leucocephalus*, 413–414; **Hemitriccus cinnamomeipectus**, 445; *Henicorhina leucophrys*, 378; *Hylophilus decurvatus*, 379; *Icterus auratus*, 183–186; *I. chrysater*, 183–186; *I. gularis*, 183–186; *I. mesomelas*, 183–186; *I. prosthemelas*, 183–186; *Ixoreus naevius*, 805; *Lanius ludovicianus*, 161–162, 798–799; *Larus delawarensis*, 806–807; *Lophortyx californicus*, 161–162; *Lysurus crassirostris*, 379; *Megarynchus pitangua*, 183–186; *Menura superba*, 489; *Micrastur ruficollis*, 349; *Mimus polyglottos*, 161–162, 421–422; *Mionectes olivaceus*, 378; *Mniotilla varia*, 378; *Myadestes melanops*, 378; *Myiarchus cinerascens*, 161; *Myioborus miniatus*, 378; *Myiozetetes similis*, 183–186; *Myrmotherula schisticolor*, 379; *Nectarinia famosa*, 449–450, 453, 455, 457; *N. kilimensis*, 449–450, 455; *N. reichenowi*, 448–461; *N. venusta*, 449–450, 453, 455; *Neosomimus parvulus*, 726, 728–732; *Oporornis formosus*, 379; *Ornithodoros vetula*, 183–186; *Otus asio*, 319–327; *Pachyramphus versicolor*, 379; *Pandion haliaetus*, 408–412, 413–414, 415–417; *Parabuteo unicinctus*, 779–785; *Passerculus sandwichensis*, 399; *Phoenicopterus ruber*, 615–616; *Picoides nuttallii*, 418–420; *P. scalaris*, 161–162; *Piculus simplex*, 378; *Pinguinus impennis*, 790–792; *Piranga rubra*, 379; *Pitangus sulphuratus*, 183–186; *Platyrinchus mystaceus*, 378; *Polioptila caerulea*, 161; *P. melanura*, 161; *Premnoplex brunneus*, 378; *Psilorhinus morio*, 183–186; *Pteroglossus torquatus*, 183–186; *Pygoscelis adeliae*, 680; *P. antarctica*, 680; *Quelea quelea*, 516–517, 523; *Quiscalus quiscula*, 608–609; *Ramphastos sulfuratus*, 183–186; *Rhynchocyclus brevirostris*, 378; *Saltator atriceps*, 183–186; *Semnornis frantzii*, 378; *Setopaga ruticilla*, 269–281; *Sittasomus griseicapillus*, 378; *Sphyrapicus varius*, 195; *Stearornis caripensis*, 188; *Sterna hirundo*, 28–29; *Sturnus vulgaris*, 28–29, 806–807; *Syndactyla subalaris*, 378; *Tangara dowii*, 379; *T. icterocephala*, 378; *Thripadectes rufobrunneus*, 379; *Tityra semifasciata*, 183–186; *Troglodytes ochraceus*, 378; *Trogon citreolus*, 183–186; *Turdus grayi*, 183–186; *Tyrannus melancholicus*, 183–186; *Tyto alba*, 619–621; *Vermivora chrysopatra*, 379; *V. peregrina*, 186–187; *Wilsonia pusilla*, 379; *Xiphorhynchus erythropygius*, 378; *Zenaidura asiatica*, 161–162
- Foraging (see Food and feeding habits; Behavior, feeding)
- Formicarius nigricapillus*, 347
- Forshaw, Joseph M., Parrots of the world (rev.), 829–830
- Forsman, Eric D., and Howard M. Wight, Allopreening in owls: what are its functions?, 525–531
- Fraga, Rosendo M., Helpers at the nest in passerines from Buenos Aires Province, Argentina, 606–608
- French, Thomas W., Great Horned Owl predation on Leach's Storm-Petrels in Maine, 202
- Friedmann, Herbert, review by, 640–641
- Fringilla coelebs*, 91, 253–264
- Fulica americana*, 74, 476, 479–480, 704–713
- F. ardesiaca*, 476
- F. gigantea*, 476, 479
- Furnarius rufus*, 607
- Gabaldon, Diana J., see Clark, L., and —
- Galápagos Islands, 723–736, 800–802
- Gallinago (Capella) gallinago*, 476
- Gallinula chloropus*, 476, 479
- Gallinule, Common, 476–479
- Gauthreaux, Sidney A., Jr., Commentary: Priorities in bird migration studies, 813–815; review by, 818–819
- Gavrilov, Valery M., see Dolnik, Victor R., and —
- Geffen, Alice M., A birdwatcher's guide to the eastern United States (rev.), 648
- Gelochelidon nilotica*, 202–203
- Genetic variation in (see Genetics, variation)
- Genetics, gene flow, 163–164; variation, 548–549, 581
- Genetics of *Aechmophorus occidentalis*, 581
- Geococcyx californianus*, 158, 166, 784
- Geographic variation in molts and plumages, 575, 577, 585; nesting coloration, 800–802; sex ratio, 516–524, 532–536; vocalizations, 423–425, 462–474
- Geographic variation of *Aechmophorus occidentalis*, 575, 577, 585; *Carpodacus mexicanus*, 462–474; *Certhidea olivacea*, 800–801; *Cyanocitta cristata*, 423–425; *Geospiza conirostris*, 800–801; *G. difficilis*, 800–801; *G. fortis*, 800–801; *G. fuliginosa*, 800–801; *G. magnirostris*, 800–801; *G. scandens*, 800–801; *Junco hyemalis*, 532–536; *Quelea quelea*, 516–524; *Tyrannus couchii*, 227–232; *T. melancholicus*, 227–232
- Georgia, 417–418, 587–594
- Geositta cunicularia*, 477
- G. tenuirostris*, 477
- Geospiza conirostris*, 800–802
- G. difficilis*, 726, 729–730, 800–802
- G. fortis*, 800–802
- G. fuliginosa*, 800–802

- G. magnirostris*, 730, 800–802  
*G. scandens*, 800–802  
*Geothlypis semiflava*, 351  
*Geranospiza caerulescens*, 345, 349  
Géroudet, Paul, Grands Échassiers Gallinacés  
 Râles d'Europe [Wading birds, Galliformes, and rails of Europe] (rev.), 644  
Gill, Frank B., and Larry L. Wolf, Nectar loss by  
 Golden-winged Sunbirds to competitors, 448–  
 461  
Gill, Robert, Jr., and Gerald A. Sanger, Tufted  
 Puffins nesting in estuarine habitat, 792–794  
*Glaucis aenea*, 346  
*Glyphorynchus spirurus*, 346, 378  
Gnatcatcher, Masked, 608  
*Gnorimopsar chopi*, 179–182  
Gochfeld, Michael, see Burger, Joanna, and ———  
Godwin, Derek, Birds of man's world (rev.), 437–  
 438; Pigeons and doves of the world (rev.), 646  
Goose, Canada, 234–252; Lesser Snow, 68–72,  
 544–550; Ross', 544–550  
Gore, M. E. J., and A. R. M. Gepp, Las Aves del  
 Uruguay (rev.), 645  
Goslow, G. E., Jr., review by, 637  
Gackle, Common, 209–210, 608–609; Great-  
 tailed, 789  
*Grallaria perspicillata*, 347  
Grant, Nicola, see Grant, P. R., and ———  
Grant, P. R., and Nicola Grant, Breeding and  
 feeding of Galápagos Mockingbirds, *Nesomimus parvulus*, 723–736; P. T. Boag, and D.  
 Schlüter, A bill color polymorphism in young  
 Darwin's finches, 800–802; review by, 819–  
 821  
Grebe, Pied-billed, 706; Western, 573–586; White-  
 tufted, 476, 478  
Greenberg, Russell, Body size, breeding habitat,  
 and winter exploitation systems in *Dendroica*,  
 756–766  
Greenland, 670  
Greenlet, Lesser, 379  
Greenway, James C., Jr., Type specimens of birds  
 in the American Museum of Natural History.  
 Part 2. Otididae . . . Picidae (rev.), 642–643  
Grosbeak, Black-headed, 160; Evening, 90–91  
Ground-Dove, Bare-faced, 477, 479; Black-winged,  
 477, 480; Golden-spotted, 477, 480  
Grouse, Blue, 308, 316; Ruffed, 308–309, 316;  
 Sage, 499–505; Sharp-tailed, 308, 316; Spruce,  
 309–310, 316  
Growth and development of *Aechmophorus occidentalis*, 581–582; *Aix sponsa*, 296–305; *Anas rubripes*, 737–745; *Conopias inornata*, 773–  
 774; *Coturnix coturnix*, 10–30, 300–303; *Florida caerulea*, 116–130; *Gymnorhinus cyanocephalus*, 797; *Larus argentatus*, 809–812; *L. delawarensis*, 4, 284–295; *Pandion haliaetus*,  
 416; *Passerculus sandwichensis*, 395–399; *Sturnus vulgaris*, 10–30  
Guatemala, 227–229  
Gull, California, 6–9; Herring, 203, 264, 694–703,  
 809–812; Laughing, 203; Ring-billed, 1–5, 6–  
 9, 284–295, 806–808  
Gulledge, James L., review by, 632  
Guyana, 425  
*Gygis alba*, 74  
*Gymnopithys leucaspis*, 347  
*Gymnorhinus cyanocephalus*, 796–798  
Gyrfalcon, 669, 672  
Habitat of *Accipiter superciliosus*, 345, 349; *Aces-trura bombus*, 346, 350; *Aechmophorus occidentalis*, 574; *Agelaius phoeniceus*, 621–623;  
*A. thilius*, 475; *Agriornis albicauda*, 475; *A. montana*, 475; *Alectoris chukar*, 499–505; *Al-locotopterus deliciosus*, 347, 350; *Amazilia amabilis*, 346; *A. franciae*, 346; *Amazona farinosa*, 345; *A. festiva*, 345, 350; *Ammodramus henslowii*, 621–623; *A. savannarum*, 621–625; *Amphispiza belli*, 499–505; *A. bilineata*, 162,  
 165; *Anabacerthia variegaticeps*, 376–377;  
*Anas cyanoptera*, 475; *A. flavirostris*, 475; *A. georgica*, 475; *A. rubripes*, 737–745; *A. ver-sicolor*, 475; *Anthus correndera*, 475; *A. fur-catus*, 475; *Aphelocoma coerulescens*, 166; *Ara ambiguua*, 345, 348; *A. severa*, 345; *Aramides wolfi*, 345, 348; *Arremon aurantiirostris*, 348;  
*Asthenes dorbignyi*, 475; *A. wyatti*, 475; *At-lapetes brunneinucha*, 376–377; *Attila spadi-ceus*, 347, 376–377; *A. torridus*, 347, 350; *Aulacorhynchus haematocephalus*, 346, 350;  
*Auriparus flaviceps*, 162; *Automolus ochrolae-mus*, 346; *A. rubiginosus*, 346; *Baryphthengus ruficapillus*, 346; *Basileuterus culicivorus*, 376–377; *B. rivularis*, 347; *B. tristriatus*, 376–377; *Bolborhynchus aurifrons*, 475; *B. orbygnesius*, 475; *Bubo virginianus*, 475; *Buteo poecilochrous*, 475; *Buthraupis excimia*, 348, 350; *Cacicu uropygialis*, 347; *Calcarius lapponicus*, 131–132, 135; *Calidris bairdii*, 475; *C. melanotos*, 475; *C. pusilla*, 56–57; *Calypte costae*, 162, 165; *Campylorhynchus pu-sillus*, 376–377; *C. trochilirostris*, 346; *Cam-pylorhynchus brunneicapillus*, 162, 166;  
*Carpodacus mexicanus*, 162; *Carpodectes hopkei*, 347, 349; *Casmerodus albus*, 475; *Catamenia inornata*, 475; *Cathartes aura*, 475;  
*Catharus frantzii*, 376–377; *C. fuscater*, 376–  
 377; *C. mexicanus*, 376–377; *C. ustulatus*, 347, 351; *Celeus loricatus*, 346; *Centrocercus urophasianus*, 499–505; *Cephalopterus pen-duliger*, 347, 349; *Cercomacra tyrannia*, 347;  
*Cerorhinca monocerata*, 145, 147–150; *Chloe-phaga melanoptera*, 475; *Chlorophanes spiza*,

- 348; *Chlorophonia callophrys*, 376–377; *Chlorospingus ophthalmicus*, 376–377; *Cinclodes atacamensis*, 475; *C. fuscus*, 475; *Circus cyaneus*, 475; *Cnemarchus (=Xolmis?) rufipennis*, 475; *Coereba flaveola*, 376–377; *Colaptes auratus*, 166; *C. rupestris*, 475; *Colibri coruscans*, 475; *Colonia colonus*, 347, 350; *Columba cayennensis*, 345; *C. goodsoni*, 345; *C. subvinacea*, 345; *Conopias inornata*, 767; *Craniolæuca erythrops*, 376–377; *Crypturellus cinereus*, 345; *C. soui*, 345; *Cyanerpes caeruleus*, 348; *C. cyaneus*, 348, 350; *Cyanolyca cucullata*, 376–377; *Cymbilaimus lineatus*, 347; *Cyphorhinus arada*, 347; *Dacnis venusta*, 348, 350; *Damophila julie*, 346; *Daptrius americanus*, 345, 348; *Dendrocincus fuliginosus*, 346; *Dendroica caerulea*, 757–758; *D. castanea*, 600, 602, 757–758; *D. cerulea*, 757; *D. coronata*, 499–505, 757–758; *D. discolor*, 757; *D. dominica*, 757–758; *D. fusca*, 757–758; *D. kirtlandii*, 757; *D. magnolia*, 757–758; *D. palmarum*, 757–758; *D. pensylvanica*, 757–758; *D. petechia*, 757; *D. pinus*, 757; *D. striata*, 757–758; *D. tigrina*, 757–758; *D. virens*, 376–377, 757–758; *Diuca speculifera*, 475; *Dryocopus lineatus*, 346; *Dysithamnus mentalis*, 347, 376–377; *Electron platyrhynchum*, 346; *Empidonax flavescens*, 376–377; *E. minimus*, 266, 273–274; *Eremophila alpestris*, 499–505, 621–623; *Euphagus cyanocephalus*, 499–505; *Euphonia minuta*, 348, 350; *E. saturata*, 348, 350; *Eutoxeres aquila*, 346; *Falco femoralis*, 475; *F. peregrinus*, 663; *F. sparverius*, 475, 499–505; *Formicarius nigricapillus*, 347; *Fulica americana*, 475; *F. ardesiaca*, 475; *F. gigantea*, 475; *Gallinago (Capella) gallinago*, 475; *Gallinula chloropus*, 475; *Geococcyx californianus*, 166; *Geositta cunicularia*, 475; *G. tenuirostris*, 475; *Geranospiza caerulea*, 345, 349; *Glaucis aenea*, 346; *Glyphorynchus spirurus*, 346, 376–377; *Grallaria perspicillata*, 347; *Gymnopithys leucaspis*, 347; *Harpia harpyja*, 345, 348–349; *Heliodoxa jacula*, 346; *Heliothryx barroti*, 346; *Hemithraupis guira*, 348; *Hemitriccus cinnamomeipectus*, 443, 445; *H. kaempferi*, 443; *H. mirandae*, 443; *Henicorhina leucophrys*, 376–377; *Heterospingus xanthopygius*, 348; *Hirundo rustica*, 499–505; *Hylocistes subulatus*, 346; *Hylophilus decurtatus*, 376–377; *H. minor*, 347; *Hylophylax naevioides*, 347; *Icteria virens*, 499–505; *Icterus galbula*, 499–505; *I. mesomelas*, 347; *Iridophanes pulcherrima*, 348, 350; *Junco hyemalis*, 499–505; *Lanius ludovicianus*, 162; *Larus serranus*, 475; *Leptasthenura andicola*, 475; *Leptodon cayanensis*, 345; *Leptopogon superciliaris*, 347; *Leptotila pallida*, 345; *Lessonia rufa*, 475; *Leucopternis semiplumbea*, 345; *Lipaugus unirufus*, 347, 350; *Lophortyx californicus*, 162, 166, 499–505; *Lophotriccus pileatus*, 347; *Lunda cirrhata*, 145, 147, 150, 792–793; *Lysurus crassirostris*, 376–377; *Malacoctila panamensis*, 346, 350; *Manacus manacus*, 347; *Melanerpes pucherani*, 346; *Melospiza melodia*, 499–505; *Menura superba*, 489–490; *Metriopelia ayamara*, 475; *M. ceciliae*, 475; *M. melanoptera*, 475; *Micrastur ruficollis*, 345, 349; *Microbates cinereiventris*, 347; *Microcerulus marginatus*, 347; *Microrhopias quixensis*, 347; *Mimus polyglottos*, 162, 421–422; *Mionectes olivaceus*, 347, 349, 376–377; *Mitrospingus cassini*, 348; *Mniotilla varia*, 376–377; *Molothrus ater*, 484, 486, 499–505; *Momotus momota*, 346, 350; *Muscisaxicola alpina*, 475; *M. rufivertex*, 475; *Myadestes melanops*, 376–377; *Myioibius atricaudus*, 347; *M. barbatus*, 347; *Myioborus miniatus*, 376–377; *Myiodynastes maculatus*, 347; *Myiophobus fasciatus*, 347; *Myiotriccus ornatus*, 347; *Myrmeciza exsul*, 347; *M. immaculata*, 347; *Myrmotherula axillaris*, 347; *M. fulviventer*, 347; *M. schisticolor*, 347, 376–377; *M. surinamensis*, 347; *Notharchus macrorhynchos*, 346, 350; *N. tec-tus*, 346, 350; *Nothoprocta ornata*, 475; *N. pentlandii*, 475; *Nothura maculosa*, 475; *No-tiochelidon murina*, 475; *Nyctibius griseus*, 346, 349; *Nycticorax nycticorax*, 364, 475; *Nystalus radiatus*, 346, 348; *Ochthoeca oen-anthoides*, 475; *Oreocreatus underwoodii*, 346, 350; *Odontophorus erythrops*, 345; *Oporornis formosus*, 376–377; *Oreopholus ruficollis*, 475; *Oreotrochilus estella*, 475; *Ornithion semiflavum*, 347; *Ornithodoros erythroptera*, 345; *Otus guatemalae*, 345; *Oxyura jamaicensis*, 475; *O. sp.*, 475; *Pachyramphus albogriseus*, 347; *P. cinnamomeus*, 347; *P. versicolor*, 376–377; *Pandion haliaetus*, 415; *Parabuteo unicinctus*, 786; *Parula pityayumi*, 347; *Parus atricapillus*, 41–42; *Passerculus sandwichensis*, 392–394, 621–623; *Patagona gigas*, 475; *Penelope purpurascens*, 345, 348–349; *Perdix perdix*, 499–505; *Petrochelidon andecola*, 475; *Phaethornis longuemareus*, 346; *P. superciliosus*, 346; *P. yaruqui*, 346; *Phalacrocorax olivaceus*, 475; *Phalcobaenus megalopterus*, 475; *Phasianus colchicus*, 499–505; *Philydor erythrocercus*, 346; *Phleocryptes melanops*, 475; *Phloeoceastes guayaquilensis*, 346; *P. melanoleucus*, 346, 348; *Phoenicopar-rus andinus*, 329–330; *P. jamesi*, 329–330; *Phoenicopterus chilensis*, 329–330, 475; *Phrygilus alaudinus*, 475; *P. fruticeti*, 475; *P. gayi*, 475; *P. plebejus*, 475; *Piaya cayana*, 345; *Pica pica*, 499–505; *Picoides scalaris*, 162; *Piculus leucolaemus*, 346; *P. rubiginosus*

- sus*, 346; *P. simplex*, 376–377; *Picumnus olivaceus*, 346; *Pionopsitta haematoxis*, 345; *Pionus chalcopterus*, 345; *P. menstruus*, 345; *P. sordidus*, 345, 350; *Pipra coronata*, 347, 350; *P. mentalis*, 347; *Pipromorphaoleaginea*, 347; *Piranga ludoviciana*, 499–505; *P. rubra*, 376–377; *Pittasoma rufopileatum*, 346, 350; *Pitylus grossus*, 348; *Platycichla leucopsis*, 347; *Platyrinchus coronatus*, 347, 349; *P. mystaceus*, 347, 376–377; *Plegadis ridgwayi*, 475; *Podiceps occipitalis*, 475; *P. rolland*, 475; *Poecilotriccus ruficeps*, 347, 350; *Pogonotriccus ophthalmicus*, 347, 350; *Polioptila schistaceigula*, 347; *Pooecetes gramineus*, 499–505, 621–623, 803–804; *Popelairia conversii*, 346; *Premnoplex brunnescens*, 346, 349, 376–377; *Pteroglossus sanguineus*, 346; *Ptychoramphus aleuticus*, 143–151; *Pulsatrix perspicillata*, 345, 348; *Pyriglenaleuconota*, 347; *Quelea quelea*, 516–517; *Querula purpurata*, 347; *Rallus sanguinolentus*, 475; *Ramphastos ambiguus*, 346, 348; *R. brevis*, 346, 348; *Ramphocænus melanurus*, 347; *Regulus calendula*, 499–505; *Rhynchoscyrus brevirostris*, 347, 376–377; *R. fulvipectus*, 347, 350; *Salpinctes obsoletus*, 499–505; *Saltator cinctus*, 610–612; *Schiffornis turdinus*, 347, 349; *Sclerurus guatemalensis*, 346; *S. mexicanus*, 346, 349; *Seumnornis frantzii*, 376–377; *Setophaga ruticilla*, 266, 273–274; *Sicalis uropygialis*, 475; *Sirystes sibilator*, 347, 350; *Sittasomus griseicapillus*, 376–377; *Speotyto (Athene) cunicularia*, 475; *Spinus atratus*, 475; *S. crassirostris*, 475; *Spizella pusilla*, 621–623; *Steatornis caripensis*, 187–188; *Sturnella magna*, 622–623; *S. neglecta*, 499–505; *Sturnus vulgaris*, 499–505; *Syndactyla subalaris*, 376–377; *Tachuris rubrigastra*, 475; *Tachyphonus delatrii*, 348; *T. luctuosus*, 348; *Tangara dowii*, 376–377; *T. gyrola*, 348; *T. icterocephala*, 348, 350, 376–377; *T. lavinia*, 348, 350; *T. nigroviridis*, 348, 350; *Taraba major*, 347; *Terenotriccus erythrurus*, 347; *Thalurania furcata*, 346; *Thamnophilus punctatus*, 347, 351; *Theristicus caudatus*, 476; *Thinocorus orbignyanus*, 475; *Threnetes ruckeri*, 346; *Thripadectes rufobrunneus*, 376–377; *Tinamus major*, 345, 348–349; *Todirostrum nigriceps*, 347; *Tolmomyias assimilis*, 347; *Touit dilectissima*, 345, 350; *Tringa flavipes*, 475; *T. melanoleuca*, 475; *Troglodytes musculus* (=aedon?), 475; *T. ochraceus*, 376–377; *Trogon collaris*, 346; *T. melanurus*, 346; *T. viridis*, 346; *Turdus chiguanco*, 475; *T. migratorius*, 499–505; *T. nudigenis*, 347; *T. obsoletus*, 347; *Tyranniscus nigrocapillus*, 347, 350; *T. viridiflavus*, 347, 350; *Tyrannulus elatus*, 347; *Tyrannus couchii*, 231; *T.* melancholicus, 231; *T. tyrannus*, 499–505; *Tyto alba*, 475; *Upucerthia serrana*, 475; *U. validirostris*, 475; *Uropsalis segmentata*, 475; *Vanellus resplendens*, 475; *Veniliornis kirkii*, 346; *Vermivora chrysoptera*, 376–377; *Vireo olivaceus*, 347; *Vultur gryphus*, 475; *Wilsonia pusilla*, 376–377, 499–505; *Xenops minutus*, 346; *X. rutilans*, 346; *Xiphorhynchus erythrophysius*, 346, 376–377; *Zenaida asiatica*, 162, 166; *Z. auriculata*, 475; *Z. macroura*, 499–505; *Zonotrichia capensis*, 475; *Z. leucocephrys*, 499–505
- Haiti, 615–619
- Haliaeetus leucocephalus*, 413–414, 548
- Hamerstrom, Frances, Effect of prey on predator: voles and Harriers, 370–374
- Hancock, James, and Hugh Elliot, The herons of the world (rev.), 827–828
- Hantzsch, Bernhard, My life among the Eskimos: the Baffinland journals of Bernhard Adolph Hantzsch, 1909–1911 (rev.), 831–832
- Hardy, John William, Molt pattern as a clue to San Blas Jay ancestry, 200–201
- Harpia harpyja*, 345, 348–349
- Harrier, 370–374
- Harrison, Colin, A field guide to the nests, eggs and nestlings of North American birds (rev.), 823–825
- Hatching, success of (see Breeding success)
- Hatching, timing of, 60, 62–67, 116–130, 204, 413, 422–423, 590, 773
- Hawk, Cooper's, 782; Harris', 413, 776–788; Red-tailed, 782; Rough-legged, 669, 672–673; Semiplumbeous, 345; Variable, 476, 479
- Heard, Douglas C., see Calef, George W., and —————
- Heliodoxa jacula*, 346
- Heliothryx barroti*, 346
- Hemithraupis guira*, 348
- Hemitriccus cinnamomeipectus*, 443–447
- H. kaempferi*, 443, 445–447
- H. mirandae*, 443, 445–447
- Henicorhina leucophrys*, 378, 380–386
- Henny, Charles J., and Laurel F. VanCamp, Annual weight cycle in wild Screech Owls, 795–796
- Heron, Black-crowned Night, 364–369; Great Blue, 364, 628–630; Little Blue, 116–130; Yellow-crowned Night, 725
- Hesperiphona vespertina*, 90–91
- Heterospingus xanthopygius*, 348
- Hickey, Joseph J., Coues Award, 212
- Hirundo rustica*, 208–211, 499–505
- Hogue, Janet G., see Marti, Carl D., and —————
- Homberger, Dominique G., reviews by, 432–433, 641–642, 829–830, 834
- Home range (see Territory in)
- Horn, David J., Roger D. Mitchell, and Gordon

- R. Stairs (Eds.), Analysis of ecological systems (rev.), 833
- Horne, Jennifer F. M., see Short, Lester L., and —
- Hornero, Rufous, 607
- Hoyt, Donald F., Practical methods of estimating volume and fresh weight of bird eggs, 73–77
- Hummingbird, Costa's, 158
- Hunt, George L., Jr., see Conover, Michael R., et al.
- Hurlbert, Stuart H., and James O. Keith, Distribution and spatial patterning of flamingos in the Andean altiplano, 328–342
- Hybridization of *Chen caerulescens* and *C. rossii*, 544–550; *Tyrannus couchii* and *T. melancholicus*, 223–227, 231; *Zonotrichia albicollis* and *Z. atricapilla*, 595–597; *Zonotrichia leucophrys* and *Z. querula*, 596–599
- Hyloctistes subulatus*, 346
- Hylophilus decurtatus*, 379
- H. minor*, 347
- Hylophylax naeviooides*, 347
- Ibis, Puna, 476, 479–481
- Icteria virens*, 499–505
- Icterus auratus*, 84
- I. chrysater*, 84
- I. cucullatus*, 159
- I. galbula*, 499–505
- I. gularis*, 84
- I. mesomelas*, 84, 347
- I. prosthemelas*, 84
- Ictinia plumbea*, 351
- Idaho, 798–800
- Incubation in *Bombycilla cedrorum*, 196; *Branta canadensis*, 234–252; *Calidris pusilla*, 56–57; *Charadrius vociferus*, 604–606; *Conopias ornata*, 771; *Coturnicops noveboracensis*, 422–423; *Coturnix coturnix*, 24; *Dendroica castanea*, 601; *Dryocopus pileatus*, 589–590; *Eudynastes chrysocome*, 101–104; *E. chrysophilus*, 101; *Florida caerulea*, 118; *Megadyptes antipodes*, 103; *Menura superba*, 489; *Mimus longicaudatus*, 725–726; *Nesomimus parvulus*, 725–726; *Oceanodroma leucorhoa*, 625–627; *Parabuteo unicinctus*, 413, 778–781; *Parus atricapillus*, 43; *Pygoscelis adeliae*, 101; *P. papua*, 101; *Spheniscus demersus*, 101; *Sterna hirundo*, 24; *Sturnus vulgaris*, 24; *Tyto alba*, 204
- Indian Ocean, 100–105
- Indiana, 533
- Innes, Richard C., Club-tipped feathers in some South American tanagers, 808–809
- Iowa, 704–713
- Iridophanes pulcherrima*, 348, 350
- Islands, biogeography of, 152–167, 350–352
- Ixoreus naevius*, 805–806
- Jackson, Jerome A., see Rogers, David T., Jr., et al.; see Schardien, Bette J., and —
- Jaeger, M. E., see Jaeger, M. M., et al.
- Jaeger, M. M., W. A. Erickson, and M. E. Jaeger, Sexual segregation of Red-billed Queleas (*Quelea quelea*) in the Awash River Basin of Ethiopia, 516–524
- Jaeger, Parasitic, 135; Pomarine, 135
- Jaksic, Fabian, and José L. Yáñez, The diet of the Barn Owl in central Chile and its relation to the availability of prey, 619–621
- Janovy, John, Jr., Keith County journal (rev.), 646
- Jay, Azure-hooded, 378; Blue, 423–425; Brown, 200–201; Piñon, 796–798; San Blas, 200–201; Scrub, 158; Tufted, 200–201
- Jehl, J. R., Jr., review by, 214–215
- Jellis, Rosemary, Bird sounds and their meaning (rev.), 632
- Jenkins, J. Mark, Foraging behavior of male and female Nuttall Woodpeckers, 418–420
- Johnsgard, Paul A., Ducks, geese, and swans of the world (rev.), 643
- Johnson, Douglas H., Estimating nest success: the Mayfield method and an alternative, 651–661
- Jones, P. J., Variability of egg size and composition in the Great White Pelican (*Pelecanus onocrotalus*), 407–408
- Juncos, Dark-eyed, 78–88, 94–99, 499–505, 532–536, 630–631, 714–722
- Junco hyemalis*, 78–88, 94–99, 499–505, 532–536, 630–631, 714–722
- Kale, Herbert W., II, editor of Recent Literature, 1A–32A, 1B–32B, 1C–32C, 1D–32D
- Kantak, Gail E., Observations on some fruit-eating birds in Mexico, 183–186
- Karr, James R., review by, 434–435
- Keith, James O., see Hurlbert, Stuart H., and —
- Kemp, Thomas R., see Tramer, Elliot J., and —
- Kenya, 448–461
- Kestrel, American, 476, 479, 499–505
- Ketterson, Ellen D., Status signaling in Dark-eyed Juncos, 94–99; and Val Nolan, Jr., Seasonal, annual, and geographic variation in sex ratio of wintering populations of Dark-eyed Juncos (*Junco hyemalis*), 532–536
- Keve, Andrew, Obituary of František Josef Turček, 441
- Kiff, Lloyd F., The nest and eggs of the Black-and-yellow Silky-Flycatcher (*Phainoptila melanoxantha*), 198–199; Bird egg collections in North America, 746–755; review by, 823–825
- Kilham, Lawrence, Courtship and the pair-bond of Pileated Woodpeckers, 587–594
- Killdeer, 604–606

- King, Ben, reviews by, 216–217, 438–440, 644–645
- King, James R., review by, 428–430
- Kingbird, Eastern, 499–505; Tropical, 221–233
- Kinglet, Ruby-crowned, 499–505
- Kite, Gray-headed, 345
- Kittiwake, Black-legged, 537–543
- Kleptoparasitism, 806–807
- Knudtson, Eric P., see Day, Robert H., et al.
- Kosinski, Robert J., and Richard H. Podolsky, An analysis of breeding and mortality in a maturing kittiwake colony, 537–543
- Kramer, Howard G., and Nicholas S. Thompson, Geographic variation in the bell calls of the Blue Jay, 423–425
- Krebs, J. R., and N. B. Davies (Eds.), Behavioural ecology: an evolutionary approach (rev.), 819–821
- Kroll, James C., see Conner, Richard N., and ———
- Kroodsma, Donald E., Vocal dueling among male Marsh Wrens: evidence for ritualized expressions of dominance/subordinance, 507–515
- Lagopus lagopus*, 106–115, 313–316, 682–687
- L. leucurus*, 310–311, 316
- L. mutus*, 311–316
- Lake Superior, 1–5
- Lanius ludovicianus*, 159, 161–162, 798–800
- Lanyon, Scott M., see Thompson, Charles F., and ———
- Laphyctes satrapa*, 223, 230
- Lark, Horned, 392, 499–505, 623
- Larus argentatus*, 203, 364, 694–703, 809–812
- L. atricilla*, 203
- L. californicus*, 6–9
- L. delawarensis*, 1–5, 6–9, 284–295, 806–808
- L. heermanni*, 74
- L. occidentalis*, 74
- L. serranus*, 477, 480
- Leck, Charles F., and Frederic L. Cantor, Seasonality, clutch size, and hatching success in the Cedar Waxwing, 196–198; Avian extinctions in an isolated tropical wet-forest preserve, Ecuador, 343–352
- LeCroy, Mary, reviews by, 440, 644
- Lein, Ross M., review by, 426–428
- Lekagul, Boonsong, and Edward W. Cronin, Jr., Bird guide of Thailand (rev.), 439–440
- Lenington, Sarah, Predators and blackbirds: the “uncertainty principle” in field biology, 190–192
- Leopold, A. Starker, The California Quail (rev.), 436–437
- Leptasthenura andicola*, 477
- Leptodon cayanensis*, 345
- Leptopogon superciliaris*, 347
- Leptotila pallida*, 345
- Lessonia rufa*, 477, 480
- Leucosternis semiplumbea*, 345
- Lill, Alan, An assessment of male parental investment and pair bonding in the polygamous Superb Lyrebird, 489–498
- Limicola falcinellus*, 189–190
- Lincer, Jeffrey L., William S. Clark, and Maurice N. LeFranc, Jr., Working bibliography of the Bald Eagle (rev.), 643–644
- Lipaagus unirufus*, 347, 350
- Lloyd, Hoyes, Memorial of, 402–406
- Longevity (see Mortality and survival)
- Longspur, Lapland, 131–142, 392
- Lophophorus impejanus*, 74
- Lophortyx californicus*, 158, 161–162, 166, 499–505
- L. gambelii*, 784
- Lophotriccus pileatus*, 347
- Low, Rosemary, Lories and Lorikeets (rev.), 641–642
- Lukasiewicz, Peter, see Butler, Ronald G., and ———
- Lunda cirrhata*, 145, 150, 792–794
- Lyrebird, Superb, 489–498
- Lysurus crassirostris*, 379
- MacLean, S. F., see Seastedt, T. R., and ———
- Mader, William J., Breeding behavior of a polyandrous trio of Harris' Hawks in southern Arizona, 776–788
- Magpie, Black-billed, 192–195, 499–505
- Maine, 202, 737–745, 809–812
- Majnep, Ian Saem, and Ralph Bulmer, Birds of my Kalam country (rev.), 440
- Malacoptila panamensis*, 346, 350
- Mallard, 655–656, 706
- Manacus manacus*, 347
- Manitoba, 392–401, 600–603
- Manning, T. H., Density and volume corrections of eggs of seven passerine birds, 207–211
- Marti, Carl D., and Janet G. Hogue, Selection of prey by size in Screech Owls, 319–327
- Martin, Brown-chested, 607
- Massachusetts, 423–425
- Mating systems of *Aechmophorus occidentalis*, 573–586; *Agelaius phoeniceus*, 353–363; *Anumbius annumbi*, 607; *Calcarius lapponicus*, 133; *Dendroica castanea*, 600–601; *Furnarius rufus*, 607; *Junco hyemalis*, 630–631; *Larus argentatus*, 694–703; *L. californicus*, 6–9; *L. delawarensis*, 1–5, 6–9; *Menura superba*, 489–498; *Mimus saturninus*, 607–608; *Parabuteo unicinctus*, 776–788; *Passerculus sandwichensis*, 391–401; *Phaeoptilus tapera*, 607; *Polioptila dumicola*, 608; *Sialia currucoides*, 613–615
- Mayfield, Harold F., Commentary: The amateur in ornithology, 168–171

- Mayr, Ernst, and G. William Cottrell (Eds.), Check-list of birds of the world, Vol. 1, 2nd Ed. (rev.), 821–822
- McLandress, Ilse, see McLandress, M. Robert, and —
- McLandress, M. Robert, and Ilse McLandress, Blue-phase Ross' Geese and other blue-phase geese in western North America, 544–550
- Meadowlark, Eastern, 622–623; Western, 499–505
- Measurements (see Weights and —)
- Megadytes antipodes*, 103
- Megarynchus pitangua*, 184
- Melanerpes erythrocephalus*, 206–207
- M. pucherani*, 346
- M. uropygialis*, 158
- Melospiza melodia*, 499–505
- Memorials (see also Obituaries), Lloyd, Hoyes, 402–406; Saunders, Aretas Andrews, 173–178
- Menura superba*, 489–498
- Metriopelia aymara*, 477, 480
- M. ceciliae*, 477, 479
- M. melanoptera*, 477, 480
- Mexico, 183–186, 200–201, 221–223
- Meyer, John R., see Weise, Charles M., and —
- Meyer de Schauensee, Rodolphe, and William H. Phelps, Jr., A guide to the birds of Venezuela (rev.), 213–214
- Michigan, 196, 533, 562–572, 595–596, 630–631
- Micrastur ruficollis*, 345, 349
- Microbates cinereiventris*, 347
- Microcerculus marginatus*, 347
- Microrhopias quixensis*, 347
- Migration and orientation, commentary on, 813–815
- Migration and orientation of *Anas flavirostris*, 480; *A. georgica*, 480; *A. versicolor*, 480; *Chen caerulescens*, 548; *C. rossii*, 548; *Chloephaga melanoptera*, 480; *Fulica americana*, 480; hybrids of *Chen caerulescens* and *C. rossii*, 548; *Larus serranus*, 480; *Lessonia rufa*, 480; *Muscisaxicola alpina*, 480; *Plegadis ridgwayi*, 480; *Podiceps rolland*, 480; *Vanellus resplendens*, 480; *Zonotrichia capensis*, 480
- Miller, Alden H. (Ed.), Joseph Grinnell's philosophy of nature: selected writings of a western naturalist (rev.), 647
- Miller, Don E., and Michael R. Conover, Differential effects of chick vocalizations and bill-pecking on parental behavior in the Ring-billed Gull, 284–295; see Conover, Michael R., et al.
- Mimus longicaudatus*, 723–727, 733
- M. polyglottos*, 159, 161–162, 421–422
- M. saturninus*, 607–608
- Minnesota, 196
- Mionectes olivaceus*, 347, 349, 378
- Mississippi, 116–130, 604–606
- Mitrospingus cassinii*, 348
- Mniotilla varia*, 378, 383
- Mock, Douglas, review by, 827–828
- Mockingbird, 159, 421–422; Mockingbird, Galápagos, 723–736
- Mokhiber, Carmine, see Olson, Storrs L., et al.
- Molothrus ater*, 209–210, 483–488, 499–505, 600–602
- M. bonariensis*, 351, 774
- Molts and plumages of *Aechmophorus occidentalis*, 573–586; *Agelaius phoeniceus*, 356; *Branta canadensis*, 245–247; *Cyanocorax dickeyi*, 200–201; *C. morio*, 200–201; *C. sanblasiana*, 200–201; *Chen caerulescens*, 68–72; *Fringilla coelebs*, 91, 253–264; *Fulica gigantea*, 479; *Hemimacronyx cinnamomeippectus*, 444–445; *Hesperiphona vespertina*, 90–91; *Junco hyemalis*, 78–88, 94–99; *Lagopus lagopus*, 106–115, 682–687; *Melanerpes erythrocephalus*, 206–207; *Oxyura* sp., 479; *Saltator cinctus*, 610; *Zonotrichia albicollis*, 91; *Z. querula*, 88–90
- Momotus momota*, 346, 350
- Montana, 410
- Morrison, R. I. G., see Elliot, R. D., and —
- Mortality and survival of *Calcarius lapponicus*, 135; *Calidris pusilla*, 65; *Chen caerulescens*, 548; *C. rossii*, 548; *Florida caerulea*, 118–120; *Gymnorhinus cyanocephalus*, 797; hybrids of *Chen caerulescens* and *C. rossii*, 548; *Junco hyemalis*, 720; *Nesomimus parvulus*, 732; *Otus asio*, 795–796; *Parus atricapillus*, 52; *Rissa tridactyla*, 537–543; *Sialia sialis*, 570–571
- Moseley, Lynn J., Individual auditory recognition in the Least Tern (*Sterna albifrons*), 31–39
- Mosquin, Theodore, and Cecile Suchal (Eds.), Canada's threatened species and habitats (rev.), 433–434
- Moynihan, M., Geographic variation in social behavior and in adaptations to competition among Andean birds (rev.), 825–827
- Mueller, Helmut C., review by, 640
- Murton, R. K., and N. J. Westwood, Avian breeding cycles (rev.), 428–430
- Muscisaxicola alpina*, 477, 480
- M. rufivertex*, 477
- Myadestes melanops*, 378
- Myers, J. P., P. G. Connors, and F. A. Pitelka, Territory size in wintering Sanderlings: the effects of prey abundance and intruder density, 551–561
- Myiarchus cinerascens*, 157–158, 161
- Myioibius atricaudus*, 347
- M. barbatus*, 347
- Myiochorus miniatus*, 378, 380, 382–387
- Myiodynastes maculatus*, 347
- Myiophobus fasciatus*, 347

- Myotriccus ornatus*, 347  
*Myiozetetes inornata*, see *Conopias inornata*  
*M. similis*, 184  
*Myrmeciza exsul*, 347  
*M. immaculata*, 347  
*Myrmotherula axillaris*, 347  
*M. fulviventris*, 347  
*M. schisticolor*, 347, 379  
*M. surinamensis*, 347
- Nectarinia famosa*, 449–450, 453, 455, 457  
*N. kilimensis*, 449–450, 455  
*N. reichenowi*, 448–461  
*N. venusta*, 449–450, 453, 455
- Nelson, J. Bryan, Commentary: R. W. Schreiber's review of *The Sulidae: gannets and boobies*, 649; The Sulidae: gannets and boobies (rev.), 634–637
- Neocrex erythrops*, 425  
*Nesomimus parvulus*, 723–736
- Nest and nest site of *Agelaius phoeniceus*, 789; *Aquila chrysaetos*, 669; *Buteo lagopus*, 669; *Calidris pusilla*, 57–59, 61; *Cerorhinca monocerata*, 145; *Charadrius vociferus*, 604; *Conopias inornata*, 769–770; *Coturnicops noveboracensis*, 422–423; *Dendroica castanea*, 600–602; *Falco peregrinus*, 662–674; *F. rusticolus*, 669; *Haliaeetus leucocephalus*, 413; *Lunda cirrhata*, 145, 792–794; *Menura superba*, 489; *Metriopelia aymara*, 480; *M. melanoptera*, 480; *Mimus polyglottos*, 421–422; *Nesomimus parvulus*, 733; *Nycticorax nycticorax*, 364; *Oceanodroma leuconota*, 625–626; *Pandion haliaetus*, 413, 415; *Parabuteo unicinctus*, 776; *Parus atricapillus*, 42; *Passerculus sandwichensis*, 392, 394; *Phainopepla melanocephala*, 198–199; *Phoenicoparrus jamesi*, 332; *Pooecetes gramineus*, 803–804; *Ptychoramphus aleuticus*, 143–151; *Pygoscelis adeliae*, 680–681; *P. antarctica*, 680–681; *Quiscalus mexicanus*, 789; *Spizella arborea*, 392, 395; *Steatornis caripensis*, 188
- Nest parasitism, 7–9, 601–602, 774
- Nestling of *Aechmophorus occidentalis*, 581–582; *Aix sponsa*, 296–305; *Certhidea olivacea*, 800–802; *Conopias inornata*, 771–773; *Coturnix coturnix*, 10–30, 300–303; *Florida caerulea*, 116–130; *Geospiza conirostris*, 800–802; *G. difficilis*, 800–802; *G. fortis*, 800–802; *G. fuliginosa*, 800–802; *G. magnirostris*, 800–802; *G. scandens*, 800–802; *Gymnorhinus cyanocephalus*, 797; *Larus delawarensis*, 284–295; *Pandion haliaetus*, 415–417; *Passerculus sandwichensis*, 395–399; *Sterna hirundo*, 10–30; *Sturnus vulgaris*, 10–30
- Nevada, 574–575, 580  
New Brunswick, 625–627  
New Hampshire, 265–283
- New Jersey, 204, 694–703, 806–808  
New York, 10–30, 196, 507  
Newfoundland, 625–627, 790–792  
Nightingale-Thrush, Black-headed, 378; Ruddy-capped, 379; Slaty-backed, 378  
Nightjar, Jungle, 189; Swallow-tailed, 477, 480  
Nolan, Val, Jr., The ecology and behavior of the Prairie Warbler *Dendroica discolor* (rev.), 426–428; see Ketterson, Ellen D., and —
- North Carolina, 31–39, 202–203, 688–693  
North Dakota, 196  
Northwest Territories, 68–72, 662–674  
Norway, 411  
*Notharchus macrorhynchus*, 346, 350  
*N. tectus*, 346, 350  
*Nothropocta ornata*, 476  
*N. pentlandii*, 476  
*Nothura*, Spotted, 476, 479  
*Nothura maculosa*, 476, 479  
*Notiochelidon murina*, 477  
Nova Scotia, 410, 413–414  
*Numida meleagris*, 74  
*Nyctanassa violacea*, 725  
*Nyctea scandiaca*, 135  
*Nyctibius griseus*, 346, 349  
*Nycticorax nycticorax*, 364–369, 476  
*Nystalus radiatus*, 346, 348
- Oceanodroma leuconota*, 202, 625–627  
*Ochthoeca oenanthoides*, 477  
*Ocreatus underwoodii*, 346, 350  
*Odontophorus erythrops*, 345  
Ohio, 196, 533, 795–796  
Oilbird, 187–189  
Olson, Storrs L., Camm C. Swift, and Carmine Mokhiber, An attempt to determine the prey of the Great Auk (*Pinguinus impennis*), 790–792
- O'Neill, John P., review by, 213–214; and Thomas S. Schulenberg, Notes on the Masked Saltator, *Saltator cinctus*, in Peru, 610–613; see Fitzpatrick, John W., and —
- Ontario, 196, 207–211, 422–423, 483–488, 596–599
- Oporornis formosus*, 379–380, 383  
Oregon, 410, 527–531, 574–575, 577  
*Oreopholus ruficollis*, 476  
*Oreoscoptes montanus*, 799  
*Oreotrochilus estella*, 477  
Oring, Lewis W., review by, 639–640  
Oriole, Hooded, 159; Northern, 499–505  
*Ornithion semiflavum*, 347  
*Ornithion semiflavum*, 347  
*O. vetula*, 184  
Ortiz-Crespo, Fernando I., Discovery of an Oilbird colony in the western drainage of the Ecuadorian Andes, 187–189

- Osborne, David R., and Steven R. Beissinger, The Paint-billed Crake in Guyana, 425
- Osprey, 408–412, 413–414, 415–417
- Oster, George F., and Edward O. Wilson, Caste and ecology in the social insects (rev.), 633–634
- Otus asio*, 319–327, 795–796
- O. guatemalae*, 345
- O. scops*, 189
- Owl, Barn, 204, 477, 480, 619–621; Great Horned, 202; Scops, 189; Screech, 319–327, 795–796; Short-eared, 725; Snowy, 135; Spotted, 525–531
- Oxyura jamaicensis*, 476, 479
- O. sp.*, 476, 479
- Pachyramphus albogriseus*, 347
- P. cinnamomeus*, 347
- P. versicolor*, 379
- Packard, Gary C., see Sutherland, Paul R., et al.
- Panama, 759–760
- Pandion haliaetus*, 408–412, 413–414, 415–417
- Parabuteo unicinctus*, 413, 776–788
- Parakeet, Andean, 477, 480
- Parasitism (see Nest parasitism, see also Kleptoparasitism)
- Parkes, Kenneth C., see Short, Lester L., and
- 
- Partridge, Grey, 499–505
- Parula pitiayumi*, 347
- Parus atricapillus*, 40–55
- Pasquier, Roger F., review by, 437–438
- Passerculus sandwichensis*, 391–401, 623–624
- Passerina ciris*, 417–418
- Patagona gigas*, 477
- Pavo muticus*, 74
- Payne, Robert B., review by, 436; Two apparent hybrid *Zonotrichia* sparrows, 595–599
- Pedioecetes phasianellus*, 308, 316
- Pelecanus occidentalis*, 74, 407–408
- P. onocrotalus*, 407–408
- Pelican, Brown, 407–408; Great White, 407–408
- Penelope purpurascens*, 345, 348–349
- Penguin, Adelie, 101, 675–681; Chinstrap, 675–681; Gentoo, 101, 675–676; Jackass, 101; Macaroni, 101; Rockhopper, 100–105; Yellow-eyed, 103
- Pennsylvania, 10–30, 196
- Perdix perdix*, 499–505
- Peru, 328–342, 443–447, 475–482, 610–613
- Pesticides (see Pollutants)
- Petrochelidon andecola*, 477
- Phaeoptilus tapera*, 607
- Phaethon rubricauda*, 74
- Phaethornis longuemareus*, 346
- P. superciliosus*, 346
- P. yaruqui*, 346
- Phainopepla*, 160
- Phainopepla nitens*, 160, 199
- Phainoptila melanoxantha*, 198–199
- Phalacrocorax auritus*, 204–206, 364, 476
- P. olivaceus*, 204–206
- Phalcobaeus megalopterus*, 476
- Phasianus colchicus*, 74, 499–505
- Pheasant, Ring-necked, 499–505
- Pheucticus melanocephalus*, 160
- Phillips, W. W. A., Annotated checklist of the birds of Ceylon (Sri Lanka) (rev.), 644–645
- Philomachus pugnax*, 189
- Philydor erythrocercus*, 346
- Phleocryptes melanops*, 477
- Phloeoeastes guayaquilensis*, 346
- P. melanoleucus*, 346, 348
- Phoebe, Eastern, 209–210
- Phoenicoparrus andinus*, 328–342
- P. jamesi*, 328–342
- Phoenicopterus chilensis*, 328–342, 476, 481
- P. ruber*, 615–619
- Phrygilus alaudinus*, 478
- P. fruticeti*, 478
- P. gayi*, 478
- P. plebejus*, 478, 480
- Physiology, body composition, 296–303; egg-laying, 248–250; energy cost of activities, 58–72, 234–252, 253–264; growth, 10–30, 296–305, 742–744; hormones, 106–115, 686; incubation, 234–252, 625–627; metabolism, 234–252; molt, 68–72, 237–247, 253–264, 682–687; nutrient reserves, 68–72, 234–252, 296–305; plumage development, 106–115; thermal regulation, 10–30, 121–123, 253–264, 604–606; water balance, 298–303; water loss from eggs, 192–195
- Physiology of *Aix sponsa*, 296–305; *Anas rubripes*, 742–744; *Bonasa umbellus*, 308–309, 316; *Branta canadensis*, 234–252; *Canachites canadensis*, 309–310, 316; *Chen caerulescens*, 68–72; *Coturnix coturnix*, 10–30, 300–303; *Dendragapus obscurus*, 308, 316; *Florida caerulea*, 121–123; *Fringilla coelebs*, 253–264; *Lagopus lagopus*, 112–114, 313–316, 682–687; *L. leucurus*, 310–311; *L. mutus*, 311–316; *Pedioecetes phasianellus*, 308, 316; *Pelecanus onocrotalus*, 407–408; *Sterna hirundo*, 10–30; *Sturnus vulgaris*, 10–30
- Piaya cayana*, 345
- Pica pica*, 192–195, 499–505
- Picoides nuttallii*, 418–420
- P. scalaris*, 157–158, 161–162
- Piculus leucolaemus*, 346
- P. rubiginosus*, 346
- P. simplex*, 378
- Picumnus olivaceus*, 346
- Pinguinus impennis*, 790–792
- Pinkowski, Benedict C., Annual productivity and

- its measurement in a multi-brooded passerine, the Eastern Bluebird, 562–572
- Pionopsitta haematocephala*, 345
- Pionus chalcopterus*, 345
- P. menstruus*, 345
- P. sordidus*, 345, 350
- Pipilo chlorurus*, 160
- P. erythrourhynchus*, 688–693
- P. fuscus*, 159
- Pipra coronata*, 347, 350
- P. mentalis*, 347
- Pipromorpha oleaginea*, 347
- Piranga ludoviciana*, 499–505
- P. rubra*, 379
- Pitangus sulphuratus*, 184
- Pitelka, F. A., see Myers, J. P., et al.
- Pittasoma rufopileatum*, 346, 350
- Pitylus fuliginosus*, 612
- P. grossus*, 348, 612
- Platycichla leucopsis*, 347
- Platyrinchus coronatus*, 347, 349
- P. mystaceus*, 347, 378, 380
- Plegadis ridgwayi*, 476, 479, 480–481
- Plover, Mongolian, 189
- Plumages (see Molts and ——)
- Podiceps occipitalis*, 476
- P. rolland*, 476, 478, 480–481
- Podilymbus podiceps*, 706
- Podolsky, Richard H., see Kosinski, Robert J., and ——
- Poecilotriccus ruficeps*, 347, 350
- Pogonotriccus ophthalmicus*, 347, 350
- Polioptila caerulea*, 157–158, 161
- P. dumicola*, 608
- P. melanura*, 157–158, 161
- P. schistaceigula*, 347
- Pollutants in *Larus argentatus*, 810–812
- Polyandry (see Mating systems)
- Polygamy (see Mating systems)
- Polygyny (see Mating systems)
- Pooecetes gramineus*, 499–505, 622–624, 802–805
- Poole, Alan, Sibling aggression among nestling Ospreys in Florida Bay, 415–417
- Popelairia conversii*, 346
- Powell, George V. N., Structure and dynamics of interspecific flocks in a neotropical mid-elevation forest, 375–390
- Power, Harry W., Is displacement a sign of female dominance or only a response to close following by males trying to avoid being cuckolded?, 613–615
- Predation by *Agriornis montana*, 480; *Aquila chrysaetos*, 548; *Asio flammeus*, 725; *Bubo virginianus*, 202; *Buteo poecilochrous*, 479; *Circus cyaneus*, 370–374; *Corvus brachyrhynchos*, 367, 804; *Haliaeetus leucocephalus*, 413–414, 548; *Lanius ludovicianus*, 798–799; *Larus argentatus*, 203; *L. atricilla*, 203; *Nesomimus parvulus*, 726, 729; *Nyctanassa violacea*, 725; *Nyctea scandiaca*, 135; *Otus asio*, 319–327; *Pandion haliaetus*, 413–414; *Parabuteo unicinctus*, 784; *Pinguinus impennis*, 790–792; *Stercorarius parasiticus*, 135; *S. pomarinus*, 135; *Tyto alba*, 620
- Predation on *Agelaius phoeniceus*, 190–191; *Calocitta lapponica*, 135; *Certhidea olivacea*, 726, 729; *Chen caerulescens*, 548; *C. rossii*, 548; *Gelochelidon nilotica*, 203; *Geococcyx californianus*, 784; *Geospiza difficilis*, 726, 729; *Harpia harpyja*, 349; hybrids of *Chen caerulescens* and *C. rossii*, 548; *Lophortyx gambelii*, 784; *Metriopelia ceciliae*, 479; *Momotus momota*, 350; *Nesomimus parvulus*, 725; *Nothura maculosa*, 479; *Nycticorax nycticorax*, 367; *Oceanodroma leucorhoa*, 202; *Penelope purpurascens*, 349; *Phoenicopterus ruber*, 617–618; *Tinamus major*, 349; *Toxostoma curvirostre*, 784
- Premnoplex brunnescens*, 346, 349, 378, 380–387
- Prévost, Yves, Osprey-Bald Eagle interactions at a common foraging site, 413–414
- Psilorhinus morio*, 184
- Ptarhigan, Rock, 311–316; White-tailed, 310–311, 316; Willow, 106–115, 313–316, 682–687
- Pteroglossus sanguineus*, 346
- P. torquatus*, 184
- Pterocnemia pennata*, 74
- Ptilogonyx caudatus*, 199
- P. cinereus*, 199
- Ptychoramphus aleuticus*, 143–151
- Puerto Rico, 204–206
- Puffin, Tufted, 145, 792–794
- Pulsatrix perspicillata*, 345, 348
- Pygoscelis adeliae*, 101, 675–681
- P. antarctica*, 675–681
- P. papua*, 101, 675–676
- Pyriglena leuconota*, 347
- Pyrrhuloxia, 159
- Quail, California, 158, 499–505; Gammel's, 784; Japanese, 10–30, 300–303
- Quebec, 196, 364–369
- Quelea, Red-billed, 516–524
- Quelea quelea*, 516–524
- Querula purpurata*, 347
- Quiscalus mexicanus*, 789
- Q. quiscula*, 209–210, 608–609
- Raffaele, Herbert A., Status of the genus *Phalacrocorax* in Puerto Rico including the first records of *P. auritus*, 204–206
- Rahn, Hermann, see Ricklefs, Robert E., and ——
- Rail, Yellow, 422–423
- Rallus sanguinolentus*, 476
- Ramphastos ambiguus*, 346, 348

- R. brevis*, 346, 348  
*R. sulfuratus*, 184  
*Ramphocænus melanurus*, 347  
 Ratti, John T., Reproductive separation and isolating mechanisms between sympatric dark- and light-phase Western Grebes, 573–586  
 Raveling, Dennis G., The annual cycle of body composition of Canada Geese with special reference to control of reproduction, 234–242  
 Raven, Common, 159; White-necked, 782  
 Rawal, U. M., Certain aspects of the functional anatomy of the feeding apparatus of some pectoral groups of birds (rev.), 647  
 Redhead, 706  
 Redpoll, Common, 392; Hoary, 392  
 Redstart, American, 265–283; Slate-throated, 378, 380, 382–387  
 Rees, William C., see Roe, Nicholas A., and —  
     *Regulus calendula*, 499–505  
 Reinecke, Kenneth J., Feeding ecology and development of juvenile Black Ducks in Maine, 737–745  
 Reynolds, Timothy D., The impact of Loggerhead Shrikes on nesting birds in a sagebrush environment, 798–800  
*Rhynchoscyclus brevirostris*, 347, 378  
*R. fulvipectus*, 347, 350  
 Richards, Douglas G., Recognition of neighbors by associative learning in Rufous-sided Towhees, 688–693  
 Rickard, W. H., see Rotenberry, J. T., et al.  
 Ricklefs, Robert E., Patterns of growth in birds. V. A comparative study of development in the Starling, Common Tern, and Japanese Quail, 10–30; and Hermann Rahn, The incubation temperature of Leach's Storm-Petrel, 625–627  
*Rissa tridactyla*, 74, 537–543  
 Roadrunner, 158, 784  
 Robin, American, 208–211, 499–505  
 Roe, Nicholas A., and William E. Rees, Notes on the puna avifauna of Azángaro Province, Department of Puno, southern Peru, 475–482  
 Rogers, David T., Jr., Jerome A. Jackson, Bette J. Schardien, and Martha S. Rogers, Observations at a nest of a partial albino Red-headed Woodpecker, 206–207  
 Rogers, Martha S., see Rogers, David T., Jr., et al.  
 Romesburg, H. Charles, see Balph, Martha Hatch, et al.  
 Rotenberry, J. T., R. E. Fitzner, and W. H. Rickard, Seasonal variation in avian community structure: differences in mechanisms regulating diversity, 499–505  
 Roth, Roland R., Foraging behavior of Mockingbirds: the effect of too much grass, 421–422  
 Ruff, 189  
 Rutledge, James T., and Randall S. Chandler, Nest-site competition between Red-winged Blackbirds and Great-tailed Grackles, 789  
 Ryan, Mark R., and James J. Dinsmore, A quantitative study of the behavior of breeding American Coots, 704–713  
 Ryder, John P., and Patricia Lynn Somppi, Female-female pairing in Ring-billed Gulls, 1–5  
 Safriel, Uriel N., see Ashkenazie, Shoshana, and —  
     *Salpinctes obsoletus*, 499–505  
 Saltator, Masked, 610–613  
*Saltator albicollis*, 612  
*S. atriceps*, 184, 612  
*S. atricollis*, 612  
*S. atripectoralis*, 612  
*S. cinctus*, 610–613  
*S. coeruleopectoralis*, 612  
*S. maximus*, 612  
*S. nigriceps*, 612  
*S. orenocensis*, 612  
*S. rufiventris*, 612  
*S. similis*, 612  
 Sanderling, 190, 551–561  
 Sandpiper, Broad-billed, 189–190; Semipalmated, 56–57; Spoon-bill, 180  
 Sanger, Gerald A., see Gill, Robert, Jr., and —  
     Sapsucker, Yellow-bellied, 195  
 Saskatchewan, 544–550  
 Saunders, Aretas Andrews, Memorial of, 173–178  
*Sayornis phoebe*, 209–210  
 Schardien, Bette J., and Jerome A. Jackson, Belly-soaking as a thermoregulatory mechanism in nesting Killdeers, 604–606; see Rogers, David T., Jr., et al.  
*Schiffornis turdinus*, 347, 349  
 Schluter, D., see Grant, P. R., et al.  
 Schreiber, Ralph, review by, 634–637  
 Schulenberg, Thomas S., see O'Neill, John P., and —  
     Schulmeister, Robert P., see Day, Robert H., et al.  
*Sclerurus guatemalensis*, 346  
*S. mexicanus*, 346, 349  
 Scott, D. M., and C. Davison Ankney, Evaluation of a method for estimating the laying rate of Brown-headed Cowbirds, 483–488  
 Scythebill, Brown-billed, 379–380  
 Sealy, Spencer G., Extralimital nesting of Bay-breasted Warblers: response to forest tent caterpillars?, 600–603  
 Searcy, William A., Male characteristics and pairing success in Red-winged Blackbirds, 353–363  
 Sears, Harold F., Colonial nesting as an anti-pred-

- ator adaptation in the Gull-billed Tern, 202–203
- Seastedt, T. R., and S. F. MacLean, Territory size and composition in relation to resource abundance in Lapland Longspurs breeding in arctic Alaska, 131–142
- Semnornis frantzii*, 378
- Serle, W., and G. J. Morel, A field guide to the birds of West Africa (rev.), 436
- Setophaga ruticilla*, 265–283
- Severinghaus, Sheldon R., and Kenneth T. Blackshaw, A new guide to the birds of Taiwan (rev.), 216
- Sherry, Thomas W., Competitive interactions and adaptive strategies of American Redstarts and Least Flycatchers in a northern hardwoods forest, 265–283
- Short, Lester L., and Kenneth C. Parkes, The status of *Agelaius forbesi* Sclater, 179–183; and Jennifer F. M. Horne, review by, 634
- Shrike, Loggerhead, 798–800
- Shrike-Tyrant, Black-billed, 477, 480
- Sialia currucooides*, 613–615
- S. sialis*, 562–572
- Sicalis uropygialis*, 478, 480
- Sierra-Finch, Ash-breasted, 478, 480
- Silky-Flycatcher, Black-and-yellow, 198–199
- Silver, Rae (Ed.), Parental behavior in birds (rev.), 215–216
- Sirystes sibilator*, 347, 350
- Sittasomus griseicapillus*, 378, 380–387
- Skead, C. J., The sunbirds of South Africa. Also the sugarbirds, the white-eyes and the spotted creeper (rev.), 647
- Snow, D. W. (Ed.), An atlas of speciation in African nonpasserine birds (rev.), 638–639
- Solitaire, Black-faced, 378
- van Someren, V. G. L., The birds of East Africa. Vol. 1. *Ploceidae-1* (rev.), 634
- Somppi, Patricia Lynn, see Ryder, John P., and
- 
- Song (see Vocalizations)
- Sonograms of *Carpodacus mexicanus*, 464, 466–467; *Cistothorus palustris*, 508; *Conopias ornata*, 769; *Cyanocitta cristata*, 424; *Pipilo erythrrophthalmus*, 689; *Sturna albifrons*, 33, 36; *Thryothorus ludovicianus*, 689
- Sotherland, Paul R., Gary C. Packard, and Theodore L. Taigen, Permeability of magpie and blackbird eggshells to water vapor: variation among and within nests of a single population, 192–195
- Soucy, Leonard J., Jr., A nearly synchronous hatching of Barn Owls, 204
- South Africa, 100–105
- South Carolina, 533
- Sparrow, Black-throated, 158, 162–163; Brewer's, 160, 799; Field, 623; Golden-crowned, 595–597; Grasshopper, 621–625; Harris', 80–90, 596–599; Henslow's, 623; Lark, 160; Sage, 499–505, 799; Savannah, 391–401, 623–624; Song, 499–505; Tree, 392; Vesper, 499–505, 622–624, 802–805; White-crowned, 160, 392, 499–505, 596–599; White-throated, 91, 595–597
- Speotyto (Athene) cunicularia*, 477
- Spheniscus demersus*, 101
- Sphyrapicus varius*, 195
- Spinetail, Red-faced, 378, 380–386
- Spinus atratus*, 478
- S. crassirostris*, 478
- Spizella arborea*, 392, 395
- S. breweri*, 160, 799
- S. pusilla*, 623
- Starling, 10–30, 209–210, 407–408, 499–505, 806–807
- Steatornis caripensis*, 187–189
- Steele, John, The researcher's field notebook (rev.), 647
- Stegmann, Boris C., Relationships of the superorders Alectromorphae and Charadriomorphae (Aves): a comparative study of the avian hand (rev.), 637
- Stercorarius parasiticus*, 135
- S. pomarinus*, 135
- Sterling, Keir B. (Ed.), Biologists and their world (rev.), 637–638; review by, 831–832
- Sterna albifrons*, 31–39
- S. hirundo*, 10–30
- Stokkan, Karl-Arne, Testosterone and daylength-dependent development of comb size and breeding plumage of male Willow Ptarmigan (*Lagopus lagopus*), 106–115; The effect of permanent short days and castration on plumage and comb growth in male Willow Ptarmigan (*Lagopus lagopus*), 682–687
- Stonehouse, Bernard, and Christopher Perrins (Eds.), Evolutionary ecology (rev.), 434–435
- Storm-Petrel, Leach's, 202, 625–627
- Strix occidentalis*, 525–531
- Struthio camelus*, 74
- Sturnella magna*, 622–623
- S. neglecta*, 499–505
- Sturnus vulgaris*, 10–30, 209–210, 407–408, 499–505, 806–807
- Sula leucogaster*, 74
- Summers, Kenneth R., see Vermeer, Kees, et al.
- Sunbird, Bronzy, 449–450, 455; Golden-winged, 448–461; Malachite, 449–450, 453, 455, 457; Variable, 449–450, 453, 455
- Sveriges Ornitologiska Förening (SOF), Sveriges fåglar (rev.), 646–647
- Swallow, Barn, 208–211, 499–505
- Swenson, Jon E., The relationship between prey species ecology and dive success in Ospreys, 408–412

- Swift, Camm C., see Olson, Storrs L., et al.
- Syndactyla subalaris*, 378, 380–387
- Syrmaticus soemmerringii*, 74
- Tachuris rubrigastra*, 477
- Tachyphonus delatrii*, 348
- T. luctuosus*, 348
- Taigen, Theodore L., see Sotherland, Paul R., et al.
- Tanager, Silver-throated, 378; Spangle-cheeked, 379; Summer, 379; Western, 499–505
- Tangara arthus*, 809
- T. chilensis*, 809
- T. cyanocephala*, 809
- T. dowii*, 379
- T. fastuosa*, 809
- T. gyrola*, 348
- T. icterocephala*, 348, 350, 378
- T. lavinia*, 348, 350
- T. nigroviridis*, 348, 350
- T. parzudakii*, 809
- T. ruficervix*, 809
- T. xanthocephala*, 809
- Tanner, James T., Guide to the study of animal populations (rev.), 440
- Taraba major*, 347
- Taxonomy of *Agelaius forbesi*, 179–183; *Chlorochrysa calliparaea*, 808–809; *C. nitidissima*, 809; *C. phoenicotis*, 809; *Conopias inornata*, 774; *Curaeus forbesi*, 179–183; **Hemitriccus cinnamomeipectus**, 443, 445–446; *H. kaempferi*, 445–446; *H. mirandae*, 445–446; *Laphyctes satrapa*, 223, 230; *Pitylus fuliginosus*, 612; *P. grossus*, 612; *Saltator albicollis*, 612; *S. atriceps*, 612; *S. atricollis*, 612; *S. atripennis*, 612; *S. aurantirostris*, 612; *S. cinctus*, 612–613; *S. coerulescens*, 612; *S. maximus*, 612; *S. nigriceps*, 612; *S. orenocensis*, 612; *S. rufiventris*, 612; *S. similis*, 612; *Tangara arthus*, 809; *T. chilensis*, 809; *T. cyanocephala*, 809; *T. fastuosa*, 809; *T. parzudakii*, 809; *T. ruficervix*, 809; *T. xanthocephala*, 809; *Tyrannus chloronotus*, 221, 223, 230; *T. couchii*, 221–233; *T. melancholicus*, 221–233; *T. occidentalis*, 221, 223, 230
- Teal, Blue-winged, 655–656, 659; Speckled, 476, 479–480
- Temple, Stanley A., review by, 436–437
- Tennessee, 628–630
- Terenotriccus erythrurus*, 347
- Tern, Common, 10–30; Gull-billed, 202–203; Least, 31–39
- Territory in *Agelaius phoeniceus*, 353–363; *Ara ambiguia*, 348; *Aramides wolfi*, 348; *Atlapetes brunneinucha*, 382; *Basileuterus culicivorus*, 382; *B. tristriatus*, 382; *Calcarius lapponicus*, 131–142; *Calidris alba*, 551–561; *C. pusilla*, 63–64; *Chlorospingus ophthalmicus*, 382; *Cis tothorus palustris*, 513; *Conopias inornata*, 769–770; *Cranioleuca erythrops*, 382; *Empidonax minimus*, 267; *Fulica americana*, 710–711; *Harpia harpyja*, 348; *Henicorhina leucophrys*, 382; *Menura superba*, 494–496; *Myioborus miniatus*, 382; *Nectarinia reichenowi*, 448–461; *Nesomimus parvulus*, 733–734; *Parus atricapillus*, 40–55; *Passerculus sandwichensis*, 392; *Penelope purpurascens*, 348; *Phloeoceastes melanoleucus*, 348; *Premnoplex brunnescens*, 382; *Pulsatrix perspicillata*, 348; *Ramphastos ambiguus*, 348; *R. brevis*, 348; *Setophaga ruticilla*, 267; *Sittasomus griseicapillus*, 382; *Spizella arborea*, 395; *Syndactyla subalaris*, 382; *Tinamus major*, 348; *Xiphorhynchus erythropygius*, 378, 380–387
- Texas, 195, 221–223, 421–422, 789
- Thalurania furcata*, 346
- Thamnophilus punctatus*, 347, 351
- Theristicus caudatus*, 476
- Thinocorus orbignyianus*, 476
- Thomas, Betsy Trent, Behavior and breeding of the White-bearded Flycatcher (*Conopias inornata*), 767–775
- Thompson, Charles F., and Scott M. Lanyon, Reverse mounting in the Painted Bunting, 417–418
- Thompson, Nicholas S., see Kramer, Howard G., and ——
- Thrasher, Curve-billed, 784; Gray, 159; Sage, 799
- Thraupis episcopus*, 351
- Threnetes ruckeri*, 346
- Thripadectes rufobrunneus*, 379
- Thrush, Varied, 805–806
- Thryothorus ludovicianus*, 688–689
- Thurber, Walter A., Cien Aves de El Salvador (rev.), 645
- Tinamus major*, 345, 348–349
- Tityra semifasciata*, 184
- Todd, Frank S., Waterfowl: Ducks, geese and swans of the world (rev.), 830–831
- Todirostrum nigriceps*, 347
- Tody-tyrant, Cinnamon-breasted**, 443–447
- Tolmomyias assimilis*, 347
- Touit dilectissima*, 345, 350
- Towhee, Brown, 159; Green-tailed, 160; Rufous-sided, 688–693
- Toxostoma cinereum*, 159
- T. curvirostre*, 484
- Tramer, Elliot J., and Thomas R. Kemp, Diet-correlated variations in social behavior of wintering Tennessee Warblers, 186–187
- Taylor, Melvin A., Jr., Two sibling species of *Tyrannus* (Tyrannidae), 221–233; review by, 638–639; (Ed.), Check-list of birds of the world, Vol. 8 (rev.), 822–823
- Treehunter, Streak-breasted, 379
- Tremblay, Jacqueline, and Laurence N. Ellison,

- Effects of human disturbance on breeding of Black-crowned Night Herons, 364–369
- Tringa flavipes*, 476
- T. melanoleuca*, 476, 479
- Trivelpiece, Wayne, and Nicholas J. Volkman, Nest-site competition between Adelie and Chinstrap penguins: an ecological interpretation, 675–681
- Troglodytes musculus* (= *aedon*?), 477
- T. ochraceus*, 378
- Trogon citreolus*, 184
- T. collaris*, 346
- T. melanurus*, 346
- T. viridis*, 346
- Turček, Frantiček Josef, Obituary of, 441
- Turdus chiguanco*, 477
- T. grayi*, 84
- T. migratorius*, 74, 208–211, 499–505
- T. nudigenis*, 347
- T. obssoletus*, 347
- Turnstone, Ruddy, 189
- Tyranniscus nigrocipillus*, 347, 350
- T. viridiflavus*, 347, 350
- Tyrannulus elatus*, 347
- Tyrannus chloronotus*, 221, 223, 230
- T. couchii*, 221–233
- T. melancholicus*, 184, 221–233
- T. occidentalis*, 221, 224, 230
- T. tyrannus*, 499–505
- Tyto alba*, 204, 480, 619–621
- Upucerthia serrana*, 477
- U. validirostris*, 477
- Uropsalis segmentata*, 477, 480
- Utah, 78–88, 533, 573–586, 714
- VanCamp, Laurel F., see Henny, Charles J., and —
- Vanellus resplendens*, 476, 480
- Venezuela, 767–775
- Veniliornis kirkii*, 346
- Verdin, 158
- Vermeer, Kees, Rebecca A. Vermeer, Kenneth R. Summers, and Raymond R. Billings, Numbers and habitat selection of Cassin's Auklet breeding on Triangle Island, British Columbia, 143–151
- Vermeer, Rebecca A., see Vermeer, Kees, et al.
- Vermivora celata*, 160
- V. chrysoptera*, 379–380, 383
- V. peregrina*, 186–187
- Viney, Clive, and Karen Phillips, A colour guide to Hong Kong birds (rev.), 216–217
- Vireo olivaceus*, 347
- Virginia, 196
- Vocalization, geographic variation of, 423–425, 464–474
- Vocalizations of *Basileuterus tristriatus*, 385, 388; *Carpodacus mexicanus*, 464–474; *Cistothorus palustris*, 507–515; *Conopias inornata*, 769; *Cyanocitta cristata*, 423–425; *Dryocopus pileatus*, 590–591; hybrids of *Tyrannus couchii* and *T. melancholicus*, 266; *Larus delawarensis*, 284–295; *Pipilo erythrorththalmus*, 688–693; *Sterna albifrons*, 31–39; *Tyrannus couchii*, 226; *T. melancholicus*, 226
- Vogt, Richard C. Cleaning/feeding symbiosis between grackles (*Quiscalus*: Icteridae) and map turtles (*Graptemys*: Emydidae), 608–609
- Volkman, Nicholas J., see Trivelpiece, Wayne, and —
- Vuilleumier, François, review by, 825–827
- Vultur gryphus*, 476
- Walter, Hartmut, Eleonora's Falcon: adaptations to prey and habitat in a social raptor (rev.), 832–833
- Warbler, Bay-breasted, 600–603; Black-and-white, 379–383; Black-throated Green, 379, 383; Blackpoll, 392; Golden-crowned, 378, 380–386; Golden-winged, 379–380, 383; Kentucky, 379–380, 383; Kirtland's, 651–654, 655–658; Orange-crowned, 160; Tennessee, 186–187; Three-striped, 378, 380, 383–388; Wilson's, 379, 383, 499–505; Yellow, 159, 392; Yellow-rumped, 499–505
- Washington, 6–9, 284–295, 353–363, 499–505, 512, 533, 805–806
- Waxwing, Cedar, 196–198
- Weather, influence on (see Environment, influence on)
- Weatherhead, Patrick J., Ecological correlates of monogamy in tundra-breeding Savannah Sparrows, 391–401
- Weeden, Robert B., Relative heart size in Alaskan Tetraonidae, 306–318
- Weights and measurements of *Agelaius phoeniceus*, 208–211; *Aix sponsa*, 74, 298–303; *Amazonetta brasiliensis*, 74; *Anas platyrhynchos*, 74; *A. rubripes*, 742–743; *Anous stolidus*, 74; *A. tenuirostris*, 74; *Anser fabalis*, 74; *Bonasa umbellus*, 308–309, 316; *Branta canadensis*, 236–248; *Buteo jamaicensis*, 74; *Canachites canadensis*, 309–310, 316; *Chen caerulescens*, 69–70; *C. rossii*, 546; *Chrysophorus amherstiae*, 74; *C. pictus*, 74; *Conopias inornata*, 773; *Coturnix coturnix*, 13–27, 300–303; *Curaeus forbesi*, 182; *Dendragapus obscurus*, 308, 316; *Dromaius novaehollandiae*, 74; *Empidonax minimus*, 270; *Fringilla coelebs*, 253–264; *Fulica americana*, 74; *Gygis alba*, 74; *Hemimacrurus cinamomeipectus*, 444–445; *H. kaempferi*, 445; *H. mirandae*, 445; *Hirundo rustica*, 208–211; hybrids of *Chen caerulescens* and *C. rossii*, 546; hybrids of *Tyrannus couchii* and *T. melancholicus*, 225–229; *La-*

- gopus lagopus*, 313–316; *L. leucurus*, 310–311, 316; *L. mutus*, 311–316; *Larus argentatus*, 697–698, 810–811; *L. heermanni*, 74; *L. occidentalis*, 74; *Lophophorus impejanus*, 74; *Molothrus ater*, 209–210; *Neocrex erythrops*, 425; *Numida meleagris*, 74; *Oceanodroma leuconota*, 626–627; *Otus asio*, 795–796; *Pandion haliaetus*, 416; *Passerculus sandwichensis*, 395, 398–399; *Pavo muticus*, 74; *Pedioecetes phasianellus*, 308, 316; *Pelecanus occidentalis*, 74; *P. onocrotalus*, 407–408; *Phaethon rubricauda*, 74; *Phainoptila melanoxantha*, 198; *Phalacrocorax auritus*, 204; *Phasianus colchicus*, 74; *Pica pica*, 192–195; *Pitylus fuliginosus*, 612; *P. grossus*, 612; *Pteroicnemia pennata*, 74; *Quiscalus quiscula*, 209–210; *Rissa tridactyla*, 74; *Saltator albocollis*, 612; *S. atriceps*, 612; *S. atricollis*, 612; *S. atripennis*, 612; *S. aurantiirostris*, 612; *S. cinctus*, 611–612; *S. coerulescens*, 612; *S. maximus*, 612; *S. nigriceps*, 612; *S. orenocensis*, 612; *S. rufiventris*, 612; *S. similis*, 612; *Sayornis phoebe*, 209–210; *Setophaga ruticilla*, 270; *Sterna hirundo*, 13–27; *Struthio camelus*, 74; *Sturnus vulgaris*, 13–27; *Struthio camelus*, 74; *Turdus migratorius*, 74, 208–211; *Tyrannus couchii*, 222–229; *T. melancholicus*, 222–229; *Xanthocephalus xanthocephalus*, 192–195
- Weise, Charles M., and John R. Meyer, Juvenile dispersal and development of site-fidelity in the Black-capped Chickadee, 40–55
- Weller, Milton W., review by, 830–831
- Wershkul, David F., Nestling mortality and the adaptive significance of early locomotion in the Little Blue Heron, 116–130
- West Virginia, 621–625, 802–805
- Whaley, Wayne H., see Ellis, David H., and —
- Whitmore, Robert C., Short-term change in vegetation structure and its effect on Grasshopper Sparrows in West Virginia, 621–625; see Wray, Thomas, II, and —
- Wiens, John A., reviews by, 430–434, 440, 633–634, 643–644, 646–647, 833–834
- Wight, Howard M., see Forsman, Eric D., and —
- Wiley, Beth Nethery, see Wiley, James W., and —
- Wiley, James W., and Beth Nethery Wiley, Status of the American Flamingo in the Dominican Republic and Eastern Haiti, 615–619
- Williams, A. J., see Burger, A. E., and —
- Wilsonia pusilla*, 379, 383, 499–505
- Wisconsin, 40–55, 196, 370–374, 608–609
- Wolf, Larry L., see Gill, Frank B., and —
- Wood-Wren, Gray-breasted, 378, 380–386
- Woodcreeper, Olivaceous, 378, 380–387; Spotted, 378, 380–387; Wedge-billed, 378
- Woodpecker, Gila, 158; Ladderbacked, 158, 162; Nuttall, 418–420; Pileated, 587–594; Red-headed, 206–207; Rufous-winged, 378
- Wolfenden, Glen E., review by, 215–216
- Woolington, Dennis W., see Day, Robert H., et al.
- Wray, Thomas, II, and Robert C. Whitmore, Effects of vegetation on nesting success of Vesper Sparrows, 802–805
- Wren, Cactus, 159; Canyon, 159; Carolina, 688–689; Marsh, 507–515; Ochraceous, 378
- Wyoming, 410
- Xanthocephalus xanthocephalus*, 192–195
- Xenops minutus*, 346
- X. rutilans*, 346
- Xiphorhynchus erythropygius*, 346, 378, 380–387
- Yáñez, José L., see Jaksić, Fabian, and —
- Yellow-Finch, Bright-rumped, 478, 480
- Yellowlegs, Greater, 476, 479
- Youngstrom, Kenneth A., see Clay, Deborah L., et al.
- Yukon, 306–318
- Zenaida asiatica*, 158, 161–162, 166
- Z. auriculata*, 477
- Z. macroura*, 158, 499–505
- Zonotrichia albicollis*, 91, 595–597
- Z. atricapilla*, 595–597
- Z. capensis*, 478, 480
- Z. leucophrys*, 160, 392, 499–505, 596–599
- Z. querula*, 88–90, 596–599