

- MYERS, J. P. 1981. A test of three hypotheses for latitudinal segregation of the sexes in wintering birds. *Can. J. Zool.* 59: 1527-1534.
- NICHOLS, J. D., & G. M. HARAMIS. 1980. Sex-specific differences in winter distribution patterns of Can-vasbacks. *Condor* 82: 406-416.
- , & J. E. HINES. 1987. Winter distribution patterns and survival rates of winter-banded Mallards. VIII: Population ecology of the Mallard. *U.S. Fish and Wildl. Serv. Resour. Publ.* 162.
- , K. J. REINECKE, & J. E. HINES. 1983. Factors affecting the distribution of Mallards wintering in the Mississippi Alluvial Valley. *Auk* 100: 932-946.
- OWEN, R. B., JR., & W. B. KROHN. 1973. Molt patterns and weight changes of the American Woodcock. *Wilson Bull.* 85: 31-41.
- PACE, R. M., III, & G. W. WOOD. 1979. Observations of woodcock wintering in coastal South Carolina. *Proc. Southeast Assoc. Fish and Wildl. Agencies.* 33: 72-80.
- PERDECK, A. C., & C. CLASON. 1983. Sexual differences in migration and winter quarters of ducks in the Netherlands. *Wildfowl* 34: 137-143.
- PULLIAM, H. R., & T. A. PARKER III. 1979. Population regulation of sparrows. *Fortschr. Zool.* 25: 137-147.
- PURSCLOVE, S. R., JR., & G. L. DOSTER. 1970. Potentialities of the woodcock as a game bird resource in the southeastern United States. *Proc. Southeast Assoc. Fish Game Comm.* 24: 223-231.
- ROBOSKI, J., & M. K. CAUSEY. 1981. Incidence, habitat use, and chronology of woodcock nesting in Alabama. *J. Wildl. Manage.* 45: 793-797.
- ROBSON, D. S. 1968. The effects of ties on critical values of some two-sample rank tests. Ithaca, New York, Cornell Univ., Biometrics Unit, BU-258-M.
- SHELDON, W. G. 1967. The book of the American Woodcock. Amherst, Univ. Mass. Press.
- STAMPS, R. T., & P. D. DOERR. 1976. Woodcock on North Carolina wintering grounds. *Proc. Southeast Assoc. Game Fish Comm.* 30: 392-399.
- , & —. 1977. Reproductive maturation and breeding of woodcock in North Carolina. *Proc. Woodcock Symp.* 6: 185-190.
- STICKEL, W. H., W. G. SHELDON, & L. F. STICKEL. 1965. Care of captive woodcock. *J. Wildl. Manage.* 29: 161-172.
- STRIBLING, H. L., & P. D. DOERR. 1985. Characteristics of American Woodcock wintering in eastern North Carolina. *N. Am. Bird Bander* 10: 68-72.
- TERRILL, S. B., & R. D. OHMART. 1984. Facultative extension of fall migration of Yellow-rumped Warblers (*Dendroica coronata*). *Auk* 101: 427-438.
- WHITING, R. M., JR., & T. G. BOGGUS. 1982. Breeding biology of American Woodcock in east Texas. *Proc. Woodcock Symp.* 77: 132-138.
- WOOD, G. W., M. K. CAUSEY, & R. M. WHITING JR. 1985. Perspectives on American Woodcock in the southern United States. *Trans. N. Am. Wildl. Conf.* 50: 573-585.

ERRATUM

In "Renesting by American Woodcocks (*Scolopax minor*) in Maine" by Daniel G. McAuley, Jerry R. Longcore, and Greg F. Sepik (1990, *Auk* 107: 407-410), the range of distance moved by females that abandoned nests or had nests destroyed by predators (p. 408) should read: "range = **1.0-15.5**" and the values in the "Distance moved (km)" column in Table 2 (p. 409) should read down the column: "**1.01, 11.34, 1.22, 4.17, 15.54, 0.65, 0.36, 1.14, 0.88, 0.24, 0.80, 0.16**." Corrected values are in boldface type.