

PIRACY OF NESTING MATERIALS FROM AND BY THE BROAD-TAILED HUMMINGBIRD

WILLIAM A. CALDER

Rocky Mountain Biological Laboratory and
Department of Biological Sciences
University of Arizona
Tucson, Arizona 85721

In the course of nesting microclimatic studies of the Broad-tailed Hummingbird (*Selasphorus platycercus*), I observed three cases of intra- or interspecific piracy of nesting materials. These observations were made in the vicinity of the Rocky Mountain Biological Laboratory, Gothic, Gunnison County, Colorado, at an elevation of 2920 m, where this is apparently the only breeding trochilid.

On 17 and 18 June 1971, Broad-tailed females were observed building nests # 1 and 3 approximately 10 m above the ground in quaking aspen trees (*Populus tremuloides*) approximately 40 m apart on a hillside. A storm with rain, hail, and gusty winds occurred the night of 18 June. The following day, nest 1 was apparently abandoned, while a female was removing nest material from nest 1 to nest 3.

On 20 June, a Warbling Vireo (*Vireo gilvus*) was removing large beakfuls of material from the walls of this same hummingbird nest 3. The damage was repaired on 21 June. On 18 July, after at least 6 days during which the female was observed feeding young, the nest walls were again suddenly diminished in height, far more extensively than occurs from normal spreading of the nest cavity as the young grow to fledging size. From 18 July until fledging, 31 July, there was no evidence of major addition to the nest

walls, although the female did attempt some repair or adjustment of what remained. The young birds completed their development in a noticeably shallower nest than others under observation, but were on schedule with other Broad-tailed Hummingbirds that came from nests begun at the same time.

On 20 July, a female hummingbird was removing fine material from the nest of the Western Wood Pewee (*Contopus sordidulus*). The adult peewees made passes on the hummingbird, periodically driving her from their nest. Her approaches to the nest were halting and cautious, apparently because of the large peewee nestlings in the nest. The hummingbird nest, to which the material was presumably taken, could not be located.

Female hummingbirds spend a considerable amount of time hovering under the eaves of buildings and up and down tree trunks to acquire the fine spiderwebs used in the nest lining and walls. The energy saving when nesting material is taken from concentrations in other nests is qualitatively obvious, as is the cost to the loser. If such piracy is widespread, protection not only of eggs and young from predators but of nest material from pilferage should exert a significant effect in natural selection as related to nesting behavior. Legg and Pitelka (Condor 58:393, 1956) observed a female Anna's Hummingbird (*Calypte anna*) who moved nesting material from one of her nests to a second and later to a third. They also observed one clear case of nesting thievery in this species. Nests were not reused or rebuilt at the same site, but material from old nests was transported elsewhere for "recycling."

These observations were supported by NSF grant GB-13249.

Accepted for publication 7 October 1971.

NUTS, BONES, AND A NESTING OF RED CROSSBILLS IN THE PANAMINT MOUNTAINS, CALIFORNIA

ROBERT B. PAYNE

Museum of Zoology
University of Michigan
Ann Arbor, Michigan 48104

In the Panamint Mountains of Death Valley National Monument, California, on 25 March 1970, I saw Red Crossbills (*Loxia curvirostra*) nesting in association with a flush of nuts of pinon pine (*Pinus edulis*). At Mahogany Flats (elevation about 2600 m), the winter snow was melting from the slopes facing south, although banks of snow covered the north-facing slopes. On the exposed soil were thousands of pinon nuts, under each of scores of pine trees. Most of the nuts had fallen free of their cones. It seemed the nuts had scarcely been harvested before snow had covered them earlier in winter, as nearly all of them were intact, although with the water and warm sunlight of spring thaw many nuts were germinating. Feeding on the fallen nuts as well as on nuts still in the cones high in the trees were several Red Crossbills.

In the center of the abundant pine nuts, a male Red Crossbill was singing a trilled song—a soft, descending series of whistled notes. I approached to

within 10 m of the singing male. A female crossbill flew into the next pine with a pine twig and then disappeared into a dense cluster of pine needles in the terminal branches where she added the stick to a nest. The nest tree was about 3 m from the male's song tree, and the nest was 5 m above the ground, which here was covered by a deep snowbank. The nest was a shallow cup, built of pine twigs, not yet lined, and contained no eggs.

Other Red Crossbills sang in trees about 20 m from the nest, but I found no more nests in the area, although some may have been nearby.

Earlier reports of crossbill nesting in California have been in summer, perhaps because of the seasonal activity of local ornithologists. At least two of these three summer records indicate nesting in association with a local abundance of nuts of lodgepole pine (*Pinus contorta*) (Butts 1940; Smith 1940; McMillan 1948). Elsewhere in western North America, the Red Crossbills may nest almost any time of year, as in Colorado they have been seen nesting in winter through spring and early summer (Bailey et al. 1953) and also in late summer (Snyder and Cassel 1951). In eastern North America, 14 of the 15 egg dates recorded by Austin (*in Bent* 1968) fell between February and May. In Europe, the breeding season appears to be better known than it is in North America. Newton (1970, and pers. comm.) found the nesting pattern to be seasonal, with nesting beginning sometimes in late autumn, after com-

pletion of the molt, and sometimes in winter or spring. Molt occurs in summer and breeding does not; in summer the birds often disperse from their breeding grounds, apparently in response to the increased numbers of young following a successful breeding and to the local depletion of conifer seed. Crossbills nest locally in forests with abundant crops of cones (Newton 1970).

Crossbills at Mahogany Flats in the Panamints may have been nesting in response to the flush of food provided by the uncovering of the pine nuts when the spring thaw had melted the snow. At least 20,000 m² of ground there were exposed by the thaw, and most of this area was littered with pinon nuts. Tordoff and Dawson (1965:419-420) have suggested that crossbills may breed in response to an abundance of food, such as a plentiful local cone crop in a year when the pines produce cones, and the appearance of the pinon nuts after spring thaw provides a similar local abundance of crossbill food. Red Crossbills are known to undergo gonadal development in response to spring daylengths when they are tested experimentally in winter (Tordoff and Dawson 1965). It would be of interest to test also the direct effect of pine seeds on their nesting.

An instance of bone-eating at Mahogany Flats may also have been related to the nesting observed there. The melting of snow had uncovered several old feces of a carnivore (perhaps of a coyote, *Canis latrans*), and I saw a female crossbill picking at one. She extracted several bone fragments from it. Examination of the old dropping showed it full of hair and bones of small rodents; the bones were weathered, soft, and crumbling. Crossbills are known to feed on charcoal, salt licks, salt on roads, and urine-stained snow (McMillan 1948; Dawson et al. 1965; Bent 1968), and bones may provide other minerals to the crossbills,

whose seed diet may be low in certain elements. Because the bird eating the bones appeared to be the same female that was building the nest, it seems possible that the extra calcium from the rodent bones might be important in supplying this mineral for formation of egg shells.

LITERATURE CITED

- BAILEY, A. M., R. J. NIEDRACH, AND A. L. BAILY. 1953. The Red Crossbills of Colorado. Denver Mus. Natur. Hist., Mus. Pictorial 9.
- BENT, A. C. 1968. Life histories of North American cardinals, grosbeaks, buntings, towhees, finches, sparrows, and allies. U.S. Nat. Mus. Bull. 237, Pt. 1.
- BUTTS, E. 1940. A sierra crossbill family. Yosemite Nature Notes 19:23-24.
- DAWSON, W. R., V. H. SHOEMAKER, H. B. TORDOFF, AND A. BORUT. 1965. Observations on the metabolism of sodium chloride in the Red Crossbill. Auk 82:606-623.
- McMILLAN, E. 1948. A nesting record for the Red Crossbill in California. Condor 50:44.
- NEWTON, I. 1970. Irruptions of crossbills in Europe, p. 337-357. In A. Watson [ed.] Animal populations in relation to their food resources. Blackwell Sci. Publ., Oxford.
- SMITH, L. M. 1940. The crossbill in Tuolumne Meadows. Yosemite Nature Notes 19:24.
- SNYDER, D. P., AND J. F. CASSEL. 1951. A late summer nest of the Red Crossbill in Colorado. Wilson Bull. 63:177-180.
- TORDOFF, H. B., AND W. R. DAWSON. 1965. The influence of daylength on reproductive timing in the Red Crossbill. Condor 67:416-422.

Accepted for publication 8 October 1971.

NEW RECORDS OF BIRDS FROM BRITISH HONDURAS (BELIZE), INCLUDING A SKUA

JON C. BARLOW,

J. A. DICK,

Department of Ornithology
Royal Ontario Museum
Toronto, Ontario, Canada

DORA WEYER,

AND

W. FORD YOUNG

Belize City
British Honduras

Since the publication of Russell's (1964) distributional account of the birds of that country and of two supplemental papers (Barlow et al. 1969, 1970), important records of birds in British Honduras have continued to accumulate. Two of us (Weyer and Young, as permanent residents of the country, with the aid of several collaborators) have been able to secure most of the specimens mentioned beyond. Likewise, the Royal Ontario Museum (ROM) has maintained a continuing interest in the avifauna of the area. J. A. Dick and Peter L. McLaren, in the course of studying vireos there for that institution between 10 and 31 March 1971, also obtained several

specimens of note. Dick and McLaren worked primarily near the city of Belize and on Ambergris Cay, whereas Weyer and Young have ranged over most of the country. Unfortunately, data for many of the specimens are minimal.

Agami agami. Agami Heron. Dora Weyer secured an adult individual (ROM 109141) of unknown sex at Spanish Creek Lagoon (Western Lagoon on current maps) on 15 April 1969. Russell (op. cit.) regards this species as a local or rare resident and mentions five specimens and several sight records.

Botaurus pinnatus. Pinnated Bittern. An adult of unknown sex (ROM 109140) was found dead by Mrs. Weyer, impaled on a barb-wire fence on 29 March 1970, 1 mile N Hattieville, Belize District. She saw another individual nearby. There is no previous record for the country, but Russell (op. cit.) notes that it has been recorded by Paynter (1955) in nearby Quintana Roo.

Dendrocygna autumnalis fulgens. Black-bellied Tree Duck. Weyer obtained an adult (ROM 109143), apparently a female by measurements and plumage, at the Mussel Creek Rice Station, 4 miles W of Burrell Boom, Belize District. Russell (op. cit.) mentioned three unsexed specimens and two sight records. Barlow et al. (1969) reported two females taken from a flock of 30 in 1967. Continued presence of this species in the country adds support to Russell's suggestion that it may be increasing in numbers there.

Anas carolinensis. Green-winged Teal. Weyer obtained a female (ROM 109142) from a small flock