

lacked the white tips to the primary coverts and white and yellow tips to the primaries that are such a conspicuous feature of the Bohemian Waxwing.

The occurrence of the Cedar Waxwing in central Alaska well over 700 miles north of its recorded range in the state is quite extraordinary. The likely explanation is that the two Cedar Waxwings joined a flock of migrating Bohemian Waxwings and accompanied them north to the Hess Creek area. From the recorded range of the two species, it seems that there is some degree of overlap in winter and un-

doubtedly also at certain times during migration.

It is unlikely that this record represents a northward extension of the range of the Cedar Waxwing in Alaska, but that possibility exists. A similar problem occurs with the Rufous Hummingbird (*Selasphorus rufus*), whose range is given by Gabrielson and Lincoln (op. cit.) as southeastern Alaska north to Prince William Sound. It is now known to occur as far north as the upper Yukon River area (White and Haugh, Can. Field Nat. 83:257, 1969).

Accepted for publication 5 May 1972

OBSERVATIONS ON THE URBAN FEEDING HABITS OF THE ROADRUNNER (*GEOCOCCYX CALIFORNIANUS*)

ROBERT E. WRIGHT

3209-A Via Buena Vista
Laguna Hills, California 92653

During the years 1970 and 1971, extensive observations were made of a pair of Roadrunners (*Geococcyx californianus*) and their young. The birds were frequent, almost daily visitors in the backyard of a home in Woodland Hills, Los Angeles County, California. The home is located on the northern slope of the Santa Monica Mountains at an elevation of about 1200 ft, and the backyard is contiguous with the chaparral growth native to this region. The yard, planted to dichondra with a border of shrubs and annuals, has a population of introduced European snails (*Helix pomata*).

Roadrunners were first noticed in March 1970 (sometimes alone, sometimes as a pair) feeding on the snails, and they have been observed many times during the following months. The bird forages under the shrubbery and flowers in the garden for snails which it carries in its beak to the nearest large rock (sometimes several meters away). With a sideways, snapping motion of its head, it hammers the snail against the rock until the shell is broken into fragments. The soft body is then eaten. After feeding, mucous from the snails is carefully cleaned from the beak by rubbing against the soil. In both 1970 and 1971 parent birds were seen feeding snails to their

half-grown young. This previously unreported feeding behavior may indicate an unsuspected ability of these primarily chaparral- and desert-adapted birds to find acceptable food even after man has altered their normal habitat. A somewhat similar behavior has been reported for the Blackbird (*Turdus merula*) which uses rocks as anvils.

The Roadrunners were also observed capturing small birds. A cement block wall separates the yard from a steep slope which is covered with dense chaparral in the upper portion and cleared of brush in the lower portion. Just beyond the wall a bird feeder, kept supplied with chicken scratch, is utilized by many passerine birds, Mourning Doves (*Zenaidura macroura*), and California Quail (*Lophortyx californicus*). A Roadrunner, attracted by this bountiful prey, takes a position a few feet from the feeder and waits until a group of birds is within about 6 ft of its position, at which time it makes a sudden rush, head down, wings slightly extended. If the attack is successful, the Roadrunner kills its prey with a few stabbing motions of its beak and carries it to the dense chaparral above. On three occasions when prey could be identified, a House Finch (*Carpodacus mexicanus*) and two young quail were captured. Zimmerman (Condor 72:475, 1970) has also described predation on passerine birds by the Roadrunner.

Although many new homes have been built in the immediate area in the last few months, the Roadrunners are still present. Hopefully they may be able to supplement their customary diet with the snails and adapt to the man-made changes in their environment.

Accepted for publication 24 February 1972

TONGUE STRUCTURE OF THE PLUMED WHISTLING DUCK (*DENDROCYGNA EYTONI*)

M. KENT RYLANDER

AND

ERIC G. BOLEN

Department of Biology
and

Department of Range and Wildlife Management
Texas Tech University
Lubbock, Texas 79409

The eight species of whistling or tree ducks (*Dendrocygna* spp.) form a closely related group whose phylogenetic status is unchallenged (cf. Delacour

1954:26; Johnsgard 1968:6; and others). Despite their taxonomic similarity, however, a significant degree of ecological and behavioral diversity exists among these forms as shown in the functional anatomy of the North American species discussed by Rylander and Bolen (1970).

Our observations of the feeding apparatus of this group indicate a general uniformity with regard to the tongue structure consistent with a basic similarity in their diets (i.e., seeds from either aquatic or terrestrial sources). We were thus surprised to note the presence of a remarkably developed fimbriated lateral margin on an anterior process of the tongue of the Plumed Whistling Duck (*D. eytoni*). This condition, not previously reported elsewhere, was discovered on specimens kindly sent to us from Australia by H. J. Frith. A fimbriated tongue margin in this position has not been reported for anatids, and