

## WINTERING WADERS IN SOME COASTAL WETLANDS OF NORTH-EAST ITALY

Roberto Tinarelli

In December, January and February 1984-85 and 1985-86 new census work was undertaken between Adige mouth (45°08'N - 12°20'E) and Saline di Cervia (44°16'N - 12°23'E), with the aim of obtaining further data on the numbers of wintering waders in the area. The surveyed area includes sandy beaches, shallow bays and lagoons subject to the tidal movements, as well as brackish and freshwater lakes up to 10 km inland. Counts were made at least once a month. When several counts were made within the same month, minimum and maximum values are reported. Data were collected on a total of 28 days (14 each winter).

Totals of birds counted and estimated in both winters are summarized in Tables 1 and 2. A more detailed analysis of these results will soon be available (Tinarelli in press).

The decrease in bird numbers after the December counts, a feature already found for the wintering population of the Avocet

*Recurvirostra avosetta* (Tinarelli 1987), was probably due to late migratory movements.

In the winter of 1984-85, an unusual cold spell severely affected the number of birds observed in January. Temperatures remained well to below -15°C for half a month, and caused complete freezing of inland waters and salt-pans. By the end of the same month, consequently, waders were only present along sandy beaches.

In normal conditions, large numbers of waders (45-50%) were observed on sandy islands in the Po Delta, salt-pans (40% of the total counted) and inland lagoons (10-15%). Salt-pans, in particular, with 600 ha suitable for waders, seemed to be one of the best habitats.

Hunting activity and poaching are probably the main factors affecting the winter population, followed by beach erosion.

Data obtained during this research indicate

Table 1. Total waders counted in the winter 1984-85.

	December	January	February
<i>Haematopus ostralegus</i>	5-6	1	7
<i>Pluvialis squatarola</i>	579-602	281-326	280-310
<i>Charadrius hiaticula</i>	4		
<i>Charadrius alexandrinus</i>	135-165	40-50	35-55
<i>Numenius arquata</i>	298-323	21-25	182-222
<i>Limosa lapponica</i>	16		
<i>Limosa limosa</i>	25		
<i>Philomachus pugnax</i>	83-85		
<i>Tringa totanus</i>	86-107	32-52	2
<i>Tringa erythropus</i>	243-273		8-12
<i>Tringa nebularia</i>	9	3	
<i>Calidris alpina</i>	4 724-5 146	2 345-2 655	2 780-3 130
<i>Calidris minuta</i>	535-585		
<i>Calidris temminckii</i>	21		
<i>Calidris alba</i>	174-194	140	286-392
<i>Recurvirostra avosetta</i>	110-150	6	10
Total inds. counted	7 042-7 711	2 869-3 258	3 590-4 150
Estimated number	8 900-9 500	4 250-4 800	5 600-6 250

Table 2. Total waders counted in the winter 1985-86.

	December	January	February
<i>Haematopus ostralegus</i>	4-6	5-8	
<i>Pluvialis squatarola</i>	698-738	707-747	612-647
<i>Charadrius hiaticula</i>	6	12	
<i>Charadrius alexandrinus</i>	80-99	80-125	130-150
<i>Numenius arquata</i>	177-197	159-174	185-218
<i>Limosa lapponica</i>			
<i>Limosa limosa</i>	22	5	30
<i>Philomachus pugnax</i>	8	24	29
<i>Tringa totanus</i>	110-120	144-164	57-60
<i>Tringa erythropus</i>	206	120-135	41-50
<i>Tringa nebularia</i>	2	3	2
<i>Calidris alpina</i>	5 908-6 140	5 950-6 380	5 466-5 836
<i>Calidris minuta</i>	600-800	560-800	490-599
<i>Calidris temminckii</i>	43	18	
<i>Calidris alba</i>	147-150	80-100	95
<i>Recurvirostra avosetta</i>	217-297	200-220	205-210
Total inds. counted	8 232-8 834	8 067-8 915	7 342-7 927
Estimated number	11 900-12 700	9 900-10 900	9 800-10 600

significantly higher numbers than the 2 000-5 400 birds reported by Perco (1984) for the same area. According to Smit's (1986) estimates, the areas hold 25-30% of the total Italian wintering population.

#### REFERENCES

Perco, F. 1984. Estimates of wader numbers during midwinter in northern Adriatic coastal wetlands. *Wader Study Group Bull.* 40: 49-50.

Smit, C.J. 1986. Wintering and migrating waders in the Mediterranean. *Wader Study Group Bull.* 46: 13-15.

Tinarelli, R. 1987. Aspetti della biologia invernale dell'Avocetta *Recurvirostra avosetta* in alcune zone umide costiere del nord Adriatico. *Avocetta* 11: 37-45.

Tinarelli, R. in press. Svernamento dei limicoli in alcune zone umide costiere dell'Adriatico settentrionale. *Avocetta*.

Robert Tinarelli, Via Vasari 17, 40128 Bologna, Italy

## AN HISTORICAL OVERVIEW OF WADER RINGING IN ITALY

Nicola Bacetti, Lionello Bendini & Lorenzo Serra

Bird ringing started in Italy in 1929, thanks to the efforts of several independent ringing stations, each one producing and using their own rings. A more centralized organization, and the existence of a co-ordinated national scheme, gradually changed the early situation. However, a complete change was only achieved quite recently.

The totals of birds ringed during the whole period are not known, and only for the last 10 years could some figures be collated. Only recent data have been stored in computerized archives (the first available year was 1982).

An analysis of microfilm containing the original lists of rings used, combined with a few unpublished early reports, allowed us to assess the actual number of waders ringed up to 1985, with the aim of providing useful indications for planning future research.

A total figure of 35 163 shorebirds ringed was found. An historical analysis of these data, arranged in 5-year periods, showed a very irregular distribution, with numbers exceeding 4 500 as early as 1935-40. This total was only exceeded in 1966-70 and the following periods. Due to the low numbers concerned, it is difficult to explain the reasons for such temporal patterns. The activity of individual ringers may have strongly affected the overall national results. However, it seems quite understandable that the lowest totals were during the two 5-year periods covering the last war and the following years (1 274 and 953 birds respectively), when a different use of trapped birds was much more likely to be preferred.

The most frequently ringed species, Lapwing *Vanellus vanellus* and Ruff *Philomachus pugnax*, with 35.5% and 23.7% respectively of the total numbers, show historical patterns quite similar to the general one. On the other hand, some more interesting situations are noted here. Both the Common Snipe *Gallinago gallinago* and Jack Snipe *Lymnocyptes minimus* were ringed in good numbers in the 1930s thanks to traditional methods which ceased later. Ringing results decreased quite suddenly afterwards: 5-year totals dropped from 174 and 261 birds of the former species (1931-40), and 53, 58, 20 of the latter (1931-45), to 3-32 and 0-1 respectively in all but the most recent periods. The recent more widespread use of mist-nets by Italian ringers has increased the number of Common Snipe ringed to 259 in 1980-85. However, only 3

Jack Snipe were ringed in the same period, which provides some evidence of a real decrease of this bird.

The ringing of Avocets *Recurvirostra avosetta* and Black-winged Stilts *Himantopus himantopus* started quite late, in the 1950s and 1960s respectively. An increase of the Italian breeding populations (the Avocet was not previously breeding in Italy at all) has resulted in a gradual increase to the present totals of 338 Avocets and 437 Stilts.

The month of capture of the latter two species falls in most cases within the range of the breeding season, since unfledged juveniles represent about 80% of Black-winged Stilts, and probably 100% of Avocets ringed. For other species, when adult birds are more regularly involved, the months of capture reflect quite precisely the phenology of migration or the pattern of presence in the country. This is the case, for instance, of a typical spring migrant, the Black-tailed Godwit *Limosa limosa*: all 379 birds were ringed in February-May (53.5% in March).

As a final comment, it should be stressed that the numbers of Italian-ringed waders are in most cases still too low (and wader-ringers still not so specialized) to be of major importance in internationally co-ordinated programs. However, what seems possible is to increase the value of results by concentrating the ringers' attention on very few species at one time. The Ruff and Black-winged Stilt, after first promising experiences, will probably deserve major attention from ringers during the next few years.

Nicola Bacetti, Lionello Bendini and Lorenzo Serra, Istituto Nazionale Biologia Selvaggina, Via Stradelli Guelfi, I-40064 Ozzano Emilia BO, Italy