
The primary moult of breeding Dunlins *Calidris alpina* in the central Taymyr in 1989

Wojciech Kania

Kania, W. 1990. The primary moult of breeding Dunlins *Calidris alpina* in the central Taymyr in 1989. *Wader Study Group Bull.* 60: 17 - 19

A total of 14 incubating Dunlins were caught at the mouth of Malaja Logata, central Taymyr, in July 1989. All were moulting the primaries, starting with the simultaneous loss of the first three, four or five primaries. Primary moult lasts probably c. 50 days, considerably less than the primary moult period reported for post-breeding Dunlins.

Wojciech Kania, Stacja Ornitologiczna IZ PAN, 80-680 Gdansk 40, Poland.

INTRODUCTION

In some populations Dunlins moult while they are breeding (Kozlova 1962; Gromadzka 1989). Moulting at breeding sites has been found east of the Urals (Greenwood 1983; Danilov *et al.* 1984; Gromadzka 1989), although in the Yamal Peninsula only a small fraction of the breeding birds was involved (Ryabitsev *in litt.*, cited by Gromadzka 1989).

Kozlova, Greenwood and Gromadzka based their statements on only a few observations (Greenwood for the area from Yenisey to Kamchatka - 11 cases, Gromadzka for Taymyr - 8 cases) from various years. These were mainly from museum specimens, and some specimens examined by each author were often the same. This paper reports data on primary moult collected during the breeding season in Taymyr in 1989. The sample size is not much bigger than those of earlier studies, but the data has the advantage of being gathered at one site and during one season.

This paper is one of the results of an ornithological investigation organised by the Soviet Academy of Sciences, and carried out jointly by the WWF-Wattenmeerstelle Schleswig-Holstein, the Ukrainian Black/Azov Seas Ornithological Station and the Gdansk Ornithological Station (Polish Academy of Sciences).

MATERIALS AND STUDY AREA

The material consists of 17 moult records and sets of measurements for 14 Dunlins caught, while incubating, at nine nests at the mouth of the Malaja Logata (73°25' N 98°25'

E) in the Taymyr tundra, in July 1989. According to members of the staff of the "Taymyrskiy" Nature Reserve, the summer of 1989 began 10-15 days later than usual.

METHODS

Dunlins were caught at their nests, using walk-in traps. Sex was determined according to the presence (male) or absence (female) of contrast between the greyish hindneck and brownish cap (Soikkeli 1966). Measurements of males and females (Table 1) placed in a bivariate plot of bill versus wing length, form separate clusters that confirm sex determination - males being smaller than females. Wing-length was measured using the maximum chord method to 1 mm. Bill-length, was measured as exposed culmen from the feather base. This and head and bill length was usually taken to 0.1 mm.

Moult score (Ashmole 1962) was noted for all 10 main primaries (Table 1).


To illustrate the moult process (Figure 1), I estimated the amount of new feather mass produced (Underhill & Zucchini 1988). Moult score for each bird was recalculated for feather mass by the formula

$$y = \sum_{i=1}^{10} m_i p_s$$

where y is the proportion of total mass of ten primaries in relation to their hypothetical mass when fully grown, m_i is the



Figure 1. Primary moult of incubating Dunlins at the mouth of the Malaja Logata, Taymyr, 1989 in relation to date. Moulting stage expressed as a percentage of total mass of ten fully grown primaries (see text). Shaded circles are males; open circles are females. Multiple records of the same individual are joined by a line. The code numbers (see Table 1) of the birds caught on the nest with known hatching dates are given.



mass of the i th primary relative to the total mass of all the primaries, p_s = the mass of a primary with score s relative to its mass when it is fully grown (0, 0.125, 0.375, 0.625, 0.875, 1 for successive scores 0, 1, ..., 5 after Underhill & Zucchini 1988, with a little simplified notation). Data for the primary feather masses of Dunlins were taken from measurements on five birds caught in Poland during autumn migration (Table 2).

RESULTS AND DISCUSSION

All breeding Dunlins found at the mouth of the Malaja Logata

in 1989 moulted their primaries during incubation. They started to moult at the end of June or the beginning of July (Figure 1).

The beginning of the moult in relation to the start of breeding can be assessed for four Dunlins caught at three nests with known hatching dates (Table 1). In *C. alpina schinzii* in Finland incubation lasts c.22 days (Soikkeli 1967). In a more severe climate it may take a few days more, as Tomkovich (1988) found for Temminck's Stint *Calidris temminckii*. In these four Dunlins the beginning of the moult coincided more or less with egg laying, which is in agreement with the Kozlova's (1962) statement.

Almost all Dunlins had the first three, four or five (usually four) primaries in the same stage of development. In all cases the inner primaries with the same score were of equal length. In Dunlin no. 10, the vanes of the second and third primaries were just emerging from the sheath. This points to the fact that corresponding old feathers were lost simultaneously. Kozlova (1962) claimed simultaneous moult usually for the first three or four primaries, but sometimes the first five or six. Such a rapid start of primary moult has never been recorded in migrating Dunlins caught in the first stages of moult at the Vistula mouth in Poland (J. Gromadzka, pers. comm.).

The moult progressed very quickly in the Dunlins in the Taymyr which were studied in 1989. It is not known if moult was suspended after the end of incubation. If moult continues at a similar rate in the second part of the moult period, the

Table 1. Primary moult scores and measurements of breeding Dunlins at the mouth of the Malaja Logata, Taymyr, 1989.

| Specific number of bird | nest | Sex | Length of wing | bill* head +bill | Weight | Date (July) | Hour | total | Moult score for individual feathers | | | | | | | | | | Hatching date (July) | | |
|-------------------------|------|-----|----------------|------------------|--------|-------------|------|-------|-------------------------------------|---|---|---|---|---|---|--------------|---|----|----------------------|----|--|
| | | | | | | | | | 1 | 2 | 3 | 4 | 5 | 6 | 7 | 8 | 9 | 10 | | | |
| 1 | 1 | M | 119 | 30.3 | 54.4 | 46 | 2 | 16 | 4 | 1 | 1 | 1 | 1 | | | | | | | 20 | |
| 2 | 2 | M | 121 | 31.7 | 56.5 | 54 | 4 | 15 | 8 | 2 | 2 | 2 | 2 | | | | | | | | |
| 3 | 3 | M | 119 | 31.8 | 55.5 | 51 | 4 | 23 | 8 | 2 | 2 | 2 | 2 | | | | | | | | |
| 4 | 4 | F | 119 | 34.9 | 58.3 | 53 | 5 | 23 | 4 | 1 | 1 | 1 | 1 | | | | | | | | |
| 5 | 4 | M | 114 | 32.3 | 57.4 | 41 | 6 | 15 | 9 | 2 | 2 | 2 | 2 | 1 | | | | | | | |
| 6 | 5 | F | 119 | 33.5 | 58.3 | 48 | 6 | 11 | 18 | 4 | 4 | 4 | 3 | 2 | 1 | OLD FEATHERS | | | | | |
| 7 | 6 | F | 124 | 37.2 | 61.6 | 60 | 9 | 17 | 9 | 3 | 3 | 2 | 1 | | | | | | | | |
| 8 | 3 | F | 118 | 34.1 | 58.6 | 53 | 9 | 18 | 9 | 2 | 2 | 2 | 2 | 1 | | | | | | | |
| 9 | 7 | F | 120 | 33.3 | 57.5 | 58 | 9 | 21 | 9 | 2 | 2 | 2 | 2 | 1 | | | | | | 26 | |
| 10 | 8 | F | 121 | 36.6 | 61.4 | 55 | 9 | 23 | 7 | 1 | 2 | 2 | 1 | 1 | | | | | | | |
| 11 | 7 | M | 119 | 32 | 56.0 | 49 | 12 | 17 | 14 | 3 | 3 | 3 | 2 | 2 | 1 | | | | | 26 | |
| 11 | | | | | | 47 | 21 | 22 | 29 | 5 | 5 | 5 | 5 | 4 | 4 | 1 | | | | | |
| 11 | | | | | | 52 | 26 | 14 | 32 | 5 | 5 | 5 | 5 | 5 | 4 | 3 | | | | | |
| 12 | 9 | F | 120 | 32 | 57.6 | 52 | 12 | 22 | 22 | 4 | 4 | 4 | 4 | 3 | 2 | 1 | | | | | |
| 12 | | | | | | | 14 | 3 | 24 | 4 | 4 | 4 | 4 | 4 | 3 | 1 | | | | | |
| 13 | 1 | F | 124 | 36.6 | 61.3 | 59 | 13 | 4 | 21 | 4 | 4 | 4 | 4 | 3 | 2 | 1 | | | | 20 | |
| 14 | 9 | M | 114 | 33 | 57.2 | 49 | 14 | 20 | 28 | 5 | 5 | 5 | 5 | 4 | 3 | 1 | | | | | |

* correct to 0.1 mm or 1 mm.



Table 2. Mass of the primaries of two adult (more than one year old) specimens of Dunlins caught at the Vistula mouth, Poland, on 17 August 1988 and 29 August 1989.

| Primary moult (<i>i</i>) | 1 | 2 | 3 | 4 | 5 | 6 | 7 | 8 | 9 | 10 | 1-10 |
|---|-----|------|------|------|------|------|------|------|------|------|-------|
| minimal | 6.5 | 7.6 | 9.1 | 11.1 | 13.4 | 16.4 | 19.7 | 23.4 | 26.6 | 31.9 | 165.7 |
| Mass (mg) | | | | | | | | | | | |
| maximal | 8.3 | 10.3 | 11.6 | 14.2 | 17.5 | 20.5 | 23.8 | 27.6 | 32.5 | 36.9 | 203.2 |
| Mean percentage of the primary mass (m_i) | 4.0 | 4.7 | 5.6 | 6.8 | 8.4 | 10.3 | 11.8 | 13.8 | 15.8 | 18.7 | 100.0 |

primary moult would take c.50 days. This is a markedly shorter period than the 59-94 days reported for primary moult in post-breeding Dunlins (Ginn & Melville 1983).

ACKNOWLEDGEMENTS

Jadwiga Gromadzka, Przemyslaw Chylareck, Nick Davidson and Hermann Hötcker gave valuable comments on earlier drafts of this paper. Members of the WWF expedition from Husum (FRG) - Peter Prokosch, Holger A. Bruns, H. Hoetker, Willi Knief, Johan Mooij - and members of the expedition of the Black/Azov Seas Ornithological Station (Ukraine), Valerij Siokhin, Igor Belashkov and Tatiana Kirikova - participated in the collection of data. I am grateful to all individuals and organisations who made our research in Taymyr possible and gave us logistic support. These included Professor Evgeniy Syroechkovsky and his staff at the Institute of Evolutionary Animal Morphology and Ecology, USSR Academy of Sciences, and Director Juriy Karbainov, Warden Victor Khristophorof and staff of the "Taymyrskiy" Nature Reserve.

REFERENCES

- Ashmole, N.P. 1962. The Black Noddy *Anous tenuirostris* on the Ascension Island. Part I. General Biology. *Ibis* 103: 235-273.
- Ginn, H.B., & Melville, D.S. 1983. *Moult in birds*. BTO Guide 19. British Trust for Ornithology, Tring.
- Greenwood, J.G. 1983. Post-nuptial primary moult in Dunlin *Calidris alpina*. *Ibis* 125: 221-228.
- Danilov, N.N., Ryzhanovskij, V.N., & Ryabitsev, V.K. 1984. *Birds of Yamal*. Nauka, Moskva (in Russian).
- Gromadzka, J. 1989. Breeding and wintering areas of Dunlin

migrating through southern Baltic. *Ornis Scand.* 20: 132-144.

Kozlova, E.V. 1962. *Fauna SSSR. Birds*. Vol. 2, subvol. 1, part 3. Nauka, Moskva, Leningrad (in Russian).

Soikkeli, M. 1966. On the variation in bill- and wing-length of the Dunlin (*Calidris alpina*) in Europe. *Bird Study* 13: 256-269.

Soikkeli, M. 1967. Breeding cycle and population dynamics in the Dunlin (*Calidris alpina*). *Ann. Zool. Fenn.* 4: 158-198.

Tomkovich, P.S. 1988. On the originality of breeding biology of Temminck's Stint *Calidris temminckii* at the northern limit of its area. *Ornitologiya* 23: 188-193.

Underhill, L.G., & Zucchini, W. 1988. A model for avian primary moult. *Ibis* 130: 358-372.

