

RECENT PUBLICATIONS ON WADERS 64

compiled by
Henk Koffijberg

In trying to make these lists as complete as possible, and the next one even more complete and up to date, your help is most welcome. Please send new titles, abstracts, reprints, copies, reports, and (important too!) omissions or corrections to:

Henk Koffijberg, Glenn Millerweg 102,
1311 RW Almere, The Netherlands.

Remember to include, if possible, the address of the first author, and translations of titles if they are in other languages than English.

For help with this issue I am very grateful to J. Van Impe, J. Meltzer, O. Opermanis, A. Aunins, J. Figueroa, D.A. Stroud, I. Bainbridge, T. Piersma and H. Hötker.

Translated titles from languages other than English are bracketed; with round brackets if the translation is by the author(s), square brackets if not.

N.B. a copy of your paper or abstract would be very helpful to enable me to add some keywords or give added annotation to the titles.

So far, about 4,200 titles have been stored in the WSG/RPW database. Keep sending more!

BREEDING

Bairlein, F. & Bergner, G. 1995. Vorkommen und Bruterfolg von Wiesen-vögeln in der nördlichen Wesermarsch, Niedersachsen. (Occurrence and breeding success of meadow birds in northwestern Lower Saxony). *Vogelwelt* 116(2): 53-59. In German with English summary. (Inst. Vogelforschung Vogelwarte Helgoland, An der Vogelwarte 21, D-26386 Wilhelmshaven, Germany). *Vanellus vanellus*, *Limosa limosa*, *Tringa totanus*, *Haematopus ostralegus* and *Gallinago gallinago*.

Broyer, J. & Benmergui, M. 1995. [Breeding of the Lapwing *Vanellus vanellus* in The Dombes, southeastern France: results and reasons for failures]. (In: Dejonghe, J.-F. (ed.). Proceedings 21st French Ornithological Symposium, Paris, 5-6 March 1994). *Alauda* 63(1): 68. In French (abstract only). (O.N.C. Stn. Dombes,

F-01330 Villars Dombes, France).

Brunner, H. 1994. [Juvenile development of the Dotterel *Charadrius morinellus* in the Austrian Alps]. *Limicola* 8(1): 15-27. In German with English summary. (Karl-Franzens Univ., Inst. Zool., Abt. Morphol. & Oekol., Universitätsplatz 2, A-8010 Graz, Austria).

Buczek, T. & Buczek, A. 1993. [Breeding waders of carbonate marshes near Chełm]. *Notatki Ornitol.* 34(3-4): 287-298. In Polish with English summary. (ul. W. Zywnego 1/32, 20-854 Lublin, Poland). Data on eight wader species.

Davis, A. 1994. Breeding biology of the New Zealand Shore Plover *Thinornis novaeseelandiae*. *Notornis* 41(Suppl.): 195-208. (Dept. Zool., Univ. Auckland, Private Bag, Auckland, New Zealand).

Flodin, L.-A., Hirsimäki, H. & Norén, L.-G. 1995. Samhäckande vadare på strandängar i sydvästra Sverige: predationsskydd eller habitatval? (Associative breeding of waders on shore meadows in southwestern Sweden: predator protection or habitat selection?) *Ornis Svecica* 5(1): 23-30. In Swedish with English summary. Tofta 4224, S-432 95 Varberg, Sweden). Association, breeding success and habitat selection of three non-aggressive species (*Tringa totanus*, *Calidris alpina schinzii* and *Philomachus pugnax*) were studied in an area where two aggressive species (*Vanellus vanellus* and *Limosa limosa*) also bred. Better nest protection by higher vegetation was more important than predator protection given by the aggressive species.

Minton, C., Pearson, G. & Lane, J. 1995. History in the mating: Banded Stilts do it again. *Wingspan* 5(2): 13-15. (165 Dalgety Rd., Beaumaris, VIC 3193, Australia). About 35,000 nests of *Cladorhynchus leucocephalus* were counted at recently flooded salt lakes in the Kalgoorlie region, Western Australia.

Opermanis, O. & Aunins, A. 1995. (Breeding biology of lapwing *Vanellus vanellus* in habitats with different human impact). *Putni daba* 5(1): 2-16. In Latvian with Russian and English summary. (Zool. Mus., Latvijas Univ., Kronvalda bulv. 4, Riga, LV-1586, Latvia). Mean clutch size, egg volume and hatching success figures were obtained for nesting *Vanellus vanellus* and compared with arable farmland and other breeding habitats such as coastal meadows, lake islands and pastures.

Paassen, A. van 1995. Aanzet tot het bepalen van het broedsucces van de Grutto. [How to determine the breeding success of Black-tailed Godwits *Limosa limosa*]. *Vogeljaar* 43(3): 97-104. In Dutch. (Actiegroep Grutto, c/o P.O. Box 558, NL-2800 AN Gouda, Netherlands).

Smith, R.D. & Whitfield, D.P. 1995. Renesting by male Dotterel *Charadrius morinellus* after successfully rearing chicks. *Bird Study* 42(2): 174-175. (Appl. Ornithol. Unit, Dept. Zool., Univ. Glasgow, Glasgow G12 8QQ, UK).

Tomkovich, P.S. 1995. Second report on research on the Great Knot *Calidris tenuirostris* on the breeding grounds. *Stilt* 26: 58-60. (Dept. Ornithol., Zool. Mus., Moscow State Univ., Herzen St. 6, Moscow 103009, Russia). Information on site faithfulness and mate fidelity, breeding density, phenology, breeding success, food and food resources, on the eastern spur of the Shchuchy Mountain Range, north-eastern Siberia.

Zöllner, T. 1995. Kiebitze (*Vanellus vanellus*) kopulieren während der Brutablösung. (Lapwings (*Vanellus vanellus*) copulate during breeding relief). *J. Orn.* 136(3): 296-299. In German with English summary. (Hauptstr. 46, D-02694 Guttau, Germany).

BEHAVIOUR, ECOLOGY & EVOLUTION

Balachandran, S. & Natarajan, V. 1992. Unusual behaviour or adaptation against predation in Terek Sandpiper *Tringa terek* (Latham). *J. Bombay Nat. Hist. Soc.* 89(3): 373-374.

Belanger, R.A. 1995. Weyburn shorebird study: 1990-1992. *Blue Jay* 52(1): 36-42. (412 McGillivray Dr., Weyburn, SN, Canada S4H 1M1).

Boettcher, R., Haig, S.M. & Bridges, W.C. 1995. Habitat-related factors affecting the distribution of nonbreeding American Avocets in coastal South Carolina. *Condor* 97(1): 68-81. (Natl. Biol. Surv., Oregon State Univ., 3200 S.W. Jefferson Way, Corvallis 97331, USA). *Recurvirostra americana*.

Bukacinski, D. & Bukacinska, M. 1994. (Factors influencing changes in the abundance and distribution of gulls, terns and plovers nesting on the mid-Vistula). *Notatki Ornitol.* 35(1-2): 79-97. In Polish with English summary. (Inst. Ekologii PAN, Dziekanow Lesny, 05-092 Lomianki, Poland). *Charadrius dubius* and *C. hiaticula*.

Clausager, I. 1994. *Wing survey from the hunting season 1993/94 in Denmark*. Report No. 115. Danish Ministry for Environment. 52 pp. Danish, French, and English summary. *Scolopax rusticola*, *Gallinago gallinago* and *Lymnocryptes minimus*.

Fair, J.M., Kennedy, P.L. & McEwen, L.C. 1995. Diet of nesting Killdeer in North Dakota. *Wilson Bull.* 107(1): 174-177. *Charadrius vociferus*.

Fiske, P. & Kálás, J.A. 1995. Mate sampling

- and copulation behaviour of Great Snipe females. *Anim. Behav.* 49(1): 209-219. (Dept. Zool., Univ. Trondheim, N-7055 Dragvoll, Norway). *Gallinago media*, Norway.
- Fiske, P., Kálás, J.A. & Saether, S.A. 1994. Correlates of male mating success in the lekking Great Snipe (*Gallinago media*): results from a four-year study. *Behav. Ecol.* 5(2): 210-218.
- Gavrilov, V.V. 1994. Dependency of the maximal life span on the type of social organization in plovers (Charadriidae, Aves). *Dokl. Biol. Sci.* 339(1-6): 569-571. (Koltsov Inst. Dev. Biol., Russian Acad. Sci., Moscow, Russia).
- Gerritsen, G.J. & Groen, N.M. (eds.) 1995. *Icelandic Black-tailed Godwit Project 1993*. Zeist: WIWO (Foundation Working Group International Wader and Waterfowl Research). WIWO-report 51. 44 pp. (Report can be ordered by paying Dfl. 15, plus Dfl. 15 administration costs, to postal giro account 2 666 009 or to ABN bank account 57.02.16.613 of WIWO, Lindengracht 9, NL-3633 AS Vreeland, Netherlands). *Limosa limosa islandica*. Data were collected on breeding densities, habitat and nest descriptions, clutch and egg sizes, timing of breeding, biometrics and blood samples of breeding birds. On the basis of recent counts the Icelandic population is estimated to be between 21,900 - 28,000 birds.
- Goss-Custard, J.D., Caldow, R.W.G., Clarke, R.T., Durell, S.E.A. Le V. Dit & Sutherland, W.J. 1995. Deriving population parameters from individual variations in foraging behaviour. I. Empirical game theory distribution model of Oystercatchers *Haematopus ostralegus* feeding on mussels *Mytilus edulis*. *J. Anim. Ecol.* 64(2): 265-276. (Inst. Terrestrial Ecol., Furzebrook Res. Stn., Wareham, Dorset BH20 5AS, UK).
- Goss-Custard, J.D., Caldow, R.W.G., Clarke, R.T. & West, A.D. 1995. Deriving population parameters from individual variations in foraging behaviour. II. Model tests and population parameters. *J. Anim. Ecol.* 64(2): 277-289. *Haematopus ostralegus*.
- Goss-Custard, J.D., Caldow, R.W.G., Clarke, R.T., Durell, S.E.A. Le V. Dit, Urfi, J. & West, A.D. 1995. Consequences of habitat loss and change to populations of wintering migratory birds: predicting the local and global effects from studies of individuals. In: *Bird conservation: the science and the action: proceedings of the annual conference of the British Ornithologists' Union, 6-10 April 1994, Shuttleworth College, Bedfordshire, UK*. Coulson, J. & Crockford, N.J. (eds.). *Ibis* (Suppl. 1): 56-66. *Haematopus ostralegus*.
- Harradine, J. 1994. Woodcock wing survey 1993/94 in Britain and Ireland. *IWRB Woodcock and Snipe Res. Group Newsletter* 20: 3-6. (BASC, Marford Mill, Rossett, Wrexham, Clwyd LL12 0HL, UK). Information on weight, wing length and condition of adult and juvenile *Scolopax rusticola* shot in the UK.
- Harrison, F. 1995. Prolonged aggression between Eastern Curlews *Numenius madagascariensis*. *Stilt* 26: 35. (4/6 Albert St., Cranbook, Qld 4814, Australia).
- Hötker, H. 1995. Aktivitätsrhythmus von Brandgänsen (*Tadorna tadorna*) und Watvögeln (Charadrii) an der Nordseeküste. (Activity rhythms of Shelducks (*Tadorna tadorna*) and waders (Charadrii) on the North Sea coast). *J. Orn.* 136(2): 105-126. In German with English summary. (Forschungs- und Technologiezentrum Westküste Univ. Kiel, Hafenstr. 1, D-25761, Büsum, Germany). *Haematopus ostralegus*, *Charadrius hiaticula*, *Pluvialis squatarola*, *Calidris canutus*, *C. alpina*, *Philomachus pugnax*, *Limosa lapponica*, *Numenius arquata*, *Tringa erythropus*, *T. totanus* and *T. nebularia*.
- James, R.A. Jr. 1995. Natal philopatry, site tenacity, and age of first breeding of the Black-necked Stilt. *J. Field Ornithol.* 66(1): 107-111. (US Fish & Wildl. Serv., Ecol. Serv., 2730 Loker Ave. W., Carlsbad, CA 92008, USA). *Himantopus mexicanus*, USA.
- Jukema, J., Piersma, T., Louwsma, L., Monkel, C., Rijpma, U., Visser, K. & Zee, D. van der 1995. Rui en gewichtsveranderingen van doortrekende Kemphanen in Friesland in 1993 en 1994. (Moult and mass changes of northward migrating Ruffs in Friesland, March-April 1993 and 1994). *Vanellus* 48(3): 55-61. In Dutch with English summary. (Haerdewei 44, NL-8854 AC Oosterbierum, Netherlands). Progress report on a co-operative study on moult, plumage and body mass variation in northward migrating *Philomachus pugnax* in the province of Friesland, The Netherlands.
- Kasperek, M. 1990. On the migration of the Whimbrel, *Numenius phaeopus*, in Turkey. *Zool. Middle East* 4: 25-32.
- Kim, E.-Y. & Won, P.-O. 1993. [Ecology of the waders migrating to Kanghwa and Yongjong Islands on the west coast of Korea]. *Bull. Inst. Ornithol. Kyung Hee Univ.* 4: 25-46. In Korean with English summary. (Inst. Ornithol., Kyung Hee Univ., Seoul 130-701, Korea). Information on 32 wader species.
- Kivit, H., Nijmeijer, H. & Ovaa, A. (eds.) 1994. *Wader and waterfowl migration in the Çukurova delta, South Turkey, spring 1990*. Zeist: WIWO (Foundation Working Group International Wader and Waterfowl Research). WIWO-report 48. 161 pp. (Report can be ordered by paying Dfl. 25, plus Dfl. 15 administration costs, to postal giro account 2 666 009 or to ABN bank account 57.02.16.613 of WIWO, Lindengracht 9, NL-3633 AS Vreeland, Netherlands). Aim of the project was to study the international importance of the deltas for waders and other waterbirds; to obtain data about the migration strategies of passage waders especially information on origins and timing, and with special attention to those waders using both the east Atlantic flyway and the Mediterranean flyway; to study the use of different wetland habitats by these birds in connection with changing weather conditions and changes in habitat; and to use the results to promote better protection.
- Krementz, D.G. & Pendleton, G.W. 1994. Diurnal habitat use of American Woodcock wintering along the Atlantic coast. *Can. J. Zool.* 72(1): 1945-1950. (Natl. Biol. Serv., Southeast Res. Group, D.B. Warnell School For. Resour., Univ. Georgia, Athens 30602-2152, USA). *Scolopax minor*, USA.
- Le dean Quenec'Hdu, S., Boret, P. & Mahéo, R. 1994. [The importance and spatial use of the Mont Saint-Michel Bay, western France - a study of three wader species: Black-tailed and Bar-tailed Godwits *Limosa limosa* and *L. lapponica* and the Knot *Calidris canutus*]. *Alauda* 62(4): 257-268. In French with English summary. (Lab. Evol. Syst. Nat. & Modifies, Campus Sci. Beaulieu, Univ. Rennes I, F-35042 Rennes Cedex, France).
- Meltofte, H. 1993. (Wader migration through Denmark: populations, non-breeding phenology and migratory strategies). *Dansk. Orn. Foren. Tidsskr.* 87: 1-180. In Danish with English summary. See review in *Bulletin* 76.
- Meran, P. 1994. Some observations on Woodcock (*Scolopax rusticola*) migration in Austria and western Hungary in 1993. *IWRB Woodcock and Snipe Res. Group Newsletter* 20: 27. (Elisabethstr. 41, A-8020 Graz, Austria). Information on spring and autumn migration, with some data on age, sex, weight, and bill length.
- Miller Mundy, A. 1995. Aggressive behaviour between Common Greenshanks and Common Redshanks. *Brit. Birds* 88(6): 296. (Manish, Isle of Harris PA85 3EZ, UK). *Tringa nebularia* and *T. totanus*.
- Minton, C. 1995. Unusual feeding mode for Eastern Curlew. *Stilt* 26: 36. (165 Dalgety Rd., Beaumaris, Vic 3193, Australia). *Numenius madagascariensis*, Australia.
- Morrier, A. 1990. Activités diurnes du Pluvier de Wilson (*Charadrius wilsonia*) et du Pluvier semipalmé (*Charadrius semipalmatus*) dans les lagunes de Chacopata, Venezuela. [Diurnal activity of Wilson's Plover (*Charadrius wilsonia*) and Semipalmented Plover (*Charadrius semipalmatus*) in the Chacopata lagoons, Venezuela]. Mémoire de maîtrise, Dept. Sci.

- Biol., Univ. Montréal. In French. (Dept. Sci. Biol., Univ. Montreal, C.P. 6128, Succ. Centre-Ville, Montreal, PQ, Canada H3C 3J7).
- Oosterveld, E. 1995. Man en vrouw Tureluur *Tringa totanus totanus* zijn gedurende de broedtijd in het veld te herkennen. (Male and female Redshanks *Tringa t. totanus* are distinguishable in the field). *Limosa* 68(2): 78-80. In Dutch with English summary. (Topweer 7, NL-9865 VD, Opende, Netherlands). Information also on *Limosa limosa*.
- Owens, I.P.F., Dixon, A., Burke, T. & Thompson, D.B.A. 1995. Strategic paternity assurance in the sex-role reversed Eurasian dotterel (*Charadrius morinellus*): behavioral and genetic evidence. *Behav. Ecol.* 6: 14-21. (Dept. Zool., Univ. Leicester, Leicester LE1 7RH, UK). Aims were to test for the existence of mate guarding behaviour predicted by operational sex-ratio theory; to test whether the pattern of copulations in the Dotterel is consistent with that predicted by the sexual conflict theory for a sex-role reversed species; and to assess the realized incidence of extrapair paternity using DNA fingerprinting.
- Paton, P.W.C., Kneedy, C. & Sorensen, E. 1992. Chronology of shorebirds and ibis use of selected marshes at Great Salt Lake. *Utah Birds* 8(1): 1-20. (Co-op. Fish & Wildl. Res. Unit, Dept. Fish & Wildl., Utah State Univ., Logan 84322, USA).
- Piersma, T., Gils, J. van, Goeij, P. de & Meer, J. van der 1995. Holling's functional response model as a tool to link the food-finding mechanism of a probing shorebird with its spatial distribution. *J. Anim. Ecol.* 64: 493-504. (Neth. Inst. Sea Res., P.O. Box 59, NL-1790 AB Den Burg, Texel, Netherlands). A test of whether the intake rate of *Calidris canutus* searching for buried molluscs can be adequately described by a functional response model (disc equation of Holling) assuming prey detection by direct touch.
- Piersma, T., Cadée, N. & Daan, S. 1995. Seasonality in basal metabolic rate and thermal conductance in a long-distance migrant shorebird, the knot (*Calidris canutus*). *J. Comp. Physiol. B* 165: 37-45. Investigations on whether changes in morphology and physiology are a response to the natural annual cycle or based on a seasonal programme exhibited also in captivity.
- Ramenofsky, M., Piersma, T. & Jukema, J. 1995. Plasma corticosterone in Bar-tailed Godwits at a major stop-over site during spring migration. *Condor* 97: 580-585. (Dept. Zool., NJ-15, Univ. Washington, Seattle, WA 98195, USA). Study investigated the regulatory role of corticosterone during stop-over in a long-distance migratory shorebird (*Limosa lapponica*).
- Rogers, K.G. 1995. Eastern Curlew biometrics: based on bivariate separation of the sexes. *The Stilt* 26: 23-34. (340 Ninks Rd., St. Andrews, Vic 3761, Australia). Data on *Numenius madagascariensis* are used to illustrate the method using culmen length and wing length for the bivariate separation and estimating head-bill lengths and weight from them. Weight estimates for adults are given separately for each sex by month up to the time of northward migration.
- Rompré, G. 1993. Etude du comportement diurne et nocturne du Chevalier semipalmé (*Catoptrophorus semipalmatus*) et autres limicoles dans le nord-est du Venezuela. [Study of diurnal and nocturnal behaviour of the Willet (*Catoptrophorus semipalmatus*) and other waders in northeastern Venezuela]. Mémoire de maîtrise, Univ. Montréal. In French. (Dept. Sci. Biol., Univ. Montreal, C.P. 6128, Succ. Centre-Ville, Montreal, PQ, Canada H3C 3J7).
- Schekkerman, H. & Roomen, M. van (eds.) 1995. *Breeding waders at Pronchishcheva Lake, northeastern Taimir, Siberia, in 1991*. Zeist: WIWO (Foundation Working Group International Wader and Waterfowl Research). WIWO-report 55. 89 pp. (Report can be ordered by paying Dfl. 20, plus Dfl. 15 administration costs, to postal giro account 2 666 009 or to ABN bank account 57.02.16.613 of WIWO, Lindengracht 9, NL-3633 AS Vreeland, Netherlands). Information presented on species composition and densities of breeding waders, habitat preference, chick-rearing period, breeding success, population characteristics, and seasonal abundance of tundra invertebrates. Breeding waders: *Pluvialis fulva*, *P. squatarola*, *Charadrius hiaticula*, *Calidris canutus*, *C. alba*, *C. alpina*, *C. ferruginea*, *C. minuta* and *Arenaria interpres*.
- Schulz, M. 1995. Observations on the Hooded Plover *Thinornis rubricollis* on Kangaroo Island, South Australia. *Stilt* 26: 54-57. (Fac. Resourc. Sci. & Manage., Southern Cross Univ., P.O. Box 157, Lismore, NSW 2480, Australia). Information on group size, habitat, nesting, and nest loss.
- Schuster, S. 1994. Untersuchungen zur Mauser des Grossen Brachvogels (*Numenius arquata*) im Vorarlberger Rheindelta. [Investigations on moult of the Curlew in the Vorarlberger Rheindelta, Austria]. *Egretta* 37(2): 60-70. In German with English summary.
- Solís, J.C. & de Lope, F. 1995. Nest and egg crypsis in the ground-nesting Stone Curlew *Burhinus oedicnemus*. *J. Avian Biol.* 26(2): 135-138. (Estación Biológica de Doñana, C.S.I.C., El Rocío, Almonte, Huelva, Spain).
- Struwe-Juhl, B. 1995. Habitatwahl und Nahrungsökologie von Uferschneepfeilen. Familien *Limosa limosa* am Hohner See, Schleswig-Holstein. (Habitat selection and feeding ecology in families of Black-tailed Godwit *Limosa limosa* in the Hohner See area, Schleswig-Holstein). *Vogelwelt* 116(2): 61-72. In German with English summary. (Staatliche Vogelschutzwarte Schleswig-Holstein, Olshausenstr. 40, D-24118 Kiel, Germany).
- Summers, R.W. 1995. Diurnal and tidal activity patterns of Purple Sandpipers on the Isle of May, Fife. *Scott. Birds* 18(1): 51-54. (Lismore, Mill Crescent, North Kessock, Inverness IV1 1XY, UK). *Calidris ferruginea*.
- Thibault, M. 1993. Etude du comportement diurne et nocturne du Pluvier de Wilson (*Charadrius wilsonia*) dans le nord-est du Venezuela. [Study of diurnal and nocturnal behaviour of Wilson's Plover (*Charadrius wilsonia*) in northeastern Venezuela]. Mémoire de maîtrise, Univ. Montréal. In French. (Dept. Sci. Biol., Univ. Montreal, C.P. 6128, Succ. Centre-Ville, Montreal, PQ, Canada H3C 3J7).
- Thorne, D. & Thorne, M. 1994. Unusual Oystercatcher nesting behaviour. *Scott. Bird News* 35: 9. *Haematopus ostralegus*.
- Tinarelli, R. 1993. [Some aspects of the wintering ecology of the Black-winged Stilt, *Himantopus himantopus*, in the inner delta of Niger (Mali)]. *Riv. Ital. Ornitol.* 63(1): 3-9. In Italian with English summary. (Via Savena Inferiore 9, 40061 Minerbio (Bologna), Italy).
- Uhlig, R. 1991 [On the migration of the Grey Plover, *Pluvialis squatarola*, in Bulgaria]. *Zool. Middle East* 5: 21-24. In German with English summary. (Revaler Str. 10, D-10245 Berlin, Germany).
- Verboven, N. & Piersma, T. 1995. Is the evaporative water loss of Knot *Calidris canutus* higher in tropical than in temperate climates?. *Ibis* 137: 308-316. (Neth. Inst. Sea Res., P.O. Box 59, NL-1790 AB Den Burg, Texel, Netherlands).
- Warnock, N., Page, G.W. & Stenzel, L.E. 1995. Non-migratory movements of Dunlins on their California wintering grounds. *Wilson Bull.* 107(1): 131-139. (Ecol., Evol. & Conserv. Biol./186, Univ. Nev., Reno 89512-0013, USA). *Calidris alpina*.

DISTRIBUTION & FAUNISTICS

- Ahmad, M.F. 1994. The wildlife of the Indus River and allied areas. *Records Zool. Sur. Pakistan* 12: 1-38. Resident waders: *Hydrophasianus chirurgus*, *Vanellus indicus*, *Charadrius dubius*, *C. alexandrinus*, *Rostratula benghalensis*, *Himantopus himantopus*, *Ibidorhyncha struthersii*, *Burhinus oedicnemus indicus*, *Glareola pratincola* and *G. lactea*.

Appleby, G., Farnes, R. & Griffith, M. 1995.

- A Grey Phalarope *Phalaropus fulicaria* at Port Fairy, Victoria. *Aust. Bird Watcher* 16(1): 34-38. (95 Broadway, Camberwell, Vic 3124, Australia).
- Argabrite, W.L. 1994. Black-necked Stilt and Glossy Ibis: two new West Virginia records in one week. *Redstart* 61(2): 57-59. (111 Jefferson Park Dr., Huntington, WV 25705, USA). *Himantopus mexicanus*, USA.
- Ashman, P. 1994. Piping Plover (*Charadrius melanotos*) 1 May 1994, Dane County, Nine Springs sewage ponds. *Passenger Pigeon* 56(4): 290. (615 E. Johnson St., Madison, WI 53703, USA).
- Austin-Smith, P.J., Flemming, S.P., Drysdale, C. & Williams, R.G. 1994. The 1991 Piping Plover census in Nova Scotia. *Can. Wildl. Serv. Occas. Pap.* No. 82: 11-15. (P.O. Box 294, Wolfville, NS, Canada B0P 1X0). *Charadrius melanotos*.
- Azam, M.H. 1994. The birds of the Salt range. *Records Zool. Sur. Pakistan* 12: 63-97. (Zool. Surv. Dept., Block 61, Pakistan Secr., Shahrah-e-Iraq, Karachi-74200, Pakistan). Information on status, abundance, and habitat of 28 wader species in Punjab, Pakistan.
- Baccetti, N. 1995. Revisione delle catture di una specie giunta all'orlo dell'estinzione: *Numenius tenuirostris* (Aves, Scolopacidae). *Ric. Biol. Selvaggina* No. 94. 18 pp. In Italian. (Inst. Nazionale Fauna Selvatica, Via Fornacetta 9, Ozzano dell'Emilia, Bologna, Italy). Revision of 141 skins collected between 1828-1974 in Italy.
- Baker, N.E. 1994. The Spur-winged Plover *Vanellus spinosus* in Tanzania. *Scopus* 18(2): 130-131.
- Balevicius, K. (ed.) 1992. *Red Data Book of Lithuania. Rare and endangered species of animals, plants and fungi*. Vilnius: Environmental Protect. Dept. Republ. Lithuania. In Lithuanian with English summary. Information on breeding numbers of *Pluvialis apricaria*, *Haematopus ostralegus*, *Tringa glareola*, *T. totanus*, *Philomachus pugnax*, *Calidris alpina*, *Gallinago media*, *Numenius arquata* and *Limosa limosa* in Lithuania.
- Baszanowski, P., Sikora, A. & Chylarecki, P. 1993. [First breeding record of the Kentish Plover (*Charadrius alexandrinus*) in Poland]. *Notatki Ornitol.* 34(3-4): 376-378. In Polish with English summary.
- Blom, R. 1991. First record of Curlew Sandpiper (*Calidris ferruginea*) in Maryland. *Md. Birdlife* 47(1): 3-4.
- Böhm, K. & Woike, M. 1995. Terekwasseraufer (*Xenus cinereus*) im Rheinland. (Terek Sandpiper in the Rhineland). *Charadrius* 31(1): 63. In German. (Spakeler Str. 57c, D-48159 Münster, Germany).
- Bukacinski, D., Cygan, J.P., Keller, M., Piotrowska, M. & Wojciak, J. 1994. (Numbers and distribution of the waterfowl nesting in the Vistula middle course - fluctuations in 1973-1993). *Notatki Ornitol.* 35(1-2): 5-47. In Polish with English summary. (Inst. Ekologii PAN, Dziekanow Lesny, 05-092 Lomianki, Poland).
- Haematopus ostralegus*, *Charadrius dubius*, *C. hiaticula*, *Vanellus vanellus*, *Limosa limosa*, *Numenius arquata*, *Tringa totanus* and *Actitis hypoleucos*.
- Busche, G. 1995. Bestandsentwicklungen von Brutvögeln in Marschen (Agrarland, Salzwiesen) des westlichen Schleswig-Holstein 1960-1994. (Population dynamics of breeding birds in cultivated coastal lowlands and saltmarshes in western Schleswig-Holstein, 1960-1994). *Vogelwelt* 116(2): 73-90. In German with English summary. (Hochfelder Weg 49, D-25746 Heide, Germany). *Haematopus ostralegus*, *Vanellus vanellus*, *Charadrius hiaticula*, *C. dubius*, *C. alexandrinus*, *Arenaria interpres*, *Gallinago gallinago*, *Numenius arquata*, *Limosa limosa*, *Tringa totanus*, *Calidris alpina*, *Philomachus pugnax* and *Recurvirostra avosetta*.
- Chandler, R.J. 1995. Photospot 36. Asian Dowitcher *Limnodromus semipalmatus*. *Brit. Birds* 88(8): 362-364. (2 Rusland Ave., Orpington, Kent BR6 8AU, UK). On identification of *Limnodromus semipalmatus*.
- Colwell, M.A., Gerstenberg, R.H., Williams, O.E. & Dodd, M.G. 1995. Four Marbled Godwits exceed the North American longevity record for Scolopacids. *J. Field Ornithol.* 66(2): 181-183. (Dept. Wildl., Humboldt State Univ., Arcata, CA 95521, USA). *Limosa fedoa*, California, USA.
- Cook, T., O'Dowd, B. & Durdin, C. 1994. Breeding Redshanks on Essex saltmarshes in 1993. *Essex Bird Report* 1993: 125-148. About 1,000 breeding pairs of *Tringa totanus* in 1993.
- Dann, P. 1994. The distribution and abundance of Palearctic and Australasian waders (Charadrii) in coastal Victoria. *Corella* 18: 148-154. (Penguin Reserve Committee of Manage., P.O. Box 97, Cowes, Phillip Island, Vic 3922, Australia).
- Darolova, A. 1993. [An unusual record of breeding of Pied Avocet (*Recurvirostra avosetta* L., 1758) in the Donau lowland (Podunajska Rovina)]. *Tichodroma* 5: 123-124. In Slovak with German summary. (Ustav Zool. & Ekosozol. SAV, Konvalinkova 3, 842 23 Bratislava, Slovakia).
- David, P.G. 1994. Wading bird nesting at Lake Okeechobee, Florida: an historic perspective. *Colon. Waterbirds* 17(1): 69-77. (Dept. Res., S. Fla. Water Manage. Dist., P.O. Box 24680, West Palm Beach 33416, USA).
- Davis, A. 1994. Status, distribution, and population trends of the New Zealand Shore Plover *Thinornis novaeseelandiae*. *Notornis* 41(Suppl.): 179-194. (Dept. Zool., Univ. Auckland, Private Bag, Auckland, New Zealand).
- Dennis, M. 1993. Essex breeding birds 1992. *Essex Bird Report* 1992: 20-34. (173 Collier Row Lane, Romford, Essex RM5 3ED, UK). *Haematopus ostralegus*, *Vanellus vanellus*, *Gallinago gallinago*, *Scolopax rusticola*, *Limosa limosa* and *Tringa totanus*. Information on number of breeding pairs, Essex, United Kingdom.
- Dennis, M. 1994. Essex breeding birds 1993. *Essex Bird Report* 1993: 20-25. *Recurvirostra avosetta*, *Haematopus ostralegus*, *Burhinus oedicnemus*, *Charadrius dubius*, *C. hiaticula*, *Vanellus vanellus*, *Gallinago gallinago*, *Scolopax rusticola* and *Tringa totanus*.
- Dionisi, V. 1993. [Nidification of the Kentish Plover, *Charadrius alexandrinus*, in Marche (central Italy)]. *Riv. Ital. Ornitol.* 63(1): 77-78. In Italian. (Cent. Studi Ecol., Assoc. Nat. Argonauta, Via L. Severi 25, 61032 Fano (Pesaro), Italy).
- Dombrowski, A., Nawrocki, P., Krogulec, J., Chmielewski, S. & Rzepala, M. 1994. (Breeding avifauna of the lateral arms of the Vistula middle course). *Notatki Ornitol.* 35(1-2): 49-78. In Polish with English summary. (Katedra Ochrony Srodowiska i Ekologii, Wydzsa Szkoła Rolniczo-Pedagogiczna, ul. Prusa 12, 08-110 Siedlce, Poland). *Haematopus ostralegus*, *Limosa limosa*, *Numenius arquata*, *Tringa totanus*, *T. nebularia*, *T. ochropus*, *Charadrius dubius*, *C. hiaticula*, *Vanellus vanellus*, *Calidris alpina*, *Philomachus pugnax*, *Gallinago gallinago* and *Actitis hypoleucos*.
- Dombrowski, A. 1994. (Comparison between the numbers of chosen bird species breeding in the lower Narew valley in 1973 and 1989). *Notatki Ornitol.* 35(3-4): 245-259. In Polish with English summary. Distinct decline in numbers of *Charadrius hiaticula* and *C. dubius*.
- Driscoll, P. 1995. Sightings of waders leg-flagged in Moreton Bay, south east Queensland. *Stilt* 26: 40-41. (Fahey Rd., Mount Glorious, Qld 4520, Australia). *Limosa lepironica*, *Calidris canutus*, *C. tenuirostris* and *Numenius madagascariensis*.
- Ferrand, Y. 1994. Rapport Becasse - coule 1994. (Report on Woodcock (France) - roding season 1994). *IWRB Woodcock and Snipe Res. Group Newsletter* 20: 7-11. In French with English summary. (O.N.C., 5 rue Thibault, St. Benoit F-78610 Auffargis, France).

- Figueroa, J. & Martí, J. 1994. Autumn migration of the Ringed Plover *Charadrius hiaticula* in North-east Iberia. *Butl. GCA* 11: 31-37. (Passeig. Sant Gervasi 59, 7e 2a. 08022 Barcelona, Spain). Main autumn passage between early August and late September. Two periods of maximum passage at the Ebro Delta but only one at the Aiguamolls de l'Emporda Natural Park.
- Flade, M. & Jahn, O. 1994. (Spring record of Buff-breasted Sandpiper (*Tryngites subruficollis*) in northeastern Poland). *Notatki Ornitol.* 35(1-2): 180-182. In Polish with English summary.
- Fleet, D.M., Frikke, J., Südbeck, P. & Vogel, R.L. 1994. Breeding birds in the Wadden Sea 1991. *Wadden Sea Ecosystem No. 1.* Wilhelmshaven: Common Wadden Sea Secr., Triilateral Monitoring & Assessment Group. 108 pp. (Common Wadden Sea Secr., Virchowstr. 1, D-2940 Wilhelmshaven, Germany). Information on distribution, habitat, population size and development, threats and protection of *Haematopus ostralegus*, *Recurvirostra avosetta*, *Charadrius dubius*, *C. hiaticula*, *C. alexandrinus*, *Vanellus vanellus*, *Calidris alpina*, *Philomachus pugnax*, *Gallinago gallinago*, *Limosa limosa*, *Numenius arquata*, *Tringa totanus* and *Arenaria interpres*.
- Frich, A.S. & Nordbjærg, L. 1994. Sjældne fugle i Danmark og Grønland i 1992. (Rare birds in Denmark and Greenland in 1992). *Dansk. Orn. Foren. Tidsskr.* 88: 99-100. In Danish with English summary. (Nyvej 63, 2. tv, DK-2620 Albertslund, Denmark). Waders: *Himantopus himantopus*, *Glareola pratincola*, *Pluvialis fulva*, *Calidris mauri*, *C. melanotos*, *Limosa limosa*, *Gallinago media* and *Xenus cinereus* in Denmark; and *Pluvialis apricaria* in Greenland.
- Ghalib, S.A. & Hanain, S.A. 1994. The waterfowl of the Karachi coast. *Records Zool. Surv. Pakistan* 12: 39-62. (Zool. Surv. Dept., Block 61, Pakistan Secr., Shahrah-e-Iraq, Karachi-74200, Pakistan). Information about 41 wader species, migratory as well as resident.
- Gibbons, B. 1994. Newborough Warren NNR, Anglesey. *Brit. Wildl.* 6(2): 109-111. Breeding waders: *Numenius arquata*, *Haematopus ostralegus*, *Charadrius hiaticula*, *Vanellus vanellus* and *Tringa totanus*, Anglesey, UK.
- Gibbons, B. 1994. The Lower Derwent Valley, Yorkshire. *Brit. Wildl.* 5(6): 381-383. Breeding waders: *Numenius arquata*, *Gallinago gallinago*, *Tringa totanus*, *Vanellus vanellus*, *Philomachus pugnax* and *Limosa limosa*, Yorkshire, UK.
- Gibbs, G. 1993. Wildfowl and waders of the Blackwater Estuary. *Essex Bird Report* 1992: 138-149. (72 Orchard Piece, Blackmore, Essex CM4 0RZ, UK). About 30,000 waders of 12 species winter.
- Internationally important populations of *Pluvialis squatarola*, *Calidris alpina* and *Limosa limosa*. Others: *Charadrius hiaticula*, *Numenius arquata*, *Tringa totanus*, *Haematopus ostralegus*, *Pluvialis apricaria*, *Vanellus vanellus*, *Limosa lapponica*, *Calidris canutus* and *Arenaria interpres*. Information also on other wader species recorded on passage or in winter.
- Glen, R.M. 1994. Notes and new distribution records of the Quail Plover *Ortyxelos meiffreni*. *Scopus* 18(1): 50.
- Gossmann, F. & Bastat-Lequerré, C. 1994. La Bécasse des bois (*Scolopax rusticola*) rapport baguage - saison 1993/94. (Woodcock ringing report 1993/94). *IWRB Woodcock and Snipe Res. Group Newsletter* 20: 12-23. In French with English summary. (O.N.C., Stn. Pays de Loire, 53 rue Russel, F-44000 Nantes, France).
- Grishanov, G. 1994. Breeding birds of the raised bog Zehlau: a survey. *Acta Ornithol. (Lituanica)* 9-10: 127-129. (Biol. Dept., Kaliningrad State Univ., Universitetskaya 2, 236040 Kaliningrad, Russia). *Pluvialis apricaria*, *Vanellus vanellus*, *Tringa glareola* and *Gallinago gallinago*.
- Groen, N.M. & Zomerdijk, P.J. (eds.) 1994. *Waders and waterbirds along part of the Atlantic coast of Morocco. Autumn 1991 - spring 1992*. Zeist: WIWO (Foundation Working Group International Wader and Waterfowl Research). WIWO-report 47. 102 pp. (The report can be ordered by paying Dfl. 20, plus Dfl. 15 administration costs, to postal giro account 2 666 009 or to ABN bank account 57.02.16.613 of WIWO, Lindengracht 9, NL-3633 AS Vreeland, Netherlands). Results of censuses of wetlands along the Moroccan coast from Tanger to Qued Massa. Data for 24 wader species.
- Haig, S.M. & Plissner, J.H. 1994. The 1991 international Piping Plover census in the United States, Mexico, and the Caribbean: breeding and winter populations. *Can. Wildl. Serv. Occas. Pap.* No. 82: 48-51. (US Fish & Wildl. Serv., Clemson Univ., Clemson, SC 29634). *Charadrius melanotos*.
- Harper, G. 1994. Where does the Slender-billed Curlew nest? New ideas from Russia. *Scott. Bird News* 35: 5. *Numenius tenuirostris*.
- Harris, K. 1995. Population monitoring counts: winter 1994. *Stilt* 26: 21-22. (59 Strickland Drive, Wheelers Hill, Victoria 3150, Australia). Numbers for 51 wader species counted in Australia.
- Hasan, A.-U. 1994. The birds of Sindh mangroves. *Records Zool. Surv. Pakistan* 12: 98-105. (Zool. Surv. Dept., Block 61, Pakistan Secr., Shahrah-e-Iraq, Karachi-74200, Pakistan). Indus delta mangrove area: most abundant waders in winter are
- Tringa totanus*, *Charadrius dubius*, *Calidris minuta*, *Charadrius mongolus* and *Calidris alba*.
- Heard, C.D.R. 1994. First record of Black-winged Pratincole *Glareola nordmanni* in Yemen. *Sandgrouse* 16(1): 57.
- Heyens, L.E. 1994. The 1991 Piping Plover census in Ontario. *Can. Wildl. Serv. Occas. Pap.* No. 82: 30-31. (Ontario Minist. Nat. Resour., P.O. Box 5080, Kenora, ON, Canada P9N 3X9). *Charadrius melanotos*.
- Hirst, P. & Proctor, B. 1995. Identification of Wandering and Grey-tailed Tattlers. *Birding World* 8(3): 91-97. *Heteroscelus incanus* and *H. brevipes*.
- Hofman, E.D. 1994. The 1991 Piping Plover census in Alberta. *Can. Wildl. Serv. Occas. Pap.* No. 82: 43-47. (Alberta Fish & Wildl. Div., #404, 4911-51 St., First Red Deer Pl., Red Deer, AB, Canada T4N 6V4). *Charadrius melanotos*.
- Hoodless, A. 1994. Woodcock - the final report. *Game Conservancy Annual Rev.* 25: 88-90. *Scolopax rusticola*, UK.
- Van Impe, J. 1995. Considérations sur les causes de la disparition du Courlis à bec grêle *Numenius tenuirostris*. (On the causes of disappearance of the Slender-billed Curlew *Numenius tenuirostris*). *Alauda* 63(2): 111-114. In French with English summary. (Dr. Van de Perrelei, 51 B, B-2140 Borgerhout, Belgium). Decline of *Numenius tenuirostris* might be related to disappearance of huge steppe areas e.g. in Kazakhstan.
- Janiszewski, T., Michalak, P. & Bargiel, R. 1993. [Report from ringing of waders at Jeziorsko Reservoir in 1989-1993]. *Notatki Ornitol.* 34(3-4): 392-393. In Polish with English summary. Data on 21 wader species.
- Jessop, R. & Minton, C. 1995. Sightings of leg-flagged waders banded in NW Australia. *Stilt* 26: 37-38. (P.O. Box 97, Cowes, Victoria 3922, Australia). *Charadrius leschenaultii*, *Himantopus himantopus*, *Arenaria interpres*, *Numenius madagascariensis*, *Heteroscelus brevipes*, *Limosa lapponica*, *Calidris canutus*, *C. tenuirostris*, *C. ruficollis*, *C. ferruginea* and *Limicola falcinellus*.
- Jessop, R. & Minton, C. 1995. Sightings of leg-flagged waders banded at other locations and seen in NW Australia. *Stilt* 26: 39. *Arenaria interpres*, *Calidris tenuirostris* and *C. ruficollis*.
- Johnson, A. & Allen, P. 1994. Black-necked Stilt at Riverton area. *Iowa Bird Life* 64(1): 20-21. *Himantopus mexicanus*, Iowa, USA.
- Kane, A. 1993. Greater Sand Plover at E. Tilbury. A first for Essex - 10th to 14th

- August 1992. *Essex Bird Report* 1992: 132-133. *Charadrius leschenaultii*, UK.
- Khanum, Z. & Ahmed, M. 1994. Some more species of birds observed in Karachi during 1987-88. *Records Zool. Sur. Pakistan* 12: 106-108. (Dept. Zool., Univ. Karachi, Karachi-75270, Pakistan). *Vanellus leucurus*, *Charadrius hiaticula*, *Calidris alpina*, *C. tenuirostris* and *Arenaria interpres*.
- Khanum, Z., Farooq, M. & Ahmed, M. 1994. A revised checklist of birds of Pakistan. *Records Zool. Sur. Pakistan* 12: 144-200.
- Kirwan, G. 1994. American Golden Plover, *Pluvialis dominica*, in Turkey. *Zool. Middle East* 10: 23-26. (6 Connaught Rd., Norwich NR2 3BP, UK).
- Knox, K., Brazil, J. & Etchberry, R. 1994. The 1991 Piping Plover census in Newfoundland and St.-Pierre-et-Miquelon. *Can. Wildl. Serv. Occas. Pap.* No. 82: 8-10. (Newfoundland & Labrador Wildl. Div., Dept. Tourism & Cult., P.O. Box 8700, St. John's, NF, Canada A1B 4J6). *Charadrius melanotos*.
- Komisja Faunistyczna Sekcji Ornitologicznej PTZool. 1994. (Rare birds recorded in Poland in 1993. Avifaunistic Commission Report No. 10). *Notatki Ornitol.* 35(3-4): 331-346. In Polish with English summary. (ul. Sienkiewicza 21, 50-335 Wroclaw, Poland). Waders: *Haematopus ostralegus*, *Himantopus himantopus*, *Burhinus oedicnemus*, *Charadrius alexandrinus*, *C. morinellus*, *Calidris bairdii*, *C. melanotos*, *C. maritima*, *Xenus cinereus* and *Phalaropus fulicarius*.
- Koonz, W.H. 1994. The 1991 Piping Plover census in Manitoba. *Can. Wildl. Serv. Occas. Pap.* No. 82: 32-34. (Manitoba Wildl. Branch, Box 14, 1495 St. James St., Winnipeg, MB, Canada R3H 0W9). *Charadrius melanotos*.
- Kurlavicius, P. 1991. Rare and new bird species in Lithuania: this century. *Acta Ornithol. (Lituanica)* 4: 81-95. *Burhinus oedicnemus*, *Charadrius morinellus*, *Xenus cinereus*, *Phalaropus lobatus* and *Calidris maritima*.
- Kuzyakin, V. 1994. An estimate of the breeding population of the Woodcock (*Scolopax rusticola*) in France. *IWRB Woodcock and Snipe Res. Group Newsletter* 20: 24-25. (Inst. Anim. Ecol., Russian Acad. Sci., Leninsky prospect 33, Moscow V - 17, Russia).
- Kuzyakin, V. 1994. Status of the Woodcock in the Far East of Russia. *IWRB Woodcock and Snipe Res. Group Newsletter* 20: 25-26. *Scolopax rusticola*.
- Laporte, P. & Shaffer, F. 1994. The 1991 Piping Plover census in Quebec. *Can. Wildl. Serv. Occas. Pap.* No. 82: 24-29. (Serv. Canadien Faune, Dir. Conserv. Environ., Reg. Quebec, 1141 Rt. Eglise, C.P. 10100, Sainte-Foy, PQ, Canada G1V 4H5). *Charadrius melanotos*.
- Lawler, W. 1994. *Shorebird counts on New South Wales north coast estuaries from the Shorebird Habitat Study*. National Parks & Wildl. Serv., Environment Survey & Res. Branch. (c/o Ecopix, P.O. Box 67, Scarborough, Qld. 4020, Australia).
- Leibak, E., Lilleleht, V. & Veromann, H. (eds.) 1994. *Birds of Estonia. Status, distribution and numbers*. Tallinn: Estonian Academy Publ. 287 pp. ISBN: 9985-50-031-8.
- Lindballe, P., Christensen, R., Munk, M.F., Skov, H., Smidt, J. & Soby, E. 1994. *Fugle i Danmark 1992. Årsrapport over observationer*. (The Danish Bird Report 1992). *Dansk. Orn. Foren. Tidsskr.* 88: 111-150. In Danish with English summary. (Dansk Ornithologisk Forening, Vesterbrogade 140, DK-1620 Copenhagen V, Denmark). Summary of occurrence in Denmark of bird species which generally are not subject to systematic monitoring schemes. Waders: *Charadrius alexandrinus*, *C. morinellus*, *Calidris minuta*, *C. temminckii*, *C. ferruginea*, *C. maritima*, *Limicola falcinellus*, *Gallinago media*, *Phalaropus lobatus* and *P. fulicarius*.
- Malinauskas, V. 1993. Bird species of the Red Data Book of Lithuania detected in the Kaisiadorys. *Acta Ornithol. (Lituanica)* 7-8: 115-116. (Gireles 55-214 Kaisiadorys, Lithuania). *Calidris alpina*, *Limosa limosa*, *Tringa glareola* and *Haematopus ostralegus*.
- Matthes, W. 1994. Limikolen im Raum Worms (Rheinland-Pfalz). (Waders in the Worms (Rhine) area). *Fauna und Flora in Rheinland-Pfalz. Zeitschr. Naturschutz Beiheft* 12. 135 pp. In German with English summary. Statements mainly about migration phenology, biometry and moult in connection with an evaluation of ringing results of Rhineland-Palatinate for 32 wader species, 1965-1980.
- McAskill, D., Hoteling, B., Ristau, C. & Flemming, S.P. 1994. The 1991 Piping Plover census in Prince Edward Island. *Can. Wildl. Serv. Occas. Pap.* No. 82: 16-23. (Nat. Hist. Soc. Prince Edward Island Ltd., P.O. Box 2346, Charlottetown, PE, Canada C1A 8C1). *Charadrius melanotos*.
- McIntyre, A., Hines, H. & Pollock, A. 1995. Colour banding Hooded Plover *Thinornis rubricollis* in East Gippsland, Victoria. *Stilt* 26: 50-53. (Dept. Conserv. & Nat. Resour., P.O. Box 260, Orbost, Victoria 3888, Australia).
- Mearns, R. 1995. Little Ringed Plovers attempt breeding in Dumfriesshire. *Scott. Birds* 18(1): 55. (Connansknowe, Kirkton, Dumfries DG1 1SX, UK). *Charadrius dubius*, Scotland.
- Meininger, P.L., Schekkerman, H. & Roomen, M.W.J. van 1995. *Populationschattingen en 1%-normen van in Nederland voorkomende watervogelsoorten: voorstellen voor standaardisatie*. (Population estimates and 1%-levels for waterbird species occurring in The Netherlands: suggestions for standardisation). *Limosa* 68(2): 41-48. In Dutch with English summary. (Rijksinst. Kust en Zee, RIJKZ, P.O. Box 8039, NL-4330 EA Middelburg, Netherlands). Information on most recently published population estimates for all waterbird species regularly occurring in the Netherlands. Data for 28 wader species.
- Minton, C., Rogers, D. & Weston, M. 1995. Catching up with Australian birds in Siberia. *Geo-Australasia* 17(5): 94-104. (165 Dalgety Rd., Beaumaris, Victoria 3193, Australia). Report of the Australian part of "Tundra Ecology '94" expedition, with information on breeding and migration of arctic waders.
- Moerbeek, D.J. 1994. [Baird's Sandpiper at Julianadorp in July-August 1993]. *Dutch Birding* 16(1): 15-18. In Dutch with English summary. *Calidris bairdii*, Netherlands.
- Morrison, R.I.G., Bourget, A., Butler, R., Dickson, H.L., Gratto-Trevor, C., Hicklin, P., Hyslop, C. & Ross, R.K. 1994. A preliminary assessment of the status of shorebird populations in Canada. *Progress Notes* No. 208. Ottawa: Canadian Wildlife Service. 19 pp. (Can. Wildl. Serv., Environ. Can., Ottawa, Ontario, Canada K1A 0H3). Review of published and unpublished information on the status and abundance of shorebird populations in Canada and on population trends and sizes (73 shorebird species).
- OAG Münster 1994. 15 Jahre "Birding" in den Rieselfeldern Münster - neue Erkenntnisse für die Avifaunistik? (Fifteen years birding at the Münster sewage farms - new discoveries for field ornithology?). *Charadrius* 30(4): 181-185. In German with English summary. (OAG Münster, Biologische Stn., Coermühle 181, D-48157 Münster, Germany). Compilation of observations of 33 rare species for Westphalia (Germany) in the Ramsar site "Rieselfeldern Münster". Waders: *Himantopus himantopus*, *Charadrius alexandrinus*, *Calidris melanotos*, *Limicola falcinellus*, *Micropalama himantopus*, *Limosa limosa islandica*, *Tringa stagnatilis* and *Xenus cinereus*.
- Nobel, W.T. de (ed.) 1995. *Birds of the Messolonghi wetlands. Eastern Mediterranean Wader Project, spring 1990*. Zeist: WIWO (Foundation Working Group International Wader and Waterfowl Research). WIWO-report 53, 144 pp. (Report can be ordered by paying Dfl. 25, plus Dfl. 15 administration costs, to postal giro account 2 666 009 or to ABN bank account 57.02.16.613 of WIWO, Lindengracht 9, NL-3633 AS Vreeland,

- Netherlands). Spring migration includes *Charadrius alexandrinus*, *Calidris alpina*, *Philomachus pugnax*, *Limosa limosa*, *Numenius arquata*, *Tringa totanus*, *Calidris minuta*, *C. ferruginea*, *Tringa erythropus*, *T. glareola*, *Arenaria interpres*. Breeding waders (high numbers): *Himantopus himantopus*, *Burhinus oedicnemus*, *Glareola pratincola*, *Charadrius alexandrinus* and *Tringa totanus*.
- Olsen, K.M. 1992. [Field identification of Marsh Sandpiper]. *Calidris* 21: 10-14. In Swedish. (Gartnerievej 3, DK-2100, Copenhagen O, Denmark). *Tringa stagnatilis*.
- Page, G.W., Stern, M.A. & Paton, P.W.C. 1995. Differences in wintering areas of Snowy Plovers from inland breeding sites in western North America. *Condor* 97(1): 258-262. *Charadrius alexandrinus*.
- Peach, W., Henderson, I. & Baillie, S. 1994. Snipe and Woodcock in Europe: an analysis of European ringing recoveries. Game Conservancy Annual Rev. 25: 91-92. *Scolopax rusticola* and *Gallinago gallinago*.
- Pietrelli, L., Biondi, M. & Guerrieri, G. 1994. Insediamenti di una colonia di Cavaliere d'Italia, *Himantopus himantopus*, nel Lazio. [First record of a breeding colony of Black-winged Stilts, *Himantopus himantopus*, in Latium (central Italy)]. *Riv. Ital. Ornitol.* 63(2): 222-225. In Italian with English summary. (Via M. Fantucci 8, I-00119 Roma, Italy).
- Plesník, J. 1994. [The Curlew (*Numenius arquata*)]. *Ochr. Prirody* 49(7): 208-209. In Czech with English summary.
- Pranašaitis, A. 1992. Avifauna of Zuvintas Nature Reserve, 1980-90. *Acta Ornithol. (Lituanica)* 5-6: 64-71. (Zuvintas Nat. Reserve, 4583 Simnas, Alytus District, Lithuania). With information on breeding, status, abundance, wintering, and migration. Information on 24 wader species.
- Reed, J.M. & Oring, L.W. 1993. Long-term population trends of the endangered Ae'o (Hawaiian Stilt, *Himantopus mexicanus knudseni*). *Trans West. Sect. Wildl. Soc.* 29: 54-60. (Ecol., Evol. & Conserv. Biol. Program, Univ. Nev., 1000 Valley Rd., Reno 89512, USA).
- Reese, J. 1992. Northernmost Chesapeake Bay nest record for Willet. *Md. Birdlife* 48(2): 75. *Catoptrophorus semipalmatus*, Maryland, USA.
- Richter, J. 1990 [Record of *Limosa lapponica* L. and *Calidris ferruginea* Pont. in the northern Upper Lausitz in autumn]. *Beitr. Vogelkd.* 36(2): 126. In German.
- Rösner, H.-U., Roomen, M. van, Südbrock, P. & Rasmussen, L.M. 1994. Migratory Waterbirds in the Wadden Sea 1992/93.
- Wadden Sea Ecosystem No. 2. Wilhelmshaven: Common Wadden Sea Secr., Trilateral Monitoring & Assessment Group. (Common Wadden Sea Secr., Virchowstr. 1, D-2940 Wilhelmshaven, Germany).* Information on distribution of *Haematopus ostralegus*, *Recurvirostra avosetta*, *Charadrius hiaticula*, *Pluvialis apricaria*, *P. squatarola*, *Vanellus vanellus*, *Calidris canutus*, *C. alba*, *C. ferruginea*, *C. alpina*, *Philomachus pugnax*, *Limosa lapponica*, *Numenius phaeopus*, *Numenius arquata*, *Tringa totanus*, *T. guttifer* and *Arenaria interpres* in the Dutch, German and Danish part of the Wadden Sea.
- Rowlands, J.A. 1994. First record of Oriental Pratincole *Glareola maldivarum* in Cyprus. *Sandgrouse* 16(1): 56.
- Sada, A.M. 1994. Additional information of Piping Plovers in Mexico. *Euphonia* 3(1): 21. (Apdo. Postal 1124, Monterrey, Nuevo Leon 64000, Mexico). *Charadrius melanotos*.
- Schulz, M. 1995. Inland record of the Sooty Oystercatcher *Haematopus fuliginosus*. *The Stilt* 26: 42. (Fac. Resourc. Sci. & Manage., Southern Cross Univ., P.O. Box 157, Lismore, NSW 2480, Australia).
- Schulz, M. 1995. A survey of shorebirds of Kangaroo Island, South Australia. *The Stilt* 26: 46-49. With information on 25 wader species.
- Shackford, J.S. 1994. Nesting of Long-billed Curlews on cultivated fields. *Bull. Okla. Ornithol. Soc.* 27(3): 17-20. (Co-op. Fish & Wildl. Res. Unit., Dept. Zool., Okla. State Univ., Stillwater 74078, USA). *Numenius americanus*, Oklahoma, USA.
- Shane, T.G. & Shane, S.J. 1995. Mountain Plover nesting in Otero County, Colorado. *Colorado Field Ornithol. J.* 29(1): 15. (P.O. Box 876, Garden City, KS 67846, USA). *Charadrius montanus*, USA.
- Skeel, M.A. 1994. The 1991 Piping Plover census in Saskatchewan. *Can. Wildl. Serv. Occas. Pap.* No. 82: 35-42. (Saskatchewan Nat. Hist. Soc., P.O. Box 4348, Regina, SK, Canada S4P 3W6). *Charadrius melanotos*.
- Stillman, R.A. & Brown, A.F. 1994. Sampling strategies for estimating the size of upland breeding bird populations. *JNCC Report No. 188*. Peterborough: Joint Nature Conservation Committee. 22 pp. (JNCC, Monkstone House, City Rd., Peterborough PE1 1UA, UK). Random sampling could have produced large reductions in survey effort for *Numenius arquata* and *Pluvialis apricaria*. Reductions in survey effort of over 25% in comparison with random sampling were obtained for *Calidris alpina* using stratified sampling by percent moorland.
- Thorpe, R.I. 1995. Grey-tailed Tattler in Wales: new to Britain and Ireland. *Brit. Birds* 88(6): 255-262. (2 Tan y Garth, Friog, Fairbourne, Gwynedd LL38 2RJ, UK). *Heteroscelus brevipes*.
- Tomkovich, P.S. 1995. Great Knot on Siberian breeding grounds. *Dutch Birding* 17(1): 15-16. (Zool. Mus., Moscow State Univ., Herzen Str. 6, 103009 Moscow K-9, Russia). *Calidris tenuirostris*.
- Uhlig, R. 1992. [*Haematopus ostralegus* in southwestern Poland in June]. *Beitr. Vogelkd.* 38(5/6): 359-360. In German. (Revaler Str. 10, D-10245 Berlin, Germany).
- Vaitkus, G., Vaitkuviene, D., Zalakevicius, M. & Svazas, S. 1994. Characteristics of summer bird migration in the Lithuanian coast of the Baltic Sea. *Acta Ornithol. (Lituanica)* 9-10: 38-51. (Inst. Ecol., Akademijos 2, 2600 Vilnius, Lithuania). Waders: *Pluvialis squatarola*, *P. apricaria*, *Vanellus vanellus*, *Charadrius hiaticula*, *C. morinellus*, *Arenaria interpres*, *Haematopus ostralegus*, *Tringa glareola*, *T. nebularia*, *T. totanus*, *Actitis hypoleucos*, *Philomachus pugnax*, *Calidris minuta*, *C. temminckii*, *C. alpina*, *C. canutus*, *C. alba*, *Gallinago gallinago*, *Scolopax rusticola*, *Numenius arquata*, *N. phaeopus* and *Limosa limosa*.
- Weston, M. & Heislers, D. 1995. An estimate of the proportion of Pied Oystercatcher pairs overwintering on open ocean beaches in eastern Victoria, Australia. *The Stilt* 26: 43-45. (28 Craig Rd., Donvale, Vic 3111, Australia). *Haematopus longirostris*.
- Wilhelm, G. 1995. Scenario of the Upland Sandpiper in western Pennsylvania. *Pa. Birds* 8(4): 204-206. (147 Cemetery Rd., Slippery Rock, PA 16057, USA). *Bartramia longicauda*, USA.
- Yovanovich, G.D.L. 1995. Collared Plover in Uvalde, Texas. *Birding* 27(2): 102-104. (48 Countryside Dr., Basking Ridge, NJ 07920, USA). *Charadrius collaris*, USA.
- Zalakevicius, M., Raudonikis, L., Grazulevicius, G. & Idzelis, R. 1994. Phenology of birds arrival to eastern Europe. *Acta Ornithol. (Lituanica)* 9-10: 12-29. (Lab. Ornithol., Inst. Ecol., Akademijos 2, 2600 Vilnius, Lithuania). Initial spring arrivals of birds at various sites in Lithuania, Estonia and Ukraine was investigated using cluster analysis, and appropriate models constructed. Waders: *Vanellus vanellus*, *Gallinago gallinago*, *Scolopax rusticola*, *Tringa ochropus*, *T. totanus*, *Haematopus ostralegus*, *Numenius arquata* and *Tringa glareola*.
- Zalakevicius, M. (comp.) 1995. Birds of Lithuania. Status, number, distribution (breeding, migration, wintering). *Acta Ornithol. (Lituanica)* 11(special issue): 105 pp. In Lithuanian and English. Information on 38 wader species in Lithuania.

MISCELLANEOUS

Anon. 1995. The Kushiro Initiative. Summary statement from the international workshop on conservation of migratory waterbirds and their wetland habitats in the East Asian Australasian Flyway: 2 December 1994, Kushiro, Japan. *Stilt* 26: 9-11. With a framework for migratory waterbird conservation in the East Asia/ Australasian flyway, and a format for an East Asian/Australasian migratory shorebird action plan.

Arnold, K.A. 1994. Common Snipe. In: Tacha, T.C. & Braun, C.E. (eds.). *Migratory shore and upland game bird management in North America*. Pp. 117-125. (Dept. Wildl. & Fish. Sci., Tex. A&M Univ., College Stn. 77843-2258, USA). *Gallinago gallinago*.

Burger, J., Seyboldt, S., Morganstein, N. & Clark, K. 1993. Heavy metals and selenium in feathers of three shorebird species from Delaware Bay. *Environ. Monit. Assess.* 28(2): 189-198. (Dept. Biol. Sci., Rutgers Univ., Piscataway, NJ 08855, USA). *Calidris alba*, *C. canutus* and *C. pusilla*, New Jersey and Delaware, USA.

Chu, P.C. 1995. Phylogenetic reanalysis of Strauch's osteological data set for the Charadriiformes. *Condor* 97(1): 174-196.

Elmes, G.W. & Free, A. (eds.) 1994. Climate change and rare species in Britain. *ITE Research Publ.* No. 8. London, HMSO. ISBN: 0-11-701802-3. Rare breeding waders: *Calidris maritima*, *C. temminckii*, *Recurvirostra avosetta*, *Burhinus oedicnemus*, *Limosa limosa*, *Tringa glareola* and *Phalaropus lobatus*.

Fair, J.M., McEwen, L.C. & Kennedy, P.L. 1994. Organochlorine residues in nesting Killdeer, *Charadrius vociferus*, and their eggs in North Dakota. *Can. Field-Nat.* 108(3): 328-331.

Hüppop, O. & Hüppop, K. 1995. Der Einfluss von Landwirtschaft und Wegenutzung auf die Verteilung von Küstenvogel-Nestern auf Salzwiesen der Hallig Nordstrandischmoor (Schleswig-Holstein). (The influence of agriculture and road usage on the distribution of coastal bird nests on saltmarshes in Schleswig-Holstein (Germany). *Vogelwarte* 38(2): 76-88. In German with English summary. (Inselstatn. Helgoland Inst. Vogelforschung 'Vogelwarte Helgoland', P.O. Box 1220, D-27494 Helgoland, Germany). *Haematopus ostralegus*.

Koenen, M.T. & Leslie, D.M. Jr. 1994. Managing for Least Terns and Snowy Plovers at Salt Plains National Wildlife Refuge. *Scissortail* 44(4): 40. (Co-op. Fish & Wildl. Res. Unit, Dept. Zool., Okla. State Univ., Stillwater 74078, USA). *Charadrius alexandrinus*, Oklahoma, USA.

Prellwitz, D.M., Erickson, K.M. & Osborne,

L.M. 1995. Translocation of Piping Plover nests to prevent nest flooding. *Wildl. Soc. Bull.* 23(1): 103-106. (US Fish & Wildl. Serv., Bowdoin Natl. Wildl. Refuge, HC 65, Box 5700, Malta, MT 59538, USA). *Charadrius melanotos*, USA.

Schweitzer, S.H. 1994. *Conservation measures for the interior population of Least Terns nesting at Salt Plains National Wildlife Refuge*. Ph.D. dissertation, Oklahoma State Univ. 117 pp. *Charadrius alexandrinus*, Oklahoma, USA.

Stillman, R.A. & Brown, A.F. 1995. Minimising effort in large-scale surveys of terrestrial birds: an example from the English uplands. *J. Avian Biol.* 26(2): 124-134. (Inst. Terr. Ecol., Furzebrook Res. Stn., Wareham, Dorset BH20 5AS, UK). Evaluation of three techniques with reference to estimating the size of upland bird populations in the south Pennines of England. Waders: *Numenius arquata*, *Calidris alpina*, *Pluvialis apricaria*, *Vanellus vanellus* and *Gallinago gallinago*.

Straw, J.A. Jr., Krementz, D.G., Olinde, M.W. & Sepik, G.F. 1994. American Woodcock. In: Tacha, T.C. & Braun, C.E. (eds.). *Migratory shore and upland game bird management in North America*. Pp. 97-114. (Krementz: Southeast Res. Group, Natl. Biol. Serv., Warnell School For. Resour., Univ. Georgia, GA, Athens 30602-2152, USA). *Scolopax minor*.

Tubbs, C.R. 1995. Sea level change and estuaries. *Brit. Wildl.* 6(3): 168-176. Results of a study in The Solent, UK. Implications for estuary conservation are discussed.

Watkins, D. 1995. The "Kushiro Initiative" - developing an international shorebird conservation program. *Stilt* 26: 7-8. On shorebird conservation in the East Asia/ Australasian flyway.

AVEIRO MEETING CONTINUED...

One must recognise with happiness that every one of these meetings has more participants and every time there are even more people from an ever wider range of countries. We hope it will continue like this and that the people from The Netherlands and Spain (the more indefatigable) continue to attend with desire to liven up the nights.

Francisco Hortas

Thank you!

Many thanks to all those who have helped recently with all stages of recent *Bulletin* publication:

EDITORIAL

Nigel Clark, Nick Davidson, Julianne Evans, John Goss-Custard, Tom van der Have, Phil Hockey, Phil Holland, John Holmes, Henk Koffijberg, Lys Muirhead, Callum Rankine, Peter Reay, Greg Ruiz, Robin Ward, Sylvia Way and Derek Yalden.

PRODUCTION AND

DISTRIBUTION

Reg Davies and Rodney West.

ILLUSTRATIONS

D. Beadle, Jens Gregersen, Eugeny A. Koblik, Achim Strache and Rolf-Rüdiger Strache.

And apologies to anyone inadvertently omitted!

DATES FOR YOUR DIARY!

FUTURE WSG MEETINGS

1996

The 1996 WSG meeting will be held in early October in Belgium and organised by the Flemish Institute for Nature Conservation and the Flemish Ornithological Society. Details of arrangements will be announced in the April 1996 *WSG Bulletin*.

1997

The dates for the 1997 WSG annual meeting will be 8-12 August in Denmark.