

RECENT PUBLICATIONS ON WADERS 68

COMPILED BY
HENK KOFFIJBERG

In trying to make these lists as complete as possible, and the next one even more complete and up to date, your help is most welcome. Please send new titles, abstracts, reprints, copies, reports, and (**important tool!**) omissions or corrections to: Henk Koffijberg, Glenn Millerweg 102, 1311 RW Almere, The Netherlands.

Remember to include, if possible, the address of the first author, and translations of titles if they are in other languages than English.

For help with this issue I'm very grateful to D.A. Stroud, I. Bainbridge, L. Slotta-Bachmayr and G. Oliso.

Translated titles from languages other than English are bracketed; with round brackets if the translation is by the author(s), square brackets if not.

N.B. a copy of your paper or abstract would be very helpful to enable me to add some keywords or give added annotation to the titles.

So far, about 6 000 titles have been stored in the WSG/RPW database. Keep sending more!!

BREEDING

Exo, K.-M., Becker, P.H., Hälterlein, B., Hötker, H., Scheuffler, H., Stiefel, A., Stock, M., Südbek, P. & Thorup, O. 1996. Bruterfolgsmonitoring bei Küstenvögeln. (Monitoring breeding success of coastal birds). *Vogelwelt* 117(4-6): 287-293. In German with English summary. (Inst. Vogelforschung, An der Vogelwarte 21, D-26386 Wilhelmshaven, Germany). Waders: *Haematopus ostralegus*, *Recurvirostra avosetta* and *Tringa totanus*.

Fraga, R.M. & Amat, J.A. 1996. Biología de la reproducción de una población de Chorlitejos Patinegros (*Charadrius alexandrinus*) en una laguna salada interior. (Breeding biology of a Kentish Plover (*Charadrius alexandrinus*) population in an inland saline lake). *Ardeola* 43(1): 69-86. In Spanish with English summary.

Gregory, R. & Bashford, R. 1996. Breeding Bird Survey: 1994-1995 index report. *BTO News* 206: 5-8. (BTO, The Nunnery, Thetford, Norfolk IP24 2PU, UK). Waders: *Haematopus ostralegus*, *Pluvialis apricaria*, *Vanellus vanellus*, *Gallinago gallinago*, *Numenius arquata*, *Tringa totanus* and *Actitis hypoleucos*.

Hälterlein, B. & Südbek, P. 1996. Brutbestands-Monitoring von Küstenvögeln an der deutschen Nordseeküste. (Monitoring of breeding seabirds and coastal birds along the German North Sea coast). *Vogelwelt* 117(4-6): 277-285. In German with English summary. (Landesamt für den Nationalpark Schleswig-Holsteinisches Wattenmeer, Schlossgarten 1, D-25832 Tönning, Germany). *Haematopus ostralegus* and *Recurvirostra avosetta* showed the most marked increase. Information on breeding numbers and trends of *Charadrius dubius*, *C. hiaticula*, *C. alexandrinus*, *Vanellus vanellus*, *Calidris alpina*, *Philomachus pugnax*, *Gallinago gallinago*, *Limosa limosa*, *Numenius arquata*, *Tringa totanus* and *Arenaria interpres*.

Harrison, F. 1996. An unusual nest of the Comb-crested Jacana *Irediparra gallinacea*. *Stilt* 29: 37. (4/6 Albert St., Cranbrook, Qld. 4814, Australia).

Kálás, J.A., Fiske, P. & Saether, S.A. 1995. The effect of mating probability on risk taking: an experimental study in lekking Great Snipe. *Am. Nat.* 146(1): 59-71. *Gallinago media*.

Kruk, M., Noordervliet, M.A.W. & Ter Keurs, W.J. 1996. Hatching dates of waders and mowing dates in intensively exploited grassland areas in different years. *Biol. Conserv.* 77(2-3): 213-218. (Dept. Environm. Biol., Leiden Univ., P.O. Box 9516, NL-2300 RA Leiden, Netherlands).

Kuczynski, L. 1994. [Geographical variation of egg size in Lapwing *Vanellus vanellus*]. *Przegl. Przyr.* 5(2): 46-56. In Polish with English summary.

Loefering, J.P. & Fraser, J.D. 1995. Factors affecting Piping Plover chick survival in different brood-rearing habitats. *J. Wildl. Manage.* 59: 646-655. *Charadrius melodus*.

Mädlow, W. & Mayr, C. 1996. Die Bestandsentwicklung ausgewählter gefährdeter Vogelarten in Deutschland - 1990-1994. (Population trends of some scarce breeding birds in Germany 1990-1994). *Vogelwelt* 117(4-6): 249-260. In German with English summary. (Am

Stadtpark 4, D-16303 Schwedt, Germany). Waders: *Limosa limosa*, *Numenius arquata* and *Tringa totanus*.

Marchant, J. & Wilson, A. 1996. Waterways Bird Survey: 1994-95 population changes. *BTO News* 205: 12-14. (BTO, The Nunnery, Thetford, Norfolk IP24 2PU, UK). Waders: *Haematopus ostralegus*, *Vanellus vanellus*, *Numenius arquata*, *Tringa totanus* and *Actitis hypoleucos*.

Noszály, G., Székely, T. & Hutchinson, J.M.C. 1995. Brood survival of Kentish Plovers (*Charadrius alexandrinus*) in alkaline grasslands and drained fish-ponds. *Ornis Hung.* 5: 15-21. (Behav. Ecol. Res. Grp., Dept. Zool., Kossuth Univ., Pf. 3, Debrecen, H-4010, Hungary).

Opermanis, O., Strazds, M., Peterhofs, E. & Strazds, A. 1996. (Recent changes, distribution and present status of birds on Latvian seacoast). *Putni daba* 6(1): 2-14. In Latvian with English summary. (Mus. Zool., Univ. Latvia, Kronvalda bulv. 4, Riga LV-1842, Latvia). Waders: *Haematopus ostralegus*, *Charadrius dubius*, *C. hiaticula*.

Ottens, H.J. & Manen, W. van 1995. Uitkomstsucces van Kievit *Vanellus vanellus* en Scholekster *Haematopus ostralegus* in Loon-West. (Nest survival in Lapwing *Vanellus vanellus* and Oystercatcher *Haematopus ostralegus* near Loon-West). *Drentse Vogels* 8: 21-25. In Dutch with English summary. (Groningerweg 34, NL-9738 AB Groningen, Netherlands). Nest survival in farmland area in central Drenthe, Netherlands.

Slotta-Bachmayr, L. 1996. Bestandsentwicklung und Habitatwahl wiesenbrütender Vogelarten im Wiesenbrütergebiet "Haarmoos" zwischen 1988 und 1992. [Population development and habitat choice of breeding meadowbirds in the "Haarmoos" area between 1988 and 1992]. *Laufener Forschungsbericht* 2: 57-88. In German. (Haus der Natur, Museumplatz 5, A-5020 Salzburg, Austria). Breeding waders: *Vanellus vanellus*, *Gallinago gallinago*, *Numenius arquata*, *Limosa limosa*, *Tringa totanus*, *Tringa glareola* and *Philomachus pugnax*. With discussion on management measures for the area.

Székely, T. 1996. Brood desertion in Kentish Plover *Charadrius alexandrinus*: an experimental test of parental quality and remating opportunities. *Ibis* 138(4): 749-755.

Whitfield, D.P., Brade, J.J., Burton, R.W., Hankinson, K.W. & Young, S. 1996. The abundance of breeding Knot *Calidris canutus islandica*. *Bird Study* 43(3): 290-299. (Scott. Natl. Heritage, Res. & Advisory Serv. Dir., 2 Anderson Place, Edinburgh EH6 5NP, UK). The effectiveness of several census methods for breeding Knot are compared, and estimates of breeding density from six sites

in west Ellesmere Island (Canada) are presented. Total breeding population was estimated to be 61 000 pairs, or 304 000 individuals in autumn. According to the authors, over-optimistic estimates of breeding numbers and productivity, however, may mean that about half of the Knots wintering in Europe may not be *C. c. islandica*.

Winiecki, A. 1992. [Changes of breeding avifauna as the effect of drainage of the Kramsk Marshes]. *Praca Zakladu Biol. Ekol. Ptakow UAM* 1: 83-91. In Polish with English summary. *Gallinago gallinago* no longer breeding.

BEHAVIOUR, ECOLOGY & EVOLUTION

Cadée, N., Piersma, T. & Daan, S. 1996. Endogenous circannual rhythmicity in a non-passerine migrant, the Knot *Calidris canutus*. *Ardea* 84(1/2): 75-84. (Lab. d'Ecol., CNRS URA 258, Univ. Pierre et Marie Curie, Bat. A, 7ième étage, 7 quai St. Bernard, Case 237, F-75252 Paris Cedex 237, France; e-mail: Niels.Cadee@snv.jussieu.fr).

Clark, N.A. 1995. Oystercatcher starvation on the Wash - a case of poor food supplies. *Wash Wader Ringing Group Report 1993-94*: 22-24. (BTO, The Nunnery, Thetford, Norfolk IP24 2PU, UK). *Haematopus ostralegus*.

Clausager, I. 1995. Wing sampling in Denmark - season 1994/95. *Woodcock and Snipe Res. Group Newsletter* 21: 14. (Nat. Environm. Res. Inst., Grenavej 12, DK-8410 Ronde, Denmark). Wings collected of *Scolopax rusticola*, *Gallinago gallinago* and *Lymnocyrtus minimus* by hunters.

Dijkstra, A. 1995. Tiendoornige stekelbaarsjes *Pungitius pungitius* als prooi van Zwarte Ruiters *Tringa erythropus*. (Ten-spined sticklebacks *Pungitius pungitius* as prey of Spotted Redshanks *Tringa erythropus*). *Drentse Vogels* 8: 49-54. In Dutch with English summary. (Dobbenwal 62, NL-9407 AG Assen, Netherlands).

Dirksen, S., Spaans, A.L. & Winden, J. van der 1996. Nachtelijke trek en vlieghoogtes van steltlopers in het voorjaar over de noordelijke havendam van IJmuiden. (Nocturnal migration and flight altitudes of waders at the IJmuiden northern breakwater during spring migration). *Sula* 10(4): 129-142. In Dutch with English summary. (Bureau Waardenburg, P.O. Box 365, NL-4100 AJ Culemborg). Results show that waders passing the coast of the Netherlands in spring, during the day, at short distances from the coast and at low altitudes, continue to do so in the late evening and at night. Observed waders: *Limosa lapponica*, *Haematopus ostralegus*, *Pluvialis apricaria*, *Calidris canutus*, *C. alpina*, *Numenius*

phaeopus, *N. arquata*, *Actitis hypoleucos* and *Tringa totanus*.

Fasola, M., Biddau, L. & Borghesio, L. 1993. Habitat preferences of resident and Palaearctic water birds at Lake Turkana. In: Wilson, R.T. (ed.). *Birds and the African Environment. Proceedings of the Eighth Pan-African Ornithological Congress. Annales Musée Royal de l'Afrique Centrale (Zoologie)* 268: 539-545. (Dipt. Biol. Anim., Univ. Pavia, Pz Botta 9, I-27100 Pavia, Italy). Information on 16 wader species. Use of certain shore types and segregation in foraging locations is described.

Ferrand, Y. & Gossmann, F. 1995. *La Becasse des bois. [The Woodcock]*. Paris: Hatier, 164 pp. In French. Breeding biology, migration, wintering, population dynamics and aspects of hunting *Scolopax rusticola* are described, based on the most recent research results.

Gill, J.A., Clark, J.A., Clark, N.A. & Sutherland, W.J. 1995. Sex differences in moult and migration patterns of Ruff. *Wash Wader Ringing Group Report 1993-94*: 35-36. (School Biol. Sci., Univ. East Anglia, Norwich, Norfolk, NR4 7JT, UK). *Philomachus pugnax*.

Gronstol, G.B. 1996. Aerobic components in the song-flight display of male Lapwings *Vanellus vanellus* as cues in female choice. *Ardea* 84(1/2): 45-55. (Dept. Ecol., Inst. Zool., Univ. Bergen, Allégt. 41, N-5007 Bergen, Norway, e-mail: Gaute.Gronstol@zoo.uib.no).

Hoodless, A. & Draycott, R. 1995. Winter Snipe density and habitat use in SW England. *Woodcock and Snipe Res. Group Newsletter* 21: 3-8. (The Game Conservancy Trust, Fordingbridge, Hampshire SP6 1EF, UK). Information on the use of farmland by wintering *Gallinago gallinago*.

Ketzenberg, C. & Exo, K.-M. 1994. Time budgets of migrating waders in the Wadden Sea: results of the interdisciplinary project Ecosystem Research Lower Saxonian Wadden Sea. *Ophelia (Suppl.)* 6: 315-321. (Inst. Vogelforschung, An der Vogelwarte 21, D-26386 Wilhelmshaven, Germany).

Kiss, J.B., Rékási, J., Sterbetz, I. & Török, Zs. 1995. Habitats and foods used by Woodcocks (*Scolopax rusticola*) during migration through North Dobrogea, Romania, 1970-1989. *Woodcock and Snipe Res. Group Newsletter* 21: 28-35. (I.C.P.D.D., Str. Babadag Nr. 165, 8800 Tilcea, Romania). Habitat use and food have been determined for 16 years during autumn migration along the Danube Delta.

Lavers, C., Haines-Young, R. & Avery, M. 1996. The habitat associations of Dunlin (*Calidris alpina*) in the Flow Country of northern Scotland and an improved model for

predicting habitat quality. *J. Appl. Ecol.* 33: 279-290. (Dept. Geogr., Univ. Nottingham, Univ. Park, Nottingham NG7 2RD, UK).

McNeil, R. & Rompré, G. 1995. Day and night feeding territoriality in Willets *Catoptrophorus semipalmatus* and Whimbrel *Numenius phaeopus* during the non-breeding season in the tropics. *Ibis* 137: 169-176. (Dept. Sci. Biol., Univ. Montréal, C.P. 6128, Succ. "Centre-ville", Montréal, Québec, Canada H3C 3J7).

Meltofte, H. 1996. Are African wintering waders really forced south by competition from northerly wintering conspecifics? Benefits and constraints of northern versus southern wintering and breeding waders. *Ardea* 84(1/2): 31-44. (Danish Polar Ctr., Strandgade 100 H, DK-1401 Copenhagen K, Denmark, e-mail: mel@dpc.dk). See elsewhere this *Bulletin* for summary of paper.

Miller, E.H. 1996. Nuptial vocalizations of male Least Seedsnipe: structure and evolutionary significance. *Condor* 98(2): 418-421. *Thinocorus rumicivorus*.

Minton, C.D.T. 1996. The migration of the Red-necked Stint *Calidris ruficollis*. *Stilt* 29: 24-35. (165 Dalgetty Rd., Beaumaris, Vic. 3193, Australia). Analysis of ringing recoveries and colour leg-flag sightings marked in the East Asian-Australasian Flyway.

Paton, P.W.C. & Edwards Jr., T.C. 1996. Factors affecting interannual movements of Snowy Plovers. *Auk* 113(3): 534-543. *Charadrius alexandrinus*.

Perez-Hurtado, A. 1995. Ecologia alimentaria de Limícolas invernantes en la bahía de Cadiz. (Feeding ecology of wintering waders in Cadiz Bay). *Airo* 6(1-2): 15-23. In Portuguese with English summary.

Ristau, C.A. 1993. The cognitive ethology of an "injury-feigning" plover: a beginning. *Etologia* 3: 57-68. (Barnard. Coll., Dept. Psychol., Columbia Univ., 3009 Broadway, New York City, NY 10027, USA). *Charadrius melodus* and *C. wilsonia*, behaviour towards intruders near nests.

Rogers, K.G., Rogers, D.I. & Minton, C.D.T. 1996. Weights and pre-migratory mass gain of the Red-necked Stint *Calidris ruficollis* in Victoria, Australia. *Stilt* 29: 2-23. (340, Ninks Rd., St. Andrews, Vic. 3761, Australia).

Sandberg, R. & Gudmundsson, G.A. 1996. Orientation cage experiments with Dunlins during autumn migration in Iceland. *J. Avian Biol.* 27(3): 183-188. (Dept. Anim. Ecol., Lund Univ., Ecology Building, S-223 62 Lund, Sweden, E-mail: Roland.Sandberg@zoekol.lu.se). *Calidris alpina*.

Schipper, C.J., Weston, M.A. & Peter, J.M. 1996. Scavenging behaviour of Ruddy

- Turnstones *Arenaria interpres*. *Stilt* 29: 39-40. (3 Orchard Dve., Croydon, Vic. 3136, Australia). Turnstones even entering buildings in search for food.
- Seppi, B.E. 1995. Hudsonian Godwit migration at Carter Spit, Alaska. *West. Birds* 26: 167. (USBLM, 6881 Abbott Loop, Anchorage, AK 99507, USA). *Limosa haemastica*.
- Starks, J. 1996. Intraspecific aggression in the Sharp-tailed Sandpiper *Calidris acuminata*. *Stilt* 29: 40. (c/o RAOU National Office, 415 Riversdale Rd., Hawthorn East, Vic. 3123, Australia).
- Tomkovich, P.S. & Soloviev, M.Y. 1996. Distribution, migrations and biometrics of Knots *Calidris canutus* on Taimyr, Siberia. *Ardea* 84(1/2): 85-98. (Zool. Mus. Moscow State Univ., Bolshaya Nikitskaya Str. 6, 103009 Moscow, Russia, e-mail: tomkovic@1.zoomus.bio.msu.ru). Data collected confirm the suggestion that Taimyr cannot harbour the entire flyway population and that an unknown breeding area should be somewhere in Siberia.
- Turpie, J.K. & Hockey, P.A.R. 1996. Foraging ecology and seasonal energy budgets of estuarine Grey Plovers *Pluvialis squatarola* and Whimbrels *Numenius phaeopus* at the southern tip of Africa. *Ardea* 84(1/2): 57-74. (Percy FitzPatrick Inst. Afr. Ornithol., Univ. Cape Town, Rondebosch 7700, South Africa, e-mail: jturpie@botzoo.uct.ac.za). Distribution, behaviour and energetics were studied at the Zwartkops estuary, South Africa.
- Underhill, L.G. 1995. The relationship between breeding and non-breeding localities of waders: the Curlew Sandpiper *Calidris ferruginea* as an extreme example. *Ostrich* 66(1): 41-46. (Avian Demography Unit, Univ. Cape Town, Rondebosch 7700, South Africa).
- Venema, P. 1995. Gedrag van pleisterende Regenwulpen *Numenius phaeopus* in Noord-Drenthe. (Behaviour of Whimbrels *Numenius phaeopus* at stop-over sites in northern Drenthe). *Drentse Vogels* 8: 26-37. In Dutch with English summary. (Poelakkers 1, NL-9321 EW Peize, Netherlands).
- Venema, P. 1995. Voedsel van de Regenwulp *Numenius phaeopus* in het voorjaar in Noord-Drenthe. (Food of Whimbrels *Numenius phaeopus* in spring in northern Drenthe). *Drentse Vogels* 8: 38-43. In Dutch with English summary.
- Venema, P. 1995. Slaaptrek en gedrag op een slaappleaats van Regenwulpen *Numenius phaeopus*. (Behaviour of Whimbrels *Numenius phaeopus* at the roost). *Drentse Vogels* 8: 44-48. In Dutch with English summary.
- Weber, T.P. & Piersma, T. 1996. Basal metabolic rate and the mass of tissues differing in metabolic scope: migration-related covariation between individual Knots *Calidris canutus*. *J. Avian Biol.* 27(3): 215-224. (Zool. Lab., Univ. Groningen, P.O. Box 14, NL-9750 AA Haren, Netherlands).
- Wieneke, J. & Cross, K. 1996. Food carrying and a new prey item of Marsh Sandpipers *Tringa stagnatilis*. *Stilt* 29: 36. (22 Bishop St., Belgian Gardens, Qld. 4810, Australia). Eating fish.
- Wolf, C. 1995. Austernfischer am Niederrhein - Ökologische Anpassungen an einen neuen Lebensraum. [Oystercatcher at the Lower Rhine - ecological adaptations in a new environment]. *Mittl. Auss. Grietherbusch Zool. Nat. Univ. Köln* 2: 9-11. In German. (Inst. Vogelforschung, An der Vogelwarte 21, D-26386 Wilhelms-haven, Germany). *Haematopus ostralegus*.

DISTRIBUTION & FAUNISTICS

- Adams, S. 1995. Curlew Sandpipers on the Wash. *Wash Wader Ringing Group Report 1993-94*: 30-34. (BTO, The Nunnery, Thetford, Norfolk IP24 2PU, UK). *Calidris ferruginea*.
- Alvarez Laó, C.M. 1995. La Ria de Aviles: un enclave de interes para las Limfcolas. (Abundance for waders in Avilés Estuary). *Airo* 6(1-2): 24-28. In Portuguese with English summary.
- Andres, B.A. & Falxa, G.A. 1995. Black Oystercatcher. In: A. Poole, P. Stettenheim & F. Gill (eds.). *The Birds of North America*, No. 155, 20 pp. (The Birds of North America, c/o The Acad. Natural Sci. Philadelphia, 1900 Benjamin Franklin Parkway, Philadelphia, PA 19103-1195, USA). *Haematopus bachmani*.
- Balachandran, S. 1995. Shore birds of the marine national park in the Gulf of Mannar, Tamil Nadu. *J. Bombay Nat. Hist. Soc.* 92(3): 303-313.
- Bhushan, B., Fry, G., Mundkur, T., Prawiradilaga, D.M., Sonobe, K. & Usui, S. 1993. *A field guide to the waterbirds of Asia*. Tokyo: Wild Bird Society of Japan, 224 pp. ISBN 4-931150-17-9, price £21.95, available from St. Ann's Bookshop, 26 Priory Rd., Malvern, WR14 3DR, UK).
- Blanco, P. 1995. (Recaptures of Kentish Plovers *Charadrius alexandrinus* in a coastal locality of Odessa (Ukraine). *Butll. GCA* 12: 11-14. In Catalan.
- Boele, A., Roomen, M. van & Winden, E. van 1996. *Midwintertelling van Watervogels in Nederland, januari 1995*. [Mid-winter counts of waterbirds in the Netherlands, January 1995]. SOVON-monitoringrapport 96/05, RIZA-rapport BM 95.28, IKC Natuurbeheer

- coproductie 11. Beek-Ubbergen : SOVON, 77 pp. In Dutch with English summary. (SOVON, Rijksweg 178, NL-6573 DG Beek-Ubbergen, Netherlands). Distribution and numbers of *Haematopus ostralegus*, *Recurvirostra avosetta*, *Pluvialis apricaria*, *P. squatarola*, *Vanellus vanellus*, *Calidris canutus*, *C. alba*, *C. alpina*, *Gallinago gallinago*, *Limosa lapponica*, *Numenius arquata*, *Tringa totanus*, *Arenaria interpres*.
- Cayford, J.T. & Waters, R.J. 1996. Population estimates for waders Charadrii wintering in Great Britain, 1987/88-1991/92. *Biol. Conserv.* 77(1): 7-18.
- Dodd, S.L. & Colwell, M.A. 1996. Seasonal variation in diurnal and nocturnal distributions of nonbreeding shorebirds at North Humboldt Bay, California. *Condor* 98(2): 196-207.
- Eubanks, T.L., Jr. 1994. The status and distribution of the Piping Plover in Texas. *Bull. Texas Ornithol. Soc.* 27(1&2): 19-25. *Charadrius melodus*.
- Fasola, M., Biddau, L., Borghesio, L., Baccetti, N. & Spina, F. 1993. Water bird populations at Lake Turkana, February 1992. In: Wilson, R.T. (ed.). *Birds and the African Environment. Proceedings of the Eighth Pan-African Ornithological Congress. Annales Musée Royal de l'Afrique Centrale (Zoologie)* 268: 529-532. (Dipt. Biol. Anim., Univ. Pavia, Pz Botta 9, I-27100 Pavia, Italy). Information on numbers of 30 wader species.
- Ferrand, Y. 1995. Monitoring the French breeding Woodcock, *Scolopax rusticola* population: first results. *Bird Census News* 8(2): 67-71.
- Ferrand, Y. 1995. Rapport Becasse - croule 1995. (Report on Woodcock (in France) - roding season 1995). *Woodcock and Snipe Res. Group Newsletter* 21: 15-22. (Office Natl. de la Chasse, Section Becasse, 5 Rue de St. Thibaut - St. Benoist, F-78610 Auffargis, France). *Scolopax rusticola*, more or less constant trend in the proportion of sites occupied by roding Woodcocks each year.
- Figuerola, J. 1995. Record of longevity for Curlew Sandpiper *Calidris ferruginea*. *Butll. GCA* 12: 15-16. In Catalan. (Dept. Biol. Anim. (vertebrat), Fac. Biol., Univ. Barcelona, Avda. Diagonal, 645, E-08028 Barcelona, Spain).
- Flowers, C. 1996. Spring feast of the Red Knots at Delaware Bay. *Birding* 28(3): 210-215. *Calidris canutus*.
- Fujimaki, Y. 1994. (Distribution and abundance of the Latham's Snipe in south-eastern Hokkaido, Japan). *Strix* 13: 73-78. In Japanese with English summary. *Gallinago hardwickii*.
- Fuller, R.J. 1995. Changes in breeding populations of curlews and lapwings in

- central Buckinghamshire, 1981-1990. *Buckinghamshire Bird Report 1994*: 11-16. (BTO, The Nunnery, Thetford, Norfolk IP24 2PU, UK). *Numenius arquata* and *Vanellus vanellus*.
- García Sánchez, E. 1995. Fenología de Correlimos Oscuro *Calidris maritima* en Asturias y primera estimación de la población invernante. (Phenology of Purple Sandpiper *Calidris maritima* in Asturias (N. Iberia) and first estimate of wintering population). *Airo* 6(1-2): 66-70. In Portuguese with English summary.
- Garcias, P. 1995. (Status of the Kentish Plover (*Charadrius alexandrinus*) in Mallorca. *Anuari Ornitol. Balears*. 10: 25-34. In Catalan with English summary.
- Gaugris, Y. & Vande Weghe, J.-P. 1993. L'importance du delta de la Rusizi (Burundi) pour les migrateurs paléarctiques. (The importance for Palaearctic migrants of the Rusizi flood plain in Burundi). In: Wilson, R.T. (ed.). *Birds and the African Environment. Proceedings of the Eighth Pan-African Ornithological Congress. Annales Musée Royal de l'Afrique Centrale (Zoologie)* 268: 546-552. (B P 2954, Bujumbura, Burundi). Information about c. 30 wader species observed in the Rusizi National Park wetlands in Burundi between 1973-1992.
- Gowthorpe, P., Lamarche, B., Binaux, R., Gueye, A., Lehlou, S.M., Sall, M.A. & Sakho, A.C. 1996. Les oiseaux nicheurs et les principaux limicoles paléarctiques du Parc National du Banc d'Arguin (Mauritanie). [Breeding birds and palearctic migrants of the Banc d'Arguin National Park]. *Alauda* 64(2): 81-126. In French with English summary. Considerable decline in wintering waders. Current and previous counts are discussed for *Pluvialis squatarola*, *Charadrius hiaticula*, *C. alexandrinus*, *Numenius phaeopus*, *N. arquata*, *Limosa lapponica*, *Tringa nebularia*, *T. totanus*, *Arenaria interpres*, *Calidris canutus*, *C. alpina*, *C. ferruginea*, *C. minuta* and *Haematopus ostralegus*.
- Hallett, C.E., Casler, B.R. & Stern, M.A. 1995. Semipalmated Plover nesting on the Oregon coast. *West. Birds* 26: 161-164. (Oregon Nat. Heritage Program, 1205 NW 25th Ave., Portland, OR 97210, USA). *Charadrius semipalmatus*.
- Harris, K. 1996. Population monitoring counts: a report on the winter 1995 and summer 1996 counts. *Stilt* 29: 43-47. (59 Strickland Drive, Wheelers Hill, Vic. 3150, Australia). 45 species of waders counted.
- Harrison, F. 1996. A Whimbrel *Numenius phaeopus* phenomenon in Cairns, Australia. *Stilt* 29: 38. (4/6 Albert St., Cranbrook, Qld. 4814, Australia). Most important Whimbrel site in Australia, with a max. of 1 027 counted on the Cairns foreshore.
- Hirschfeld, E. 1994. The first record of the Golden Plover *Pluvialis apricaria* in the United Arab Emirates. *Emirates Bird Rep.* 18: 109-110.
- Jacobsen, E.M. 1996. Engfugles bestandsændringer og -taetheder 1976-1995, belyst ved hjælp af punktoptællinger. (Point count censuses of meadow birds 1976-1995). *Dansk Orn. Foren. Tidsskr.* 90(3): 115-118. In Danish with English summary. (Fugleregistreringgruppen, Vesterbrogade 140, D-1620 Copenhagen V, Denmark). Population trends for *Gallinago gallinago*, *Tringa totanus*, *Haematopus ostralegus*, *Vanellus vanellus* breeding in meadows. The point count censuses largely confirm patterns found in other investigations, in spite of the small sample size.
- Janzekovic, F. 1995. Winter distribution of the Green Sandpiper (*Tringa ochropus*) and its habitat choice in Slovenia. *Ornis Hung.* 5: 65-66. (Dept. Biol., Univ. Maribor, Koroska c. 160, 62 000 Maribor, Slovenia).
- Komisja Faunistyczna Sekcji Ornitologicznej PTZool 1995. (Rare birds recorded in Poland in 1994: Avifaunistic Commission Report No. 11). *Notatki Ornitol.* 36(3-4): 343-358. In Polish with English summary. (ul. Sienkiewicza 21, 50-335 Wroclaw, Poland). Waders: *Haematopus ostralegus*, *Himantopus himantopus*, *Recurvirostra avosetta*, *Burhinus oedincnemus*, *Glareola nordmanni*, *Charadrius alexandrinus*, *C. morinellus*, *Calidris maritima*, *Xenus cinereus* and *Phalaropus fulicarius*.
- Kosinski, Z. [et al.] 1994. [Importance of the Jeziorsko reservoir for migratory waders]. *Przeegl. Przynr.* 5(2): 61-701. In Polish with English summary. Most numerous: *Vanellus vanellus*, *Calidris alpina*, *Philomachus pugnax*, *Limosa limosa*, *Gallinago gallinago*, *Numenius arquata*, *Tringa erythropus*, *Tringa totanus*, *T. nebularia* and *T. glareola*.
- Leitao, D. 1995. Charadriiformes dos terrenos agrícolas do estuário do Tejo. (Charadriiformes of the Tejo Estuary farmlands). *Airo* 6(1-2): 29-38. In Portuguese with English summary.
- Lorenzo, J.A. & González, J. 1995. Las aves Limícolas de Canarias: breve sinopsis y estudio actual. (The waders of the Canary Islands: short synopsis and actual studies). *Airo* 6(1-2): 1-14. In Portuguese with English summary.
- Marks, J.S. & Redmond, R.L. 1996. Demography of Bristle-thighed Curlews *Numenius tahitiensis* wintering on Laysan Island. *Ibis* 138(3): 438-447.
- Martin, A.A. & Tejada, T.V. 1992. [Wintering of waders in dams of Caceres]. *Bull. Parc. Nat. Delta de l'Ebre* 7: 4-9. In Spanish with English summary. 14 wader species found, including *Calidris temminckii*.
- Maumary, L. & Leuzinger, H. 1996. Seltene Vogelarten und ungewöhnliche Vogelbeobachtungen in der Schweiz im Jahre 1995. (Rare and unusual bird observations in Switzerland in 1995). *Orn. Beob.* 93(4): 329-342. In German with English summary. (Ancienne Poste 11, CH-1030 Bussigny, Switzerland). Waders: *Glareola pratincola*, *Charadrius dubius*, *Chettusia gregaria*, *Calidris melanotos*, *C. ferruginea*, *Gallinago media*, *Tringa erythropus* and *Phalaropus lobatus*.
- Meininger, P.L., Berrevoets, C.M. & Strucker, R.C.W. 1994. *Watervogel-tellingen in het zuidelijk Deltagebied 1987-91*. [Counts of waterbirds in the southern Delta area 1987-91]. Middelburg: Rijksinstituut voor Kust en Zee RIKZ, 381 pp. In Dutch with English summary. (RIKZ, P.O. Box 8039, NL-4330 EA Middelburg, Netherlands). Results of monthly counts of waterbirds in the southern Delta area, SW Netherlands. ISBN 90-369-0493-5, Report RIKZ 94.005. Giving detailed information on the occurrence of many wader species, 1987-1991.
- Meininger, P.L., Berrevoets, C.M. & Strucker, R.C.W. 1996. *Watervogels in de Zoute Delta 1994/95*. [Waterbirds in the salt waters of the Delta area, SW-Netherlands, 1994/95]. Middelburg: Rijksinstituut voor Kust en Zee RIKZ, 72 pp. In Dutch with English summary. Results of monthly counts of waterbirds in the southern Delta area, SW-Netherlands. ISBN 90-369-0375-0, Report RIKZ 96.009. Giving detailed information on the occurrence of many wader species.
- Meran, P. 1995. Some observations on Woodcock (*Scolopax rusticola*) migration in Austria and western Hungary. *Woodcock and Snipe Res. Group Newsletter* 21: 10-11. (Elisabethstr. 41, A-8020 Graz, Austria).
- Minton, C.D.T. 1996. Analysis of overseas movements of Red-necked Stints and Curlew Sandpipers. *Victorian Wader Study Group Bull.* 20: 39-43. (165 Dalgetty Rd., Beaumaris, Vic. 3193, Australia). *Calidris ruficollis* and *C. ferruginea*.
- Minton, C. 1996. Sightings of leg-flagged waders from Victoria, Australia. *Stilt* 29: 48-50. *Charadrius leschenaultii*, *Arenaria interpres*, *Numenius madagascariensis*, *Limosa lapponica*, *Calidris canutus*, *C. ruficollis*, *C. ferruginea* and *C. alba*.
- Minton, C. & Jessop, R. 1996. Further sightings of waders leg-flagged (yellow) in North West Australia. *Stilt* 29: 51-52. *Charadrius leschenaultii*, *Xenus cinereus*, *Heteroscelus brevipes*, *Limosa lapponica*, *L. limosa*, *Calidris tenuirostris*, *C. ruficollis* and *C. ferruginea*.
- Minton, C. 1996. Japanese leg-flagged waders seen in Australia. *Stilt* 29: 55-56. *Heteroscelus brevipes*, *Limosa lapponica*, *L.*

- limosa*, *Calidris canutus*, *C. ruficollis* and *Limicola falcinellus*.
- Minton, C. 1996. Russian colour-banded/flagged waders seen in Australia. *Stilt* 29: 54. *Calidris canutus*, *C. tenuirostris* and *C. ferruginea*.
- Moskoff, W. 1995. Solitary Sandpiper. In: A. Poole, P. Stettenheim & F. Gill (eds.). *The Birds of North America*, No. 156. 16 pp. (The Birds of North America, c/o The Acad. Natural Sci. Philadelphia, 1900 Benjamin Franklin Parkway, Philadelphia, PA 19103-1195, USA). *Tringa solitaria*.
- Mundkur, T. 1993. *A status overview of shorebirds in the East Asian-Australasian Flyway*. Kuala Lumpur: Asian Wetland Bureau. (Asian Wetland Bureau, Inst. Advanced Stud., Univ. Malaya, Lembah Pantai, 59100 Kuala Lumpur, Malaysia).
- OAG Münster 1996. Internationale Limikolen-Zählungen (1979-1994). (WSG Inland Wader Counts (1979-1994). *Vogelwelt* 117(4-6): 307-310. In German with English summary. (OAG Münster, Biol. Stn., Coermühle 181, D-48157 Münster, Germany). Data on *Philomachus pugnax*, *Tringa totanus*, *T. glareola*, *T. nebularia*, *T. ochropus*, *Limosa limosa*, *Gallinago gallinago* and *Actitis hypoleucos*.
- Niedzwiecki, S. & Staszewski, A. 1994. [Waterfowl and waders of the lower Oder between Czulin and Bielinek in August 1992]. *Przegl. Przyr.* 5(1): 65-81. In Polish with English summary.
- Ogilvie, M. and the Rare Breeding Birds Panel. 1996. Rare breeding birds in the United Kingdom in 1994. *Brit. Birds* 89(9): 387-417. (Glencairn, Bruichladdich, Isle of Islay PA49 7UN, UK). Breeding waders: *Recurvirostra avosetta*, *Burhinus oedipnemus*, *Charadrius morinellus*, *Calidris temminckii*, *Calidris maritima*, *Philomachus pugnax*, *Limosa limosa*, *Tringa glareola* and *Phalaropus lobatus*.
- Oliosio, G. 1996. Première observation du Vanneau sociable *Chettusia gregaria* en Rhône-Alpes. [First observation of the Sociable Plover *Chettusia gregaria* in the Rhône-Alpes region (France)]. *Bièvre* 14: 83. In French. (C.O.R.A., Drôme (G.D.E.R.V.), le Mail, F-26230 Grignan, France). *Vanellus gregarius*.
- Page, G.W., Stenzel, L.E., David Shuford, W. & Bruce, C.R. 1991. Distribution and abundance of the Snowy Plover on its western North American breeding grounds. *J. Field Ornithol.* 62: 245-255. *Charadrius alexandrinus*.
- Page, G.W. ...[et al.] 1995. Snowy Plover. In: A. Poole, P. Stettenheim & F. Gill (eds.). *The Birds of North America*, No. 154, 24 pp. (The Birds of North America, c/o The Acad. Natural Sci. Philadelphia, 1900 Benjamin Franklin Parkway, Philadelphia, PA 19103-1195, USA). *Charadrius alexandrinus*.
- Palomeque, S.L. 1995. Aportación a la fenología de Limícolas en los Llanos de Cáceres. (Contribution to phenology of waders in "The Plains of Cáceres"). *Airo* 6(1-2): 56-65. In Portuguese with English summary.
- Paulson, D.R. 1995. Black-bellied Plover. In: A. Poole, P. Stettenheim & F. Gill (eds.). *The Birds of North America*, No. 186, 28 pp. (The Birds of North America, c/o The Acad. Natural Sci. Philadelphia, 1900 Benjamin Franklin Parkway, Philadelphia, PA 19103-1195, USA). *Pluvialis squatarola*.
- Du Plessis, G.J. 1995. Large aggregation of Blackwinged Pratincoles *Glareola nordmanni* in the northern Orange Free State. *Ostrich* 66(1): 40-41.
- Racinskis, E. 1996. (An observation of a wintering Jack Snipe). *Putni daba* 6(2): 21-23. In Latvian with English summary. (Box 10, Riga LV-1047, Latvia). *Lymnocyptes mimimus*.
- Rebassa, M., Suarez, M., Martinez, O. & Wijk, S. 1995. (Winter census of wildfowl and waders in Mallorca and Eivissa/Formentera, January 1996). *Anuari Ornitol. Balears* 10: 45-50. In Catalan with English summary.
- Riddington, R. 1994. The Red-necked Stint on Fair Isle - the first juvenile in Europe. *Birding World* 7: 355-358. *Calidris ruficollis*, UK.
- Robson, B. 1994. The Collared Pratincole in Norfolk. *Birding World* 7: 267. *Glareola pratincola*, UK.
- Rogers, M.J. & Rarities Committee 1996. Report on rare birds in Great Britain in 1995. *Brit. Birds* 89(11): 481-531. Information on 25 wader species. With drawings of *Pluvialis dominica*, *Calidris pusilla* (adult), *Actitis macularia*. Photograph of *Gallinago media*.
- Rösner, H.U. & Günther, K. 1996. Monitoring von rastenden Wat- und Wasservögeln im Wattenmeer. (Monitoring of waterbirds staging in the Wadden Sea). *Vogelwelt* 117(4-6): 295-301. In German with English summary. (WWF-Projektbüro Wattenmeer, Norderstrasse 3, D-25813 Husum, Germany). Information about *Calidris canutus*, *Calidris alpina*, *Pluvialis squatarola* and *Numenius arquata*.
- Rousseau, J.S. 1995. Observations récentes du Pluvier guignard (*Charadrius morinellus*) en Wallonie. Statut de l'espèce en Belgique et notes sur ses habitudes et son identification. [Recent observations of Dotterel (*Charadrius morinellus*) in Wallonia. Status in Belgium and notes on behaviour and identification]. *Aves* 32(4): 177-206. In French.
- Shackford, J.S. 1995. Melanistic Mountain Plover in Cimarron County, Oklahoma. *Bull. Okla. Ornithol. Soc.* 28: 4-5. *Charadrius montanus*.
- Shuford, W.D., Page, G.W. & Hickey, C.M. 1995. Distribution and abundance of Snowy Plovers wintering in the interior of California and adjacent states. *West. Birds* 26: 82-98. (Pt. Reyes Bird Obs., 4990 Shoreline Hwy., Stinson Beach, CA 94970, USA). *Charadrius alexandrinus*.
- Soulliere, G.J. 1993. Short-billed Dowitcher nest found in Ontario. *Ontario Birds* 11: 109-110. (Mich. Dept. Nat. Resour., Wildl. Div., Box 798, Sault Ste. Marie, MI 49783, USA). *Limnodromus griseus*.
- Underhill, L.G. 1995. Migratory birds. In: Cowan, G.I. (ed). *Wetlands of South Africa*. Pretoria: Dept. Environm. Affairs and Tourism. Pp. 163-177. (Dept. Stat. Sci., Univ. Cape Town, Rondebosch 7700, South Africa). *Charadrius hiaticula*, *C. leschenaultii*, *Pluvialis squatarola*, *Arenaria interpres*, *Xenus cinereus*, *Tringa hypoleucos*, *T. glareola*, *T. stagnatilis*, *T. nebularia*, *Calidris canutus*, *C. ferruginea*, *C. minuta*, *C. alba*, *Philomachus pugnax*, *Limosa lapponica*, *Numenius arquata* and *N. phaeopus* in South Africa).
- Volet, B. & Schmid, H. 1996. Übersicht über das Brutgeschehen und andere ornithologische Ereignisse 1993 und 1994 in der Schweiz. (Rare and unusual records of breeding, migrating and wintering bird species, 1993 and 1994). *Orn. Beob.* 93(4): 315-328. In German with English and French summary. (Schweizerische Vogelwarte, CH-6204 Sempach, Switzerland). Waders: *Charadrius dubius* (breeding), *Gallinago gallinago* (suspected breeding), *Numenius arquata* (probable breeding), *Actitis hypoleucos* (breeding). Records of *Himantopus himantopus*, *Burhinus oedipnemus*, *Charadrius alexandrinus*, *C. morinellus* and *Tringa stagnatilis*.
- Warnock, S.E. & Takekawa, J.Y. 1996. Wintering site fidelity and movement patterns of Western Sandpipers *Calidris mauri* in the San Francisco Bay estuary. *Ibis* 138: 160-167.
- Zielinski, M. 1995. (Ringing of waders (Charadrii) at the Vistula River mouth in 1987-1994). *Notatki Ornitol.* 36(1-2): 181-190. In Polish with English summary. (Stacja Ornitol. IE PAN, ul. Nadwiślanska 108, 80-680 Gdansk 40, Poland). With information on *Calidris alpina*, *C. ferruginea*, *C. minuta* changes in numbers caught over the years.

MISCELLANEOUS

- Aymerich, F.R. 1995. Ecología de humedales y conservación de Limícolas en el SE de España: experiencias de investigación aplicables a la gestión. (Ecology of wetlands and wader conservation in Southeastern Spain: research experiences applicable to management). *Airo* 6(1-2): 39-55. In Portuguese with English summary.
- Burton, N.H.K., Evans, P.R. & Robinson, M.A. 1996. Effects on shorebird numbers of disturbance, the loss of a roost site and its replacement by an artificial island at Hartlepool, Cleveland. *Biol. Conserv.* 77(2-3): 193-201.
- Dickson, H.L. & Smith, A.R. 1991. Use of Landsat Thematic Mapper and multi-spectral scanning imagery to identify habitats and shorebird nesting areas on the outer Mackenzie River Delta, NWT. In: Marsh, P. & Ommanney, C.S.L. (eds.). *Mackenzie Delta: Environmental interactions and implications of development*. National Hydrology Res. Inst. Symp. No. 4, Saskatoon, Saskatchewan. Saskatchewan: National Hydrology Res. Inst., Environment Canada, pp. 91-106.
- Estelle, V.B., Mabee, T.J. & Farmer, A.H. 1996. Effectiveness of predator exclosures for Pectoral Sandpiper nests in Alaska. *J. Field Ornithol.* 67(3): 447-452. *Calidris melanotos*.
- Figuerola, J., Velarde, R., Bertolero, A. & Cerdá, F. 1996. Abwesenheit von Haematozoa bei einer Brutpopulation des Seeregenpfeifers *Charadrius alexandrinus* in Nordspanien. (Absence of haematozoa in a breeding population of Kentish Plover *Charadrius alexandrinus* in Northern Spain). *J. Orn.* 137(4): 523-525. In German with English summary. (Dept. Biol. Anim. (Vertebrats), Fac. Biol., Univ. Barcelona, Avda. Diagonal 645, E-08028, Spain).
- Figuerola, J. & Gustamante, L. 1995. Principales objetivos del marcaje de Limícolas en el Delta de l'Ebro. (Main objectives of wader ringing at the Ebro Delta). *Airo* 6(1-2): 71-75. In Portuguese with English summary.
- Gratto-Trevor, C.L. 1994. *Use of Landsat TM Imagery in determining priority shorebird habitat in the outer Mackenzie Delta, NWT*. NOGAP subproj. C.24. Unpubl. rep. Saskatoon: Canadian Wildl. Serv., 217 pp. (Canadian Wildl. Serv., Environment Canada, Prairie and Northern Wildl. Res. Ctr., 115 Perimeter Rd., Saskatoon, Saskatchewan S7N 0X4, Canada).
- Johnston, R.M. & Barson, M.M. 1993. Remote sensing of Australian wetlands: an evaluation of Landsat TM data for inventory and classification. *Aust. J. Mar. Freshwater Res.* 44: 235-252.
- Melvin, S.M., MacIvor, L.H. & Griffin, C.R. 1992. Predator exclosures: a technique to reduce predation at Piping Plover nests. *Wildl. Soc. Bull.* 20: 143-148. *Charadrius melodus*.
- Minton, C. 1996. Comparison of flag sightings versus recoveries for waders marked in Victoria, Australia. *Stilt* 29: 39. (165 Dalgetty Rd., Beaumaris, Vic. 3193, Australia).
- Olivier, G.N. 1995. Managing Snipe habitats. *Woodcock and Snipe Res. Group Newsletter* 21: 9-10. (21 rue du Colonel Moll, F-75017 Paris, France). *Gallinago gallinago*.
- Pittaway, R. 1993. Leg and bill colour of Purple Sandpipers. *Ontario Birds* 11: 107-109. *Calidris maritima*.
- Retallick, R.W.R. & Bolitho, E.E. 1996. Disturbance of Beach Thick-knee *Esacus magnirostris* by aircraft. *Stilt* 29: 36-37. (Dept. Zool., James Cook Univ., Townsville, Qld. 4811, Australia).
- Whittingham, M.J. 1996. The use of radiotelemetry to measure the feeding behavior of breeding European Golden Plovers. *J. Field Ornithol.* 67(3): 463-470. *Pluvialis apricaria*.

