

Recent Publications on Waders 74

In trying to make these lists really as complete as possible, and the next one even more complete and up to date, your help is necessary!! Please send new titles, abstracts, reprints, copies, reports, and (important too!) omissions or corrections to: **Henk Koffijberg, Glenn Millerweg 102, 1311 RW Almere, The Netherlands.**

Remember to include, if possible, the address of the first author, and translations of titles if they are in other languages than English.

For help with this issue I'm very grateful to T. Piersma, I. Bainbridge, P. Triplet, K.-M. Exo, T.W. Dougall, C. Zöckler, R.B. Lanctot, R. McNeil, W. Meissner, J. Melter, T.D. Evans, B. Nikolov, and M. Schmitz.

Translated titles from languages other than English are bracketed; with round brackets if the translation is by the author(s), square brackets if not.

N.B. a copy of your paper or abstract would be very helpful to enable me to add some keywords or give added annotation to the titles.

So far, more than 7000 titles have been stored in the WSG/RPW database. Still not enough! Keep sending more and use this database!

Breeding

Amat, J.A., Fraga, R.M. & Arroyo, G.M. 1999. Reuse of nesting scrapes by Kentish Plovers. *Condor* 101(1): 157-158. *Charadrius alexandrinus*

Amat, J.A., Fraga, R.M. & Arroyo, G.M. 1999. Brood desertion and polygamous breeding in the Kentish Plover *Charadrius alexandrinus*. *Ibis* 141(4): 596-607. (Estación Biológica de Doñana, C.S.I.C., Apartado 1056, E-41080 Sevilla, Spain)

Arnold, T.W. 1999. What limits clutch size in waders?. *J. Avian Biol.* 30(2): 216-220. (Delta Waterfowl & Wetlands Res. Stn., RR 1, Portage la Prairie, Manitoba R1N 3A1 Canada, e-mail: tarnold@deltawaterfowl.com). Clutch

enlargement in Holarctic waders leads to greater rates of nest predation, egg damage, and hatching failure. Together this is sufficient to limit clutch size among most wader species.

Bealey, C.E., Green, R.E., Robson, R., Taylor, C.R. & Winspear, R. 1999. Factors affecting the numbers and breeding success of Stone Curlews *Burhinus oedicnemus* at Porton Down, Wiltshire. *Bird Study* 46(2): 145-156. (RSPB, The Lodge, Sandy, Bedfordshire SG19 2DL, UK). High rabbit population has positive effect on breeding numbers and breeding success.

Bil, W. & Schuur, J. 1999. Overlevingskansen van Kieviten pulli op intensief gebruikt grasland. [Survival of Lapwing pulli on intensively used grassland]. *Vogeljaar* 47(2): 55-63. In Dutch with English summary. (De Buorren 91, NL-8408 HL Lippenhuizen, Netherlands, e-mail: W.Bil@hetnet.nl). *Vanellus vanellus*.

Blomqvist, D., Johansson, O.C. & Götmark, F. 1997. Parental quality and egg size affect chick survival in a precocial bird, the Lapwing *Vanellus vanellus*. *Oecologia* 110: 18-24.

Davis, W.E. Jr. & Armstrong, C. 1997. Killdeer nest on gravel roofs of office buildings in Canton, Massachusetts. *Bird Observer* 25: 94-98. (c/o BOEM, P.O. Box 236, Arlington, MA 02174, USA). *Charadrius vociferus*.

Espie, R.H.M., James, P.C., Brigham, R.M. 1998. The effect of flooding on Piping Plover *Charadrius melodus* reproductive success at Lake Diefenbaker, Saskatchewan, Canada. *Biol. Conserv.* 86(2): 215-222.

Espie, R.H.M., Brigham, R.M. & James, P.C. 1996. Habitat selection and clutch fate of Piping Plovers *Charadrius melodus* breeding at Lake Diefenbaker, Saskatchewan. *Can. J. Zool.* 74(6): 1069-1075.

Flynn, L., Nol, E. & Zharikov, Y. 1999. Philopatry, nest-site tenacity, and mate fidelity of Semipalmated Plovers. *J. Avian Biol.* 30(1): 47-55. (Watershed Ecosystems Grad. Progr., Trent Univ., Peterborough, Ontario, Canada K9J 7B8, e-mail: lflynn@sd48.mountain-inter.net). *Charadrius semipalmatus*, Manitoba, Canada.

Flynn, L. 1997. *Philopatry, nest-site*

tenacity, and mate fidelity in Semipalmated Plovers (Charadrius semipalmatus) breeding at Churchill, Manitoba. M.Sc. thesis, Trent University. (Watershed Ecosystems Grad. Progr., Trent Univ., Peterborough, Ontario, Canada K9J 7B8, e-mail: lflynn@sd48.mountain-inter.net).

Frost, D. 1997. Observations on the territorial behaviour of Kentish Plover *Charadrius alexandrinus* at Khor Dubai. *Emirates Bird Rep.* 19: 126. (1 Shakespeare Rd., St. Ives, Cambs PW17 4TR, UK)

Grave, C., Melter, J. & ten Thoren, B. 1998. Zur Siedlungsdichte der Kiebitz *Vanellus vanellus* in intensiv genutzten Agrarlandschaften Nordwestdeutschlands. (Densities of Lapwings *Vanellus vanellus* in intensively used farmland in Lower Saxony). *Vogelkd. Ber. Niedersachs.* 30(2): 77-86. In German with English summary. (Lindenstrasse 33, D-49191 Belm/OS, Germany)

Heg, D. & van Treuren, R. 1998. Female-female cooperation in polygynous Oystercatchers. *Nature* 391: 687-691. *Haematopus ostralegus*

Hegy, Z. 1996. Laying date, egg volumes and chick survival in Lapwing (*Vanellus vanellus* L.), Redshank (*Tringa totanus* L.), and Black-tailed Godwit (*Limosa limosa* L.). *Ornis Hung.* 6: 1-7.

De Jong, A. & Sandström, L. 1998. (Two Curlews *Numenius arquatus* laid eggs in the same nest?). *Ornis Svecica* 8(3): 129-130. (Hössjö 115, S-905 86, Umeå, Sweden)

Kruk, M., Noordervliet, M. & ter Keurs, W. 1999. Overleving van gruttoekikens op boerengrasland. [Survival of Black-tailed Godwit chicks in agricultural grasslands]. *Vogeljaar* 47(2): 49-54. In Dutch. (Stichting Landschapsbeheer Zuid-Holland, P.O. Box 558, NL-2800 AN Gouda, Netherlands). *Limosa limosa*.

Lanctot, R.B., Weatherhead, P.J. Kempnaers, B. & Scribner, K.T. 1998. Male traits, mating tactics and reproductive success in the Buff-breasted Sandpiper, *Tryngites subruficollis*. *Anim. Behav.* 56: 419-432. (US Geol. Surv., Biol. Res. Div., Alaska Biol. Sci. Ctr., 1011 E. Tudor Rd., Anchorage, AK 99503-6199, USA, e-mail:

richard_lanctot@nbs.gov). Study results suggest that males may use alternative mating tactics more readily than males in other lek-breeding waders because unpredictable breeding conditions in its high arctic breeding area leads to low lek stability; and males are not constrained by morphological markings that indicate status or sex. Both characteristics may reduce reproductive benefits associated with males adopting on mating tactic and result in a sort of scramble competition in which males switch between tactics as local conditions change.

Lanctot, R.B., Scribner, K.T., Kempnaers, B. & Weatherhead, P.J. 1997. Lekking without a paradox in the Buff-breasted Sandpiper. *Am. Nat.* 149(6): 1051–1070. *Tryngites subruficollis*.

Liker, A. & Székely, T. 1999. Parental behaviour on the Lapwing *Vanellus vanellus*. *Ibis* 141(4): 608–614. (Behav. Biol. Res. Grp., Dept. Ecol., Univ. Vet. Sci., Pf. 2, H–1400 Budapest, Hungary, e-mail: aliker@ns.univet.hu)

Michaelis, H. 1998. Brut des Austernfischers *Haematopus ostralegus* auf dem Flachdach eines Schulgebäudes in Mettingen, Kreis Steinfurt. (Breeding by Oystercatchers *Haematopus ostralegus* on the flat roof of a school building at Mettingen in the German federal state of North Rhine-Westphalia). *Charadrius* 34(1): 57–59. In German. (Berg up Sonn 23, D–49497 Mettingen, Germany).

Möller, D. 1998. Brutvorkommen der Uferschnepfe (*Limosa limosa*) in Ackerflächen. (Black-tailed Godwits (*Limosa limosa*) breeding in arable fields). *Charadrius* 34(1): 60–63. In German with English summary.

Nol, E., Sullivan Blanken, M. & Flynn, L. 1997. Sources of variation in clutch size, egg size and clutch completion dates of Semipalmated Plovers in Churchill, Manitoba, Canada. *Condor* 99: 389–396. (Biol. Dept., Trent Univ., Peterborough, Ontario, Canada K9J 7B8, e-mail: enol@trentu.ca). *Charadrius semipalmatus*

Orr, N. 1999. Egg-burying behaviour of Kentish Plovers. *Brit. Birds* 92(7): 367–368. (4 Denham Drive, Christchurch, Dorset BH23 5AT, UK). *Charadrius alexandrinus* burying eggs suggesting temperature control. Parish, D.M.B. & Coulson, J.C. 1998.

Parental investment, reproductive success and polygyny in the Lapwing, *Vanellus vanellus*. *Anim. Behav.* 56: 1161–1167.

Rebergen, A. ...[et al.] 1998. Breeding success and predation at nests of Banded Dotterel (*Charadrius bicinctus*) on braided riverbeds in the central South Island, New Zealand. *N.Z.J. Ecol.* 22: 33–41. (Dept. Conserv., P.O. Box 191, Masterton, New Zealand). Very low breeding success. Island nests more successful than nests on the mainland.

Schekkerman, H., van Roomen, M.W.J. & Underhill, L.G. 1998. Growth, behaviour of broods and weather-related variation in breeding productivity of Curlew Sandpipers *Calidris ferruginea*. *Ardea* 86(2): 153–168. (DLO Inst. Forrestry & Nat. Res., P.O. Box 23, NL–6700 AA Wageningen, Netherlands, e-mail: h.schekkerman@ibn.dlo.nl).

Stasaitis, J. 1998. The breeding peculiarities of the Golden Plover *Pluvialis apricaria* in Lithuania. *Acta Zool. Lituanica, Ornithol.* 8(2): 141–147.

Teather, K. & Nol, E. 1997. Mixed sexual dimorphism in the Semipalmated Plover. *Condor* 99: 803–806. (Nol: Biol. Dept., Trent Univ., Peterborough, Ontario, Canada K9J 7B8, e-mail: enol@trentu.ca). *Charadrius semipalmatus*

Tuttle, D.C., Stein, R. & Lester, G. 1997. Snowy Plover nesting on Eel River gravel bars, Humboldt County [California]. *West. Birds* 28: 174–176. (Dept. Public Works, Nat. Resour. Div., 1106 Second St., Eureka, CA 95501, USA). *Charadrius alexandrinus*.

Widemo, F. 1998. Alternative reproductive strategies in the Ruff, *Philomachus pugnax*: a mixed ESS?. *Anim. Behav.* 56: 329–336. (Dept. Zool., Univ. Uppsala, Villavögen 9, S–752 36 Uppsala, Sweden).

Behaviour, ecology & evolution

Alexander, S.A., Hobson, K.A., Gratto-Trevor, C.L. & Diamond, A.W. 1996. Conventional and isotopic determinations of shorebird diets at an inland stopover: the importance of invertebrates and *Potamogeton pectinatus* tubers. *Can. J. Zool.* 74(6): 1057–1068.

Andres, B.A. 1998. Shoreline habitat use of Black Oystercatchers breeding in Prince William Sound, Alaska. *J. Field Ornithol.* 69(4): 626–634. *Haematopus bachmani*

Baccetti, N., Serra, L., Cherubini, G. & Magnani, A. 1999. Timing of attachment to wintering site as revealed by experiment displacements of Dunlins *Calidris alpina*. *J. Ornithol.* 140(3): 309–317. (Inst. Naz. Fauna Selvatica, I–40064 Ozzano Emilia BO, Italy, e-mail: infszumi@iperbole.bologna.it)

Backwell, P.R.Y., O'Hara, P.D. & Christy, J.H. 1998. Prey availability and selective foraging in shorebirds. *Anim. Behav.* 55: 1659–1667. (Smithsonian Trop. Res. Inst., P.O. Box 2072, Balboa, Panama). Waders studied a.o. *Numerius phaeopus*, *Catoptrophorus semipalmatus*, *Arenaria interpres*, *Pluvialis squatarola*, *Charadrius wilsonia*, *C. semipalmatus*.

Barbosa, A. 1997. The effects of predation risk on scanning and flocking behavior in Dunlin. *J. Field Ornithol.* 68: 607–612. (Lab. Ecol., Univ. Pierre et Marie Curie, 7 Quai St., Bernard, Case 237, F–75252 Paris Cedex 05, France). *Calidris alpina*.

Barbosa, A. 1996. Relationship between bill morphology and preening behaviour in waders. *Ethol. Ecol. & Evol.* 8: 291–296.

Brandsma, O. 1999. Het belang van bemesting voor het voedselaanbod van weidevogels. (The importance of organic fertilizers and/or lime for earthworms and meadow birds in a wetland in NW–Overijssel). *Levende Natuur* 100(4): 118–123. In Dutch with English summary. (Klosseweg 12, NL–7946 KH Waneperveen, Netherlands). Relation between biomass of earthworms and densities of breeding *Vanellus vanellus* and *Limosa limosa*.

Buchanan, J.B. ...[et al.] 1996. Body mass and lipid levels of shorebirds collected in western Washington. *Northwest Nat.* 77: 51–54. (Washington Dept. Fish Wildl., Wildl. Manage. Div., 600 Capitol Way N., Olympia, WA 98501 USA). *Charadrius semipalmatus*, *Pluvialis squatarola*, *Arenaria interpres*, *Calidris canutus*, *C. alba*, *C. mauri*, *C. alpina*.

Burger, J. & Tsipoura, N. 1998. Experimental oiling of Sanderlings (*Calidris alba*): behavior and weight

- changes. *Environ. toxicol. chem.* 17(6): 1154–1158. (Tsipoura: Rutgers Univ., Div. Life Sci., Nelson Biol. Labs., 604 Allison Rd., Piscataway, NJ 08854, e-mail: tsipoura@rci.rutgers.edu).
- Calvo, B. & Furness, R.W. 1996. Effects of environmental conditions on aerial feeding by Collared Pratincoles *Glareola pratincola* breeding in southwest Spain. *Avocetta* 20: 97–100. (Appl. Ornithol. Unit, Inst. Biomed. Life Sci., Graham Kerr Bldg., Glasgow Univ., Glasgow G12 8QQ, UK)
- Dann, P. 1999. Feeding periods and supratidal feeding of Red-necked Stints and Curlew Sandpipers in Western Port, Victoria. *Emu* 99(3): 218–222. (Penguin Reserve Comm. Manage., P.O. Box 97, Cowes, Philip Isl., Vic. 3922, Australia). *Calidris ruficollis*, *C. ferruginea*.
- Delehanty, D.J., Fleischer, R.C., Colwell, M.A. & Oring, L.W. 1998. Sex-role reversal and the absence of extra-pair fertilization in Wilson's phalaropes. *Anim. Behav.* 55: 995–1002. (Dept. Bio., Univ. Nevada, Reno, NV 89557, USA). *Phalaropus tricolor*.
- Dietz, M.W., Piersma, T. & Dekinga, A. 1999. Body-building without power training: endogenously regulated pectoral muscle hypertrophy in confined shorebirds. *J. Exp. Biol.* 202: 2831–2837. (Ctr. Ecol. Evol. Stud., Zool. Lab., Univ. Groningen, P.O. Box 14, NL-9750 AA Haren, Netherlands, e-mail: m.w.dietz@biol.rug.nl)
- Le Dréan-Quénech'du & Mahéo, R. 1997. [Impact of waders on molluscs in the Traicts du Croisic, west France]. *Alauda* 65: 131–149. In French with English summary. (Lab. Evol. Syst. Nat. Mod., Univ. de Rennes, Campus de Beaulieu, F-35042 Rennes Cedex, France)
- Goss-Custard, J.D., Cayford, J.T. & Lea, S.E.G. 1998. The changing trade-off between food finding and food stealing in juvenile Oystercatchers. *Anim. Behav.* 55: 745–760. (ITE, Furzebrook Res. Stn., Wareham, Dorset BH20 5AS, UK). *Haematopus ostralegus*
- Hitchcock, C.L. & Gratto-Trevor, C. 1997. Diagnosing a shorebird local population decline with a stage-structured population model. *Ecology* 78: 522–534.
- Hockey, P.A.R., Turpie, J.K., Plagányi, E.E. & Philips, T.E. 1999. Scaling patterns in the foraging behaviour of sympatric plovers: effects of body size and diet. *J. Avian Biol.* 30(1): 40–46. (Percy FitzPatrick Inst. Afr. Ornithol., Univ. Cape Town, Rondebosch, S. Africa 7701, e-mail: phockey@botzoo.uct.ac.za). *Dromas ardeola*, *Pluvialis squatarola*, *Charadrius leschenaultii*, *C. mongolus*, *C. hiaticula*, Kenya.
- Hughey, K.F.D. 1997. The diet of the Wrybill (*Anarhynchus frontalis*) and the Banded Dotterel (*Charadrius bicinctus*) on two braided rivers in Canterbury, New Zealand. *Notornis* 44: 185–193. (Dept. Resour. Manage., P.O. Box 56, Lincoln Univ., New Zealand).
- Johansson, O.C. & Blomqvist, D. 1996. Habitat selection and diet of Lapwing *Vanellus vanellus* chicks on coastal farmland in S.W. Sweden. *J. Appl. Ecol.* 33: 1030–1040.
- De Leon, M.T. & Smith, L.M. 1999. Behavior of migrating shorebirds at North Dakota prairie potholes. *Condor* 101(3): 645–654. (Dept. Range, Wildl. Fish. Manage., Texas Tech. Univ., Lubbock, TX 79409–2125, USA). Diurnal time-activity budgets of *Recurvirostra americana*, *Limnodromus scolopaceus*, *Calidris himantopus*, *C. pusilla*, *C. minutilla* were examined to evaluate the importance of prairie pothole wetlands for shorebirds.
- Liker, A. & Székely, T. 1997. Aggression among female Lapwings, *Vanellus vanellus*. *Anim. Behav.* 54: 797–802. (Zool. Ctr., Univ. Vet. Sci., P.O. Box 2, H-1400 Budapest, Hungary)
- Lindström, Å. 1998. Mass and morphometrics of Little Stints *Calidris minuta* on autumn migration along the arctic coast of Eurasia. *Ibis* 140(1): 171–174.
- Lunn, J. 1999. Northern Lapwings resting on urban roof. *Brit. Birds* 92(7): 368. (11 Springfields, Barnsley S75 1JS, UK). *Vanellus vanellus*.
- Mason, C.F. & MacDonald, S.M. 1999. Habitat use by Lapwings and Golden Plovers in a largely arable landscape. *Bird Study* 46(1): 89–99. (Dept. Biol. Sci., Univ. Essex, Wivenhoe Pk., Colchester CO4 3SQ, UK). *Vanellus vanellus* and *Pluvialis apricaria*.
- McNeil, R. & Rodriguez Silva, J.R. 1996. Ecological significance and sensorial aspects of nocturnal foraging in shorebirds. In: Cabana, T. (ed.). *Animals in their environment*. Orbis, Frelighsburg (Quebec), 1996, p. 23–58. (Dept. Sci. Biol., Univ. Montréal, C.P. 6128, Succ. 'Centre-ville', Montréal, PQ, H3C 3J7 Canada, e-mail: mcneilr@ere.unmontreal.ca). Review of occurrence, controlling factors, inter-relationships, success, strategies, and benefits of nocturnal foraging in Charadrii, and discusses its significance.
- Melter, J. & Sauvage, A. 1997. Measurement and moult of Ruffs *Philomachus pugnax* wintering in West Africa. *Malimbus* 19: 12–18. (OAG Münster, Biol. Stn., Coermühle 181, D-48157 Münster, Germany). Measurement of wing, bill and tarsus, and information about leg colour.
- Mihue, J.R., Trost, C.H. & Mihue, T.B. 1997. Shorebird predation on benthic macroinvertebrates in an irrigation reservoir. *Great Basin Naturalist* 57: 245–253. (Dept. Biol. Sci., Idaho State Univ., Pocatello, ID 83209–8007, USA)
- Norris, K. & Johnstone, I. 1998. Interference competition and the functional response of Oystercatchers searching for cockles by touch. *Anim. Behav.* 56: 639–650. *Haematopus ostralegus*.
- Petersen, B. & Exo, K.-M. 1999. Predation of waders and gulls on *Lanice conchilega* tidal flats in the Wadden Sea. *Mar. Ecol. Prog. Ser.* 178: 229–240. (Inst. Vogelforschung 'Vogelwarte Helgoland' An der Vogelwarte 21, D-26386 Wilhelmshaven, Germany, e-mail: exo@ifv-terramare.fh-wilhelmshaven.de). Waders investigated: *Haematopus ostralegus*, *Calidris alpina*, *Numenius arquata*, *Tringa totanus*, East Frisian Wadden Sea, Germany.
- Piersma, T., Dekker, M. & Sinninghe Damst., J. 1999. An avian equivalent of make-up? *Ecol. Letters* 2: 201–203. (NIOZ, P.O. Box 59, NL-1790 AB Den Burg, Texel, Netherlands, e-mail: theunis@nioz.nl). *Calidris canutus* makes a complete switch from commonly occurring monoester preen waxes to a much rarer class of higher-molecular-weight diester waxes at the time of take-off to the high arctic breeding grounds.
- Piersma, T., Gudmundsson, G.A. & Lilliendahl, K. 1999. Rapid changes in the size of different functional organ and

muscle groups during refueling in a long-distance migrating shorebird. *Physiol. Biochem. Zool.* 72(4): 405–415. Study of adaptive value of size changes in different organs and muscle groups in *Calidris canutus islandica* in relation to migration.

Piersma, T. & Klaassen, M. 1999. *Methods of studying the functional ecology of protein and organ dynamics in birds*. In: Adams, N.J. & Slotow, R.H. (eds.). Proc. 22 Int. Ornithol. Congr., Durban, p. 36–51. The potential of new non-terminal methods (ultrasound, nuclear magnetic resonance imaging and computer tomography), which allow repeated size estimation of various organs, was evaluated in recent studies on shorebirds and swans. These methods allow non-invasive, repeated quantification of the size of different organs in individual birds.

Rojas, L.M., McNeil, R., Cabana, T. & Lachapelle, P. 1999. Diurnal and nocturnal visual capabilities in shorebirds as a function of their feeding strategies. *Brain behav. evol.* 53: 29–43. (McNeil: Dept. Sci. Biol., Univ. Montréal, C.P. 6128, Succ. 'Centre-ville', Montréal, PQ, H3C 3J7 Canada, e-mail: mcnair@ere.unmontreal.ca). *Charadrius wilsonia*, *Limnodromus griseus*, *Scolopax minor*, *Himantopus himantopus*, *Catoptrophorus semipalmatus*.

Ruggiero, R.G. 1996. Interspecific feeding associations mutualism and semi-parasitism between *Hippopotami Hippopotamus amphibius* and African Jacanas *Actophilornis africanus*. *Ibis* 138: 346–348. (Wild. Conserv. Soc., BP 14537, Brazzaville, Republic of Congo)

Serra, L., Whitelaw, D.A., Tree, A.J. & Underhill, L.G. 1999. Molt, mass and migration of Grey Plovers *Pluvialis squatarola* wintering in South Africa. *Ardea* 87(1): 71–81. (Ist. Naz. Fauna Selvatica, via Ca' Fornacetta 9, I-40064 Ozzano Emilia BO, Italy)

Smith, H.T. & Rodgers Jr., J.A. 1996. Sanderling *Calidris alba* flight duration response to inadvertent disturbance. *Fla. Field Nat.* 24(3): 74–75.

Steiof, K. 1999. Water Rail killing and eating wader. *Brit. Birds* 92(4): 204. (Seestrasse 60, D-133347 Berlin, Germany). *Rallus aquaticus* versus *Calidris alpina* or *Gallinago gallinago*.

Sueur, F. 1999. Mue tardive chez le Bécasseau variable *Calidris alpina*. [Late moulting in the Dunlin *Calidris alpina*]. *Avifaune Picarde* 7: 32. In French. (9 rue du Champ neuf, Le Bout des Crocs, F-80120 Saint-Quentin-en-Tourmont, France).

Triplet, P., Sueur, F., Fagot, C., Oget, E. & Desprez, M. 1999. Réponses de l'Huîtrier-pie *Haematopus ostralegus* à une diminution de sa ressource alimentaire principale en Baie de Somme: la Coque. [Response of Oystercatchers *Haematopus ostralegus* to a reduction of its main food source in the Somme bay: the cockle. *Alauda* 67(2): 145–153. (SMACOPI, Réserve Naturelle Baie de Somme, Stn. Biol. Blanquetaque, 1 place de l'Amiral Courbet, F-80100 Abbeville, France, e-mail: blanquetaque@dyadel.net)

Triplet, P., Bacquet, S., Lengignon, A., Oget, E. & Fagot, C. 1999. Effets de dérangements sur l'Huîtrier-pie *Haematopus ostralegus* en Baie de Somme. (Effects of disturbance on the Oystercatcher *Haematopus ostralegus* in the Baie de Somme. *Gibier Faune Sauvage* 16(1): 45–64. In French with English and German summary. Recreation and illegal fishing (cockles) were the most important cause of human disturbance of Oystercatchers.

Triplet, P., Stillman, R.A. & Goss-Custard, J.D. 1999. Prey abundance and the strength of interference in a foraging bird. *J. Anim. Ecol.* 68: 254–265. (Stillman: Inst. Terr. Ecol., Furzebrook Res. Stn., Wareham, Dorset BH20 5AS, UK, e-mail: r.stillman@ite.ac.uk)

Tsipoura, N. & Burger, J. 1998. *Shorebird diet during spring migration stopover on Delaware Bay*. Piscataway: Rutgers University, Dept. Biol. Sci., 10 p. (Rutgers Univ., Div. Life Sci., Nelson Biol. Labs., 604 Allison Rd., Piscataway, NJ 08854, e-mail: tsipoura@rci.rutgers.edu).

Tsipoura, N. & Burger, J. 1999. Shorebird diet during spring migration stopover on Delaware Bay. *Condor* 101(3): 635–644. Horseshoe crab eggs on the main dish for *Calidris canutus*, *Arenaria interpres*, *C. alba*, *C. pusilla*. All had an increase in body mass up to 70–80% during their stay at Delaware Bay.

Violet, F. 1998. Mue tardive d'un

Pluvier dor, *Pluvialis apricaria*. [Late moulting in the Golden Plover *Pluvialis apricaria*]. *Avifaune Picarde* 6: 96. In French. (11/13 boulevard Carnot, F-80460 Ault, France)

Weston, M.A. 1998. An observation of aggression between a Hooded Plover and a Black-fronted Dotterel. *Aust. Bird Watcher* 17: 301–302. (Dept. Zool., Univ. Melbourne, Parkville, Vic. 3052, Australia). *Charadrius rubricollis*, *C. melanops*.

Distribution & faunistics

Adriaens, P. & Bekaert, L. 1997. [Least Sandpiper at Gent in August 1996]. *Dutch Birding* 19: 217–221. In Dutch with English summary. *Calidris minutilla*, Belgium.

Agee, A. 1997. Killdeer hatches in Snowy Plover nest. *West. Birds* 28: 115. (10117 Ranchwood Manor Dr., Oklahoma City, OK 73139, USA). *Charadrius vociferus*, *Charadrius alexandrinus*.

Andrews, I. 1997. The Western Sandpiper in Lothian. *Birding World* 10: 302–304. *Calidris mauri*.

Baron, N. 1999. Riders on the storm. *Can. geographic* 119(5): 49–56. *Calidris mauri*, migration strategies.

Bennett, A.J. 1996. Pacific and American golden-plovers breeding in southwestern Alaska. *Northwest. Nat.* 77: 49–51. (Lake Clark Natl. Pk. Pres., P.O. Box 2643, Kenai, AK 99611–2643, USA). *Pluvialis dominica*, *P. fulva*, range extension.

Berrevoets, C.M., Strucker, R.C.W. & Meininger, P.L. 1999. *Watervogels in de Zoute Delta 1997/98*. [Waterbirds in the salt waters of the Delta area, SW-Netherlands, 1997/98]. Middelburg: Rijksinstituut voor Kust en Zee RIKZ, 80 p. In Dutch with English summary. (RIKZ, P.O. Box 8039, NL-4330 EA Middelburg, Netherlands). Results of monthly counts of waterbirds in the southern Delta area, SW-Netherlands. ISBN 90-369-344-35, Report RIKZ 99.001. Giving detailed information on the occurrence of many wader species in the Delta area.

Bishop, M.A. & Warnock, N. 1998. Migration of Western sandpipers: links between their Alaskan stopover areas and breeding grounds. *Wilson Bull.*

- 110(4): 457–462. *Calidris mauri*.
- Borrow, N. 1997. Red-crested Bustard *Eupodotis ruficrista* and Adamawa Turtle Dove *Streptopelia hypopyrrha*, new to the Gambia, and sightings of Great Snipe *Gallinago media*. *Malimbus* 19: 36–38. (Flat 5, 63–67 St. George's Dr., Pimlico, London SW1V 4DD, UK)
- Bourne, W.R.P. 1997. Phalaropes in the Arabian Sea & Gulf of Oman. *Oman Bird News* 20: 56. *Phalaropus lobatus* and *Phalaropus fulicarius*
- Buchanan, J.B. & Evenson, J.R. 1997. Abundance of shorebirds at Willapa Bay, Washington. *West. Birds* 28: 158–168. (Cascadia Res. Coll., 218 W. Fourth Ave., Olympia, WA 98501, USA). *Calidris mauri*, *C. alpina*, *Limnodromus griseus* most numerous species.
- Carter, M.A. & Rogers, D. 1998. Little Ringed Plover *Charadrius dubius*: a Kimberley record with comments on morphological and vocal aspects of eastern populations. *Aust. Bird Watcher* 17: 269–277. (30 Canadian Bay Rd., Mt Eliza, VIC 3930, Australia).
- Cleeves, T. 1998. The Slender-billed Curlew in Northumberland – a new British record. *Birding World* 11: 181–191. *Numenius tenuirostris*.
- Devos, K., De Scheemaeker, F., Allein, S. & Rapp, G. 1998. Resultaten van steltloperstellingen langs de Vlaamse kust, winter 1996/97. (Wader counts along the Belgian coast during the winter 1996/97). *Mergus* 12(4): 187–199. In Dutch with English and French summary. (Inst. Natuurbehoud, Kliniekstr. 25, B-1070 Brussels, Belgium)
- Devos, K., De Scheemaeker, F., Allein, S. & Rapp, G. 1997. Resultaten van steltloperstellingen langs de Vlaamse kust, winter 1995/1996. (Wader counts along the Belgian coast during the winter 1995/1996). *Mergus* 11. In Dutch with English and French summary.
- Devos, K., De Scheemaeker, F., Allein, S. & Rapp, G. 1996. Resultaten van steltloperstellingen langs de Vlaamse kust, winter 1994/1995. (Wader counts along the Belgian coast during the winter 1994/1995). *Mergus* 10: 187–198. In Dutch with English and French summary.
- Devos, K., De Scheemaeker, F., Allein, S. 1994. Resultaten van steltloperstellingen langs de Vlaamse kust, winter 1993/1994. (Wader counts along the Belgian coast during the winter 1993/1994). *Mergus* 8: 242–246. In Dutch with English and French summary.
- Van Dijk, A.J., Boele, A., Zoetebier, D. & Meijer, R. 1999. Kolonievogels en zeldzame broedvogels in Nederland in 1996. (Rare and colonial breeding birds in The Netherlands in 1996). *Limosa* 72(1): 23–36. In Dutch with English summary. (Rijksstraatweg 178, NL-6573 DG Beek-Ubbergen, Netherlands). Breeding waders: *Recurvirostra avosetta*, *Charadrius dubius*, *C. hiaticula*, *C. alexandrinus*, *Calidris alpina*, *Philomachus pugnax*, *Actitis hypoleucos*.
- Dougall, T.W., Mee, A. & Yalden, D.W. 1999. Recent fluctuations in a Common Sandpiper breeding population. *Scott. Birds* 20(1): 44–45. (62 Leamington Terrace, Edinburgh EH10 4JL, Scotland). *Actitis hypoleucos*, no indication of an overall decline or increase in the population in the Moorfoot Hills, SE Scotland, between 1993–1998.
- Duckworth, J.W., Timmins, R.J. & Evans, T.D. 1998. The conservation status of the River Lapwing *Vanellus duvaucelii* in southern Laos. *Biol. Conserv.* 84(3): 215–222. (East Redham Farm, Pilning, Bristol BS12 3JG, UK). Southern Lao population threatened by hydropower development and human pressure.
- Eckert, C. & Sinclair, P. 1997. Little Stint: a Yukon first at Judas Creek. *Birders J.* 6: 183–185. (153 Grange Hall Rd., Beavercreek, OH 45430, USA). *Calidris minuta*.
- Ecsedi, Z. & Oláh, J. Jr. 1997. Spoon-billed Sandpipers in Vietnam. *Birding World* 10: 261. *Eurynorhynchus pygmeus*, Red River Delta.
- Edwards, T. 1997. The Lesser Sand Plover in Sussex – a new British record. *Birding World* 10: 294–297. *Charadrius mongolus*.
- Fernandez, G., Carmona, R. & de la Cueva, H. 1998. Abundance and seasonal variation of Western Sandpipers (*Calidris mauri*) in Baja California Sur, Mexico. *Southwest Nat.* 43: 57–61. (H. de la Cueva: P.O. Box 434844, San Diego, CA 92143–4844, USA). Most numerous wader during autumn migration and winter.
- Gantlett, S. 1997. The Spur-winged Plover in Kent – a new British bird?. *Birding World* 10: 217–219. *Vanellus spinosus*, UK and Netherlands. Escape??
- Van Gastel, A.J.G. & Van Gastel, E.R. 1999. Cream-coloured Courser *Cursorius cursor*, new for Ghana. *Malimbus* 21(1): 54–55.
- Gerdes, K., Kruckenberg, H., Reuter, M. & Voss, E. 1998. Zur Brutvogelwelt des Ems-Ästuars zwischen Emden und Leer. (Breeding birds in the Ems estuary). *Vogelkd. Ber. Niedersachs.* 30(1): 19–32. In German with English summary. (Mozartstrasse 20, D-26789 Leer, Germany). High breeding numbers of *Recurvirostra avosetta*, *Tringa totanus*, *Vanellus vanellus*, *Limosa limosa*.
- Gibson, D.D. & Kessel, B. 1997. Inventory of the species and subspecies of Alaska birds. *West. Birds* 28: 45–95. (Univ. Alaska Mus., Fairbanks, AK 99775–6960, USA). 448 species and 110 subspecies.
- Gromadzka, J. & Serra, L. 1998. Differential migration of juvenile and adult Grey Plovers *Pluvialis squatarola* at the mouth of the Vistula River, Poland. *Ornis Fenn.* 75(4): 193–199.
- Jürgens, R. 1998. 5 Stelzenläufer *Himantopus himantopus* bei Schöppenstedt beobachtet. [5 Black-winged Stilts *Himantopus himantopus* near Schöppenstedt observed]. *Vogelkd. Ber. Niedersachs.* 30(2): 109–110. In German. (Am Krähenfelde 7, D-38170 Schöppenstedt, Germany).
- Kapanen, M. 1997. Finnish birds 5. Broad-billed Sandpiper. *Alula* 3: 54–55. *Limicola falcinellus* in Finland.
- Keijl, G. 1999. Trek van Bonte Strandlopers in het voorjaar bij Castricum. [Migration of Dunlins at Castricum in spring]. *Graspieper* 19(1): 14–17. In Dutch. Spring migration of *Calidris alpina* along the North Sea coast of the Netherlands.
- Knaus, P. & Balzari, C' A. 1999. Seltene Vogelarten und ungewöhnliche Vogelbeobachtungen in der Schweiz im Jahre 1998. (Rare and unusual bird observations in Switzerland in 1998).

- Ornithol. Beob.* 96(3): 157–182. In German with English summary. (Pflanzschulstr. 49, CH–8004 Z_rich, Switzerland, e-mail: p.knaus@hotmail.com). Waders: *Charadrius hiaticula*, *Pluvialis squatarola*, *Calidris melanotos*, *Limicola falcinellus*, *Lymnocyptes minimus*, *Numenius phaeopus*, *Tringa stagnatilis*, *T. glareola*, *Phalaropus lobatus*, *Ph. fulvicarius*.
- Lakin, L. & Rylands, K. 1997. The Semipalmated Plover in Devon : the second British record. *Birding World* 10: 212–216. *Charadrius semipalmatus*
- Meier–Zwicky, C. 1999. Zweiter Schweizer Brutnachweis des Mornellregenpfeifers *Charadrius morinellus* L. in Graubünden. (Second breeding record of the Dotterel *Charadrius morinellus* in Switzerland). *Ornithol. Beob.* 96(1): 62–63. In German with English summary.
- Meissner, W. & Wlodarczak, A. 1998. (Spring migration of waders *Charadrii* in the area of the protected "Rzeczne taki" reserve at the Bay of Puck). *Notatki Ornitol.* 39: 219–229. In Polish with English summary. (Katedra Ekol. Zool. Kregowcow, Univ. Gdansk, Al. Legionow 9, P–80–441 Gdansk, Poland). Information about spring migration numbers of 25 wader species. Most numerous: *Tringa glareola*, *Philomachus pugnax*, *Gallinago gallinago*, *Vanellus vanellus*, *Calidris alpina*.
- Nankinov, D., Shurulinkov, P., Nikolov, I., Nikolov, B., Dalaktchieva, S., Hristov, I., Stanchev, R., Rogev, A., Dutsov, A. & Sarov, M. 1998. Studies of the waders (*Charadriiformes*) on the wetlands around Sofia (Bulgaria). *Riv. ital. Orn.* 68(1): 63–68. (Int. Zool., BAS 1000 Sofia, Bulgaria) Data about 40 wader species, migration as well as breeding.
- North, M.R., Dewhurst, D. & Tucker, S.S. 1996. First nest discovered for Alaska subspecies of Marbled Godwit. *Northwest Nat.* 77: 17–18. (USFWS, West. Alaska Ecol. Serv., 605 W. 4th Ave., Anchorage, AK 99501, USA). *Limosa fedoa beringiae*.
- Ogilvie, M. and the Rare Breeding Birds Panel. 1999. Rare breeding birds in the United Kingdom in 1996. *Brit. Birds* 92(3): 120–154. (Glencairn, Bruichladdich, Isle of Islay PA49 7UN, UK). Breeding waders: *Himantopus*, *Recurvirostra avosetta*, *Burhinus oedicephalus*, *Charadrius dubius*, *Calidris morinellus*, *Calidris temminckii*, *Calidris maritima*, *Philomachus pugnax*, *Limosa limosa*, *Numenius phaeopus*, *Tringa nebularia*, *Tringa ochropus*, *Tringa glareola*, *Phalaropus lobatus*.
- Ogilvie, M. and the Rare Breeding Birds Panel. 1999. Rare breeding birds in the United Kingdom in 1997. *Brit. Birds* 92(8): 389–428. Breeding waders: *Recurvirostra avosetta*, *Burhinus oedicephalus*, *Charadrius dubius*, *Calidris morinellus*, *Calidris temminckii*, *Calidris maritima*, *Philomachus pugnax*, *Limosa limosa*, *Numenius phaeopus*, *Tringa nebularia*, *Tringa ochropus*, *Tringa glareola*, *Phalaropus lobatus*.
- Page, G.W. ...[et al.] 1997. Numbers of wintering shorebirds in coastal wetlands of Baja California, Mexico. *J. Field Ornithol.* 68: 562–574. Most abundant species were *Calidris mauri*, *Limosa lapponica*.
- Page, G.W., Stenzel, L.E. & Kjelson, J.E. 1999. Overview of shorebird abundance and distribution in wetlands of the Pacific coast of the contiguous United States. *Condor* 101(3): 461–471. (Point Reyes Bird Observ., 4990 Shoreline Highway, Stinson Beach, CA 94970, USA, e-mail: gpage@prbo.org). 43 species recorded. *Calidris mauri* and *C. alpina* most numerous in all wetlands.
- Pepping, M., Piersma, T., Pearson, G. & Lavaleye, M. (eds.) 1999. *Intertidal sediments and benthic animals of Roebuck Bay, Western Australia*. NIOZ report 1999–3. Den Burg, Texel : Netherlands Inst. Sea Res. [etc.], 211 p. (NIOZ, P.O. Box 59, NL–1790 AB Den Burg, Texel, Netherlands). Detailed distribution maps of feeding waders.
- Rehfish, M. & Austin, G. 1999. Ringed Plovers go east. *BTO News* 223: 14–15. (BTO, The Nunnery, Thetford, Norfolk IP24 2PU, UK). *Charadrius hiaticula*: shift in centres of distribution may be associated with changing weather patterns.
- Reichholf, J. 1994. Die Wasservogel am unteren Inn. Ergebnisse von 25 Jahren Wasservogelzählung: Dynamik der Durchzugs- und Winterbestände, Trends und Ursachen. [Waterbirds at the lower Inn. Results of 25 years counting: dynamics of migration and resting numbers, trends and explanation.]. *Mitt. zool. Ges. Braunau* 6: 1–92. In German.
- Rogers, M.J. & Rarities Committee 1999. Report on rare birds in Great Britain in 1998. *Brit. Birds* 92(11): 554–609. Information on 25 wader species. Photographs of *Glareola pratincola*, adult *Charadrius mongolus*, adult *Pluvialis dominica*, juvenile *Gallinago media*, first winter *Tringa flavipes*, and *Xenus cinereus*.
- Rohde, J. 1998. *Kampfläufer in Schleswig–Holstein: ein Tagebuch aus den Jahren 1986–97*. [Ruffs in Schleswig–Holstein: a diary from the years 1986–97]. 207 p. In German with English summary. (Oskar–Alexanderstr. 20, D–24576 Bad Bramstedt, Germany). *Philomachus pugnax*.
- De Smet, G. 1997. [Slender–billed Curlew collected at Canisvliet in September 1896]. *Dutch Birding* 19: 230–232. In Dutch with English summary. *Numenius tenuirostris*.
- Sueur, F. 1996. [Ornithological observations in Madagascar]. *Alauda* 64: 435–442. In French with English summary. First record for *Calidris canutus*.
- Sueur, F. 1998. Synthèse ornithologique picarde 1992. [Summary of ornithological records in the Picardy in 1992]. *Avifaune Picarde* 6: 1–49. In French. (9 rue du Champ neuf, Le Bout des Crocs, F–80120 Saint–Quentin–en–Tourmont, France). Information on occurrence and breeding numbers of 31 wader species.
- Sueur, F., Carruette, P. & Triplet, P. 1998. Actes ornithologiques 1995 de la Réserve Naturelle de la Baie de Somme. [Ornithological records of 1995 at the nature reserve Baie de Somme]. *Avifaune Picarde* 6: 51–74. In French. Information on the occurrence and breeding numbers of 34 wader species.
- Sueur, F. & Triplet, P. 1998. Evolution de l'ensablement de la baie de Somme et avenir des stationnements de quelques Anatidés et Limicoles. [Silting up of the Somme bay and the future of staging ducks and shorebirds]. *Avifaune Picarde* 6: 75–88. In French. Most frequently observed waders: *Haematopus ostralegus*, *Charadrius hiaticula*, *Pluvialis squatarola*, *Calidris canutus*, *C. alpina*, *Numenius arquata*, *Tringa totanus*. Data on numbers from

1970–1997.

- Sueur, F., Carruette, P. & Triplet, P. 1999. Actes ornithologiques 1996 de la Réserve Naturelle de la Baie de Somme. [Ornithological records of 1996 at the nature reserve Baie de Somme]. *Avifaune Picarde 7*: 1–31. In French. Information on the occurrence and breeding numbers of 34 wader species.
- Sueur, F., Carruette, P. & Triplet, P. 1999. Actes ornithologiques 1997 de la Réserve Naturelle de la Baie de Somme. [Ornithological records of 1997 at the nature reserve Baie de Somme]. *Avifaune Picarde 7*: 33–60. In French. Information on the occurrence and breeding numbers of 35 wader species.
- Sueur, F., Carruette, P. & Triplet, P. 1999. Actes ornithologiques 1998 de la Réserve Naturelle de la Baie de Somme. [Ornithological records of 1998 at the nature reserve Baie de Somme]. *Avifaune Picarde 7*: 61–91. In French. Information on the occurrence and breeding numbers of 33 wader species.
- Sueur, F. 1999. Actes ornithologiques 1998 de la réserve Authie–Somme. [Ornithological records of 1998 at the reserve Authie–Somme]. *Avifaune Picarde 7*: 109–115. In French. Information on the occurrence of 12 wader species.
- Tiedemann, R. 1999. Seasonal changes in the breeding origin of migrating Dunlins *Calidris alpina* as revealed by mitochondrial DNA sequencing. *J. Ornithol.* 140(3): 319–323. (Inst. Haustierkunde, Christian–Albrechts–Univ. Kiel, Olshausenstr. 40, D–24118 Kiel, Germany, e–mail: rtiedemann@ifh.uni-kiel.de)
- Tourenq, C. ...[et al.] 1996. [Ornithological observations in south–west Mongolia]. *Alauda* 64: 355–362, 405–412. In French with English summary. (INRA, CRA, Toulouse, B.P. 27, F–31326 Castanet Tolosan Cedex, France). First breeding record for *Himantopus himantopus*.
- Tréca, B. 1997. [Ruff *Philomachus pugnax* in Senegal]. *Alauda* 65: 161–166. In French with English summary. (Lab. Ornithol., ORSTOM, B.P. 1386, Dakar, Senegal)
- Triplet, P., Robert, J.–C., Sueur, F. & Violet, F. 1999. Chronique ornithologique du Hable d'Ault (Somme): année 1998. [Ornithological records for Hable d'Ault: 1998]. *Avifaune Picarde 7*: 93–108. In French. (SMACOPI, Réserve Naturelle Baie de Somme, Stn. Biol. Blanquetaque, 1 place de l'Amiral Courbet, F–80100 Abbeville, France, e–mail: blanquetaque@dyadel.net) Information on the occurrence and breeding of 25 wader species.
- Tyler, S.J. & Tyler, L. 1997. Early nesting of Red–winged Pratincoles *Glareola pratincola* and some other records for the Boteti River [Botswana]. *Babbler* 32: 40. (Room 106, D.A.H.P., Private Bag 68–908, Gaborone, Botswana)
- Votier, S. 1997. Ageing of Collared Pratincoles. *Birding World* 10: 460. *Glareola pratincola*.
- Warnock, N., Haig, S.M. & Oring, L.W. 1998. Monitoring species richness and abundance of shorebirds in the western Great Basin. *Condor* 100: 589–600.
- Whitfield, D.P. 1998. Methods for ageing Dotterel *Charadrius morinellus*. *Ring. & Migr.* 19(3): 200–204. (Scottish Nat. Heritage, Advisory Serv., 2 Anderson Place, Edinburgh EH6 5NP, UK)
- Wilson, A. 1999. Tumbling Lapwing populations. *BTO News* 224: 14. (BTO, The Nunnery, Thetford, Norfolk IP24 2PU, UK). Population estimates for *Vanellus vanellus* in 1998.
- Zöckler, C. 1998. Patterns in biodiversity in Arctic birds. *WCMC Biodiversity Bull.* 3: 1–15. (World Conservation Monitoring Centre, 219 Huntingdon Rd., Cambridge CB3 0DL, UK, fax: +44 (0)1223 277136). Breeding distribution and population size of 20 *Calidris* waders and protected areas in the circumpolar Arctic region.

Miscellaneous

Fadat, C. 1996. [Proposal for a management plan for hunting Woodcock *Scolopax rusticola* in Europe]. *Alauda* 64: 33–44. In French. (1 rue Jean Rostand, F–34800 Clermont–l'H., rault, France)

Rattner, B.A., Capizzi, J.L. & Melancon, M.J. 1995. Exposure and effects of oilfield Brine discharges on Western Sandpipers (*Calidris mauri*) in Nueces Bay, Texas. *Bull. Environ.*

Contam. Toxicol. 54(5): 683–689.

Schmitz, M. & Osing, H. 1999. Auswirkungen der Errichtung von Wupper– und Grosser Dhünn–Talsperre auf die Avifauna. Mit Anmerkungen zur Funktion der Ausgleichs– und Ersatzmassnahmen. (Effect of the creation of the Wuppertalsperre and Grosse Dhünn–Talsperre reservoirs on the local avifauna. With comments on the effectiveness of habitat–loss –compensation and habitat creation measures). *Charadrius* 35(2): 41–60. In German with English summary. (Kölner Strasse 55, D–45145 Essen, Germany, e–mail: michael.schmitz@uni-essen.de). *Charadrius dubius*, increase in breeding pairs at first. Large numbers of waders during the first phase of construction.

Steendam, O. 1998. Weidevogelbescherming in Noord–Holland. [Meadowbird protection in Noord–Holland]. *Graspieper* 18(4): 133–139. In Dutch. (Landschapsbeheer Noord–Holland, P.O. Box 205, NL–2050 AE Overveen, Netherlands). Breeding waders: *Haematopus ostralegus*, *Vanellus vanellus*, *Limosa limosa*, *Tringa totanus*, *Philomachus pugnax*, *Gallinago gallinago*, *Recurvirostra avosetta*.

Summers, R.W. & Etheridge, B. 1998. Rates of wear of incoloy and stainless steel rings on Turnstones *Arenaria interpres*. *Ring. & Migr.* 19(2): 81–85. (Lismore, Mill Crescent, North Kessock, Inverness IV1 1XY, UK).

Taylor, L.A. & Mackay, M.F. 1997. *The effects of the 'Iron Baron' oilspill on shorebirds*. Johnstone Centre Report No. 110. Albury, N.S.W. : Charles Sturt Univ.

Tipper, R. 1998. The world's best wader–watching – and much more: birding in Hong Kong. *Birding World* 11: 138–148.