

Recent publications on waders 78

COMPILED BY HENK KOFFIJBERG

Glenn Millerweg 102, NL-1311 RW Almere, The Netherlands, e-mail: koffijberg@wanadoo.nl

In trying to make these lists as complete as possible, and the next one even more complete and up to date, your help is necessary!! Please send new titles, abstracts, reprints, copies, reports, and (important too!) omissions or corrections to: **Henk Koffijberg, Glenn Millerweg 102, NL-1311 RW Almere, The Netherlands, e-mail: koffijberg@wanadoo.nl.**

Remember to include, if possible, the address of the first author, and translations of titles if they are in other languages than English.

For help with this issue I am very grateful to J. Melter, T. Gunnarsson, W. Meissner, M. Boschert, H. Insley.

Translated titles from languages other than English are bracketed; with round brackets if the translation is by the author(s), square brackets if not.

N.B. a copy of your paper or abstract would be very helpful to enable me to add some keywords or give added annotation to the titles.

BREEDING

Adams, N.J., Kerley, G.I.H. & Watson, J.J. 1999. Disturbance of incubating African Black Oystercatchers: is heating of exposed eggs a problem? *Ostrich* 70: 225–228. *Haematopus moquini*, overheating of eggs doesn't seem to influence breeding success.

Amat, J.A., Fraga, R.M. & Arroyo, G.M. 2001. Variations in body condition and egg characteristics of female Kentish Plovers *Charadrius alexandrinus*. *Ardea* 89(2): 293–299. (Biol. Stn. Doñana, CSIC, Apartado 1056, E-41080 Sevilla, Spain, e-mail: aguilar@cica.es)

Amat, J.A., Fraga, R.M. & Arroyo, G.M. 2001. Intraclutch egg-size variation and offspring survival in the Kentish Plover *Charadrius alexandrinus*. *Ibis* 143: 17–27.

Amat, J.A., Visser, G.H., Pérez-Hurtado, A. & Arroyo, G.M. 2000. Brood desertion by female shorebirds: a test of the differential parental capacity hypothesis on Kentish Plovers. *Proc. R. Soc. Lond. B* 267: 2171–2176. *Charadrius alexandrinus*.

Blomqvist, D. & Johansson, O.C. 2000. A comment on successive clutches and breeding system in Lapwings *Vanellus vanellus*. *Ibis* 142(2): 323.

Byrkjedal, I., Gronstol, G.B., Hafsmo, J.E. & Lislewand, T. 2000. Chick punishment and chick adoption in Northern Lapwings. *Ornis Fenn.* 77(2): 89–92. *Vanellus vanellus*.

Cotter, P.A. & Andres, B.A. 2000. Nest density of shorebirds from the Beaufort Sea coast, Alaska. *Can. Field-Nat.* 114: 287–291.

Currie, D., Valkama, J., Berg, A., Boschert, M., Norrdahl, K., Hänninen, M., Korpimäki, E., Pöyri, V. & Hemminki, O. 2001. Sex roles, parental effort and offspring desertion in the monogamous Eurasian Curlew *Numenius arquata*. *Ibis* 143: 642–650. (Sect. Ecol., Dept. Biol., Univ. Turku, FIN-20014 Turku, Finland)

Davis, W.E.D. Jr. & Recher, H.F. 2000. False-brooding behaviour in the Red-capped Plover *Charadrius ruficapillus*. *Aust. Bird Watcher* 18: 248–249. (Coll. Gen. Stud., Boston Univ., Boston, MA 02215, USA)

Duncan, A., Duncan, R., Rae, R., Rebecca, G.W. & Stewart, B.J. 2001. Roof and ground nesting Oystercatchers in Aberdeen. *Scott. Birds* 22: 1–8. *Haematopus ostralegus* population in Aberdeen in 1993 of at least 275 pairs of which 205 nested on roofs. Productivity for a sample of roof nesting birds was 0.8 fledged young per pair.

French, P., Insley, H., Siriwardena, G. & Buxton, N. 2000. The breeding success of a population of Lapwings in part of Strathspey 1996–1998. *Scott. Birds* 21: 98–108. (e-mail Insley: hugh.insley@tinyworld.co.uk). 630 Nesting attempts of *Vanellus vanellus* were monitored. Mean clutch size (3.6) and brood size at hatching (3.0) remained constant over the study period. Nest failure rate didn't vary significantly between years and nest survival from laying to hatching was about 60%. However, there were highly significant differences in the daily failure rates for nests across the 8 study sites used. Combining all 8 sites, survival of chicks from hatching to fledging varied between 9.5% and 17.5%.

Handel, C.M. & Gill Jr., R.E. 2000. Mate fidelity and breeding site tenacity in a monogamous sandpiper, the Black Turnstone. *Anim. Behav.* 60: 471–481. *Arenaria melanocephala*.

Larsen, T. 2000. Influence of rodent density on nesting associations involving the Bar-tailed Godwit *Limosa lapponica*. *Ibis* 142: 476–481.

Mabee, T.J. & Estelle, V.B. 2000. Assessing the effectiveness of predator exclosures for plovers. *Wilson Bull.* 112(1): 14–20. (ABR Inc., PO Box 249, Forest Grove, OR 97116, USA). *Charadrius melanodus*, *C. alexandrinus*, *C. vociferus*.

Parish, D.M.B., Thompson, P.S. & Coulson, J.C. 2001. Effects of age, cohort and individual on breeding performance in the Lapwing *Vanellus vanellus*. *Ibis* 143(2): 288–295.

Powell, A.N. & Powell, A.N. 2001. Habitat characteristics and nest success of Snowy Plovers associated with California Least Tern colonies. *Condor* 103(4): 785–792. (USGS, West. Ecol. Res. Ctr., Dept. Biol., San Diego, CA 92182, USA). *Charadrius alexandrinus*.



- Reed, J.M., Gray, E.M., Lewis, D., Oring, L.W., Coleman, R., Burr, T. & Luscomb, P.** 1999. Growth patterns of Hawaiian stilt chicks. *Wilson Bull.* 111(4): 478–487. *Himantopus mexicanus knudseni*.
- Rusticali, R., Scarton, F. & Valle, R.** 1999. Habitat selection and hatching success of Eurasian Oystercatchers in relation to nesting Yellow-legged Gulls and human presence. *Waterbirds* 22: 367–375. (Valle: Castello 618/E, I-30122, Venezia, Italy, e-mail: robevalle@tin.it). *Haematopus ostralegus*.
- Schmechel, F.** 2001. *Aspects of habitat selection, population dynamics, and breeding biology of the endangered Chatham Island oystercatcher (Haematopus chathamensis)*. PhD thesis, Lincoln Univ. (Ecol. & Entomol. Grp., PO Box 84, Lincoln Univ., Canterbury, New Zealand, e-mail: Schmechf@lincoln.ac.nz).
- Schreiber, M.** 2001. Verbreitung und Bruterfolg des Kiebitzes *Vanellus vanellus* im südwestlichen Niedersachsen in Abhängigkeit von ausgewählten bodenkundlichen Parametern und landwirtschaftlicher Nutzung. (Distribution and breeding success of the Lapwing *Vanellus vanellus* in south-west Lower Saxony in relation to soil types and agricultural land use). *Vogelwelt* 122(2): 55–65. In German with English summary. (Blankenburger Str. 34, D-49565 Bramsche/Epe, Germany, e-mail: Schreiber.Umwelplanung@t-online.de)
- Valkama, J. & Currie, D.** 1999. Low productivity of Curlews *Numenius arquata* on farmland in southern Finland: causes and consequences. *Ornis Fenn.* 76: 65–70.
- Winton, B.R., Leslie Jr., D.M. & Rupert, J.R.** 2000. Breeding ecology and management of Snowy Plovers in North-Central Oklahoma. *J. Field Ornithol.* 71: 573–584. *Charadrius alexandrinus*.

BEHAVIOUR, ECOLOGY AND EVOLUTION

- Arcas, J.** 2001. Body weight variation and fat deposition in Common Sandpipers *Actitis hypoleucos* L. during their autumn migration in the Ria de Vigo, Galicia, north-west Spain. *Ring. & Migr.* 20(3): 216–220. (Lab. Anatomia Anim., Dept. Ecol. Biol. Anim., Fac. Sci., Univ. Vigo, Apartado 874, E-36200 Vigo (Pontevedra), Spain)
- Ausden, M.** 2001. The effects of flooding of grassland on food supply for breeding waders. *Brit. Wildl.* 12(3): 179–187.
- Battley, P.F., Piersma, T., Dietz, M.W., Tang, S., Dekkinga, A. & Hulsman, K.** 2000. Empirical evidence for differential organ reductions during trans-oceanic bird flight. *Proc. R. Soc. Lond. B* 267: R1795–R1804. (Aust. Sch. Environm. Stud., Griffith Univ., Nathan, Qld. 4111, Australia).
- Boschert, M.** 1999. Bestandsentwicklung des Kiebitzes nach partieller Wiedervernässung und Extensivierung. Untersuchungen in drei Gebieten der Oberrheinebene. (Population development of the Lapwing (*Vanellus vanellus*) after re-irrigation and changes in grassland management in the Upper Rhine Valley, Baden-Württemberg). *Natur-schutz u. Landschaftsplanung* 31(2): 51–57. In German with English summary. (BIOPLAN, Inst. angewandte Biol. & Planung GbR, Nelkenstr. 10, D-77815 Bühl, Germany).
- Boyd, H. & Piersma, T.** 2001. Changing balance between survival and recruitment explains population trends in Red Knots *Calidris canutus islandica* wintering in Brit-
- ain, 1969–1995. *Ardea* 89(2): 301–317. (Neth. Inst. Sea Res. NIOZ, PO Box 59, NL-1790 AB Den Burg, Texel, Netherlands, e-mail: theunis@nioz.nl).
- Cabral, J.A., Pardal, M.A. & Marques, J.C.** 1999. The impact of macroalgal blooms on the use of the intertidal area and feeding behaviour of waders (*Charadrii*) in the Mondego estuary (west Portugal). *Acta oecologica* 20(4): 417–427.
- Collins, P. & Jessop, R.** 2001. Arrival and departure dates and habitat of Little Curlew *Numenius minutus* at Broome, north-western Australia. *Stilt* 39: 10–12. (RMB 4009, Cowes, Vic. 3922, Australia, e-mail: moonbird@waterfront.net.au)
- De Nobel, W.T., Diadicheva, E.A., Nilsen, S.O. & Kirikova, T.** 2001. *Broad-billed Sandpiper in the Sivash, August 1998*. In: Van der Winden, J., E.A. Diadicheva, W.T. de Nobel & M.W.J. van Roomen (eds.). Counts and ecology of waterbirds in the Sivash, Ukraine, August 1998. – WIWO Report 71. – WIWO: Zeist, 2001, p. 61–71. Numbers, distribution, age, moult, biometrics, weight and feeding ecology of *Limicola falcinellus*.
- Diadicheva, E.A., Chernichko, R.N., Vinokurova, S.V. & De Nobel, W.T.** 2001. *Biometry and moult of waders in the Sivash, August 1998*. In: Van der Winden, J., E.A. Diadicheva, W.T. de Nobel & M.W.J. van Roomen (eds.). Counts and ecology of waterbirds in the Sivash, Ukraine, August 1998. – WIWO Report 71. – WIWO: Zeist, 2001, p. 39–59. Data on: *Glareola pratincola*, *Calidris minuta*, *C. ferruginea*, *C. alpina*, *Philomachus pugnax*, *Gallinago gallinago*, *Tringa stagnatilis*, *T. nebularia*, *T. glareola*.
- Drake, K.R., Thompson, J.E., Drake, K.L. & Zonick, C.** 2001. Movements, habitat use, and survival of nonbreeding Piping Plovers. *Condor* 103(2): 259–267. (Caesar Kleberg Wildl. Res. Inst., Texas A&M Univ.-Kingsville, Campus Box 218, Kingsville, TX 78363, USA). *Charadrius melanotos*.
- Durell, S.E.A. Le V. dit, Goss-Custard, J.D., Caldow, R.W.G., Malcolm, H.M. & Osborn, D.** 2001. Sex, diet and feeding method-related differences in body condition in the Oystercatcher *Haematopus ostralegus*. *Ibis* 143: 107–119.
- Elias, S.P., Fraser, J.D. & Buckley, P.A.** 2000. Piping Plover brood foraging ecology on New York barrier islands. *J. Wildl. Manage.* 64: 346–254. *Charadrius melanotos*.
- Green, R.E., Tyler, G.A. & Bowden, C.G.R.** 2000. Habitat selection, ranging behaviour and diet of the stone curlew (*Burhinus oedicnemus*) in southern England. *J. Zool. (Lond.)* 250: 161–183. (e-mail R.E. Green: reg29@hermes.cam.ac.uk).
- Gronstol, G.B., Byrkjedal, I., Hafsmo, J.E. & Lisllevand, T.** 1999. Northern Lapwing *Vanellus vanellus* copulating with a grass turf. *Ornis Norvegica* 22(1/2): 60–62.
- Heg, D., Ens, B.J., Van der Jeugd, H.P. & Bruinzeel, L.W.** 2000. Local dominance and territorial settlement of nonbreeding Oystercatchers. *Behaviour* 137: 473–530. (Zool. Lab., Univ. Groningen, Kerklaan 30, 9751 NN Haren, Netherlands) *Haematopus ostralegus*.
- Hodder, K.H.** 2000. Foraging behaviour of Eastern Curlews *Numenius madagascariensis* remaining in Roebuck Bay after the breeding migration. *Aust. Bird Watcher* 18: 183–191. (CEH Dorset, Winfrith Technology Centre, Dorchester, Dorset DT2 8ZD, UK)
- Jackson, D.B. & Green, R.E.** 2000. The importance of the introduced hedgehog (*Erinaceus europaeus*) as a preda-



- tor of the eggs of waders (*Charadrii*) on machair in South Uist, Scotland. *Biol. Conserv.* 93(3): 333–348. (e-mail R.E. Green: reg29@hermes.cam.ac.uk).
- Jessop, R. & Collins, P.** 2001. Raptor attack methods and the response of Little Curlew *Numenius minutus* to disturbance at Broome, north-western Australia. *Stilt* 39: 61–64. (Philip Island Nature Park, PO Box 97, Cowes, Vic. 3922, Australia, e-mail: rjessop@penguins.org.au).
- Jukema, J., Wymenga, E. & Piersma, T.** 2001. Opvatten en ruien in de zuidwesthoek: Kemphanen *Philomachus pugnax* op voorjaarstrek in Friesland. (Stopping over in SW Friesland: fattening and moulting Ruffs *Philomachus pugnax* during northward migration in The Netherlands). *Limosa* 74(1): 17–26. In Dutch with English summary. (Heardawei 44, NL-8854 AC Oosterbierum, Netherlands).
- Kvist, A. & Lindström, A.** 2001. Basal metabolic rate in migratory waders: intra-individual, intraspecific, interspecific and seasonal variation. *Funct. Ecol.* 15(4): 465–473.
- Lewis, L.J. & Kelly, T.C.** 2001. A short-term study of the effects of algal mats on the distribution and behavioural ecology of estuarine birds. *Bird Study* 48(3): 354–360. (Dept. Zool. Anim. Ecol., Natl. Univ. Ireland, Lee Maltings, Prospect Rowe, Cork, Ireland). Effects on *Limosa limosa* and *Tringa totanus*.
- Masero, J.A., Perez-Gonzalez, M. & Otero-Saavedra, M.** 1999. Food supply for waders (Aves: *Charadrii*) in an estuarine area in the Bay of Cadiz (SW Iberian Peninsula). *Acta oecologica* 20(4): 429–434.
- Masero, J.A. & Pérez-Hurtado, A.** 2001. Importance of supratidal habitats for maintaining overwintering shorebird populations: how Redshanks use tidal mudflats and adjacent saltworks in southern Europe. *Condor* 103(1): 21–30. (Dept. Biol. Anim., Veg. & Ecol., Fac. Cienc. Mar., Univ. Cadiz, 11510 Puerto Real, Spain). *Tringa totanus*.
- McCloskey, J.T. & Thompson, J.E.** 2000. Sex-related differences in migration chronology and winter habitat use of Common Snipe. *Wilson Bull.* 112: 143–148. (West Virginia Univ., For. Div., PO Box 6125, Morgantown, WV 26506, USA) *Gallinago gallinago*.
- Minton, C.** 2000. Curlew Sandpipers *Calidris ferruginea* swimming when feeding. *Stilt* 37: 22–23. (165 Dalgetty Road, Beaumaris 3139 Vic., Australia, e-mail: mintons@ozemail.com.au)
- Minton, C. & Whitelaw, J.** 2000. Waders roosting on mangroves. *Stilt* 37: 23–24. *Calidris ruficollis*, *C. ferruginea*, *Tringa nebularia*, *Pluvialis fulva*.
- Pfister, C., Kasprzyk, M.J. & Harrington, B.A.** 1998. Body-fat levels and annual return in migrating Semipalmated Sandpipers. *Auk* 115: 904–915. *Calidris pusilla*.
- Piersma, T., Koolhaas, A., Dekkinga, A. & Gwinner, E.** 2000. Red blood cell and white blood cell counts in sandpipers (*Philomachus pugnax*, *Calidris canutus*): effects of captivity, season, nutritional status, and frequent bleedings. *Can. J. Zool.* 78: 1349–1355. (Neth. Inst. Sea Res. NIOZ, POBox 59, NL-1790 AB Den Burg, Texel, Netherlands, e-mail: theunis@nioz.nl).
- Plissner, J.H., Haig, S.M. & Oring, L.W.** 1999. Within- and between-year dispersal of American Avocets among multiple western Great basin wetlands. *Wilson Bull.* 111(3): 314–320. *Recurvirostra americana*.
- Van Roomen, M.W.J.** 2001. Feeding behaviour of *Tringa species in the Sivash*, August 1998. In: Van der Winden, J., E.A. Diadicheva, W.T. de Nobel & M.W.J. van Roomen (eds.). Counts and ecology of waterbirds in the Sivash, Ukraine, August 1998. – WIWO Report 71. – WIWO: Zeist, 2001, p. 73–78. *Tringa erythropus*, *T. totanus*, *T. stagnatilis*, *T. nebularia*, *T. ochropus*, *T. glareola*.
- Sanders, M.D.** 2000. Enhancing food supplies for waders: inconsistent effects of substratum manipulations on aquatic invertebrate biomass. *J. Appl. Ecol.* 37(1): 66–76.
- Sasvari, L. & Hegyi, Z.** 2000. Avian predators influence the daily time budget of Lapwings *Vanellus vanellus*. *Folia Zool.* 49(3): 211–220.
- Scheiffarth, G.** 2001. The diet of Bar-tailed Godwits *Limosa lapponica* in the Wadden Sea: combining visual observations and faeces analysis. *Ardea* 89(3): 481–494. (Inst. Vogelforschung, Vogelwarte Helgoland, An der Vogelwarte 21, D-26386 Wilhelmshaven, Germany, e-mail: g.scheiffarth@t-online.de).
- Shepherd, P.C.F., Lank, D.B., Smith, B.D., Warnock, N., Kaiser, G.W. & Williams, T.D.** 2001. Sex ratios of Dunlin wintering at two latitudes on the Pacific coast. *Condor* 103(2): 352–360. (Ctr. Wildl. Ecol., Dept. Biol. Sci., Simon Fraser Univ., Burnaby, BC, V5A 1S6 Canada). *Calidris alpina*.
- Shepherd, P.C.F. & Boates, J.S.** 1999. Effects of commercial baitworm harvest on Semipalmated Sandpipers and their prey in the Bay of Fundy Hemispheric Shorebird Reserve. *Conserv. Biol.* 13: 347–356. (Ctr. Wildl. Ecol., Dept. Biol. Sci., Simon Fraser Univ., Burnaby, BC, V5A 1S6 Canada, e-mail: pshepher@sfu.ca). *Calidris pusilla*.
- Sitters, H.P., Gonzalez, P.M., Piersma, T., Baker, A.J. & Price, D.J.** 2001. Day and night feeding habitat of Red Knots in Patagonia: profitability versus safety? *J. Field Ornithol.* 72(1): 86–95. (Limos, Old Ebford lane, Ebford, Exeter EX3 0QR, UK, e-mail: hsitters@aol.com). *Calidris canutus rufa*.
- Smith, A.C. & Nol, E.** 2000. Winter foraging behavior and prey selection of the Semipalmated Plover in coastal Venezuela. *Wilson Bull.* 112: 387–398. *Charadrius semipalmatus*.
- Summers, R.W., Nicoll, M. & Peach, W.** 2001. Numbers, migration phenology and survival of Purple Sandpipers *Calidris maritima* at Gourdon, eastern Scotland. *Bird Study* 48(2): 139–146. (Lismore, Mill Crescent, North Kessock, Ross-shire, IV1 3XY, UK).
- Valle, R. & Scarton, F.** 1999. The presence of conspicuous associates protects nesting Redshank *Tringa totanus* from aerial predators. *Ornis Fenn.* 76: 145–148.
- Wolcott, D.L. & Wolcott, T.G.** 1999. High mortality of Piping Plovers on beaches with abundant ghost crabs: correlation, not causation. *Wilson Bull.* 111: 321–329. *Charadrius melanotos*.

DISTRIBUTION AND FAUNISTICS

- Alcorn, M. & Alcorn, R.** 2000. Seasonal migration of Banded Stilt *Cladorhynchus leucocephalus* to the Natimuk-Douglas salt pans in western Victoria, Australia. *Stilt* 36: 7–10. (17 Lawrence St., Horsham, 3400 Vic., Australia, e-mail: rmalcorn@netconnect.com.au)
- Auchu, C.** 2000. A Common Ringed Plover at Les Escoumins, Quebec. *Birders J.* 9: 192–195. (9B Gagnon, C.P. 853, Les Escoumins, PQ G0T 1K0, Canada). *Charadrius hiaticula*.



- Baker, A.J., Piersma, T. & Greenslade, A.D.** 1999. Molecular vs. phenotypic sexing in Red Knots. *Condor* 101: 887–893. *Calidris canutus*.
- Barter, M.A., Wilson, J.R., Li, Z.W., Li, X.Y., Liu, Y.F. & Tian, H.S.** 2000. Northward migration of shorebirds in the Shuangtaizihekou National Nature Reserve, Liaoning Province, China in 1998 and 1999. *Stilt* 37: 2–9. (21 Chivalry Av., Glen Waverley, Vic. 3150, Australia, e-mail: markbarter@optusnet.com.au)
- Barter, M.A., Wilson, J.R., Li, Z.W., Dong, Z.G., Cao, Y.G. & Jiang, L.S.** 2000. Yalu Jiang National Nature Reserve, north-eastern China – a newly discovered internationally important Yellow Sea site for northward migrating shorebirds. *Stilt* 37: 13–20.
- Barter, M.A., Li, Z.W. & Xu, J.L.** 2001. Shorebird numbers on the Tianjin Municipality coast in May 2000. *Stilt* 39: 2–9.
- Boyd, H. & Piersma, T.** 2001. Why do few Afro-Siberian Knots *Calidris canutus canutus* now visit Britain? *Bird Study* 48(2): 147–158. (Piersma: Neth. Inst. Sea Res. NIOZ, POBox 59, NL-1790 AB Den Burg, Texel, Netherlands, e-mail: theunis@nioz.nl).
- Brewster, C.A.** 2000. Breeding of Collared (Red-winged) Pratincoles *Glareola pratincola* at Shashe dam. *Babbler* 36: 21. (Private Bag 0024, Bobonong, Botswana)
- Chernichko, I.I., Van der Winden, J., Gorlov, P.I., De Nobel, W.T., Van Roomen, M.W.J. & Siokhin, V.D.** 2001. *Numbers of waterbirds in the Sivash, August 1998*. In: Van der Winden, J., E.A. Diadicheva, W.T. de Nobel & M.W.J. van Roomen (eds.). Counts and ecology of waterbirds in the Sivash, Ukraine, August 1998. – WIWO Report 71. – WIWO: Zeist, 2001, p. 17–37. Most numerous waders of the estimated total of 520,000–550,000: *Philomachus pugnax*, *Calidris ferruginea*, *Calidris alpina*, *Recurvirostra avosetta*, *Calidris alba*. With distribution maps.
- Collins, P., Boyle, A., Minton, C. & Jessop, R.** 2001. The importance of inland claypans for waders in Roebuck Bay, Broome, NW Australia. *Stilt* 38: 4–8. RMB 4009, Cowes, Vic. 3922, Australia, e-mail: moonbird@waterfront.net.au)
- Collins, P. & Jessop, R.** 2001. Waterbirds at Taylors Lagoon, near Broome, Western Australia. *Stilt* 39: 27–30. Internationally significant numbers of *Glareola maldivarum*, *Numenius minutus*.
- Craig, M., Darnell, J., Davis, C., Kirkby, T. & Singor, M.** 2001. Waders at Lake McLarty, Western Australia. *Stilt* 38: 18–32. (PO Box 141, Yungaburra, Qld. 4872, Australia)
- Crawford, D.N.** 1999. Migratory waders in south-east Asia, Australia and New Zealand: reference data and the relative abundance of each species. *Stilt* 35: 2–9. (1/6 Marrawah St., Lyons 2606 ACT, Australia). *Calidris ruficollis* and *Limosa lapponica* most abundant species.
- Crossland, A.C.** 2000. Notes on the waders wintering at three sites at the northwestern tip of Sumatra, Indonesia. *Stilt* 36: 4–6. (46 Frensham Crescent, Christchurch, 8006, New Zealand). 12 Wader species were recorded. Most numerous were *Charadrius dubius*, *Calidris ruficollis*, *Pluvialis fulva*.
- Van Dijk, A.J., Van der Weide, M., Zoetebier, D. & Plate, C.** 2001. Kolonievogels en zeldzame broedvogels in Nederland in 1998. (Rare and colonial breeding birds in The Netherlands in 1998). *Limosa* 74(1): 1–16. In Dutch with English summary. (SOVON, Rijksstraatweg 178, NL-6573 DG Beek-Ubbergen, Netherlands). Breeding waders: *Recurvirostra avosetta*, *Charadrius dubius*, *C. hiaticula*, *C. alexandrinus*, *Calidris alpina*, *Philomachus pugnax*, *Actitis hypoleucos*.
- Dinsmore, S.J.** 1999. First record of a Snowy Plover in North Carolina. *Chat* 63: 85. *Charadrius alexandrinus*.
- Dodd, S.L., Spinks, M.D. & Wilkinson, P.M.** 1999. Abundance and distribution of wintering Piping Plovers on the coast of South Carolina: findings from the 1997, 1998, and 1999 mid-winter censuses. *Chat* 63: 155–166. *Charadrius melanotos*.
- Eckert, C.** 1999. A notable movement of godwits in the Yukon, spring 1999. *Birders J.* 8: 149–150. *Limosa lapponica*, *L. haemastica*.
- Eckert, C.D. & Sinclair, P.H.** 2000. The Yukon's first Pacific Golden Plover at Lees Marsh. *Birders J.* 9: 150–152. (1402 Elm St., Whitehorse, YT, Y1A 4B6 Canada). *Pluvialis fulva*.
- Fernandez, G., De la Cueva, H. & Warnock, N.** 2001. Phenology and length of stay of transient and wintering Western Sandpipers at Estero Punta Banda, Mexico. *J. Field Ornithol.* 72(4): 509–520. (Centro de Investigacion Cientifica y de Educacion Superior de Ensenada, Km 17 Carretera Tijuana-Ensenada, Ensenada, Baja California, Mexico). *Calidris mauri*.
- Finn, P.G., Catterall, C.P. & Driscoll, P.V.** 2001. The low tide distribution of Eastern Curlew on feeding grounds in Moreton Bay, Queensland. *Stilt* 38: 9–17. (Fac. Environm. Sci., Griffith Univ., Nathan, Qld. 4111, Australia, e-mail: P.Finn@mailbox.gu.edu.au). *Numenius madagascariensis*.
- Flore, B.-O.** 2000. Mornellregenpfeifer *Charadrius morinellus* im Nationalpark Hohe Tauern/Österreich. [Dotterel *Charadrius morinellus* at the National Park Hohe Tauern/Austria]. *Monticola* 8: 192. In German. (Hafkemeyerweg 20, D-49084 Osnabrück, Germany)
- Fraser, P.A. & Rogers, M.J.** 2001. Report on scarce migrant birds in Britain in 1999. *Brit. Birds* 94(12): 560–589. (c/o 2 Churchtown Cottages, Towendack, St Ives, Cornwall TR26 3AZ, UK). Waders: *Charadrius alexandrinus*, *Calidris temminckii*, *C. melanotos*, *Tryngites subruficollis*, *Phalaropus lobatus*, *P. fulicarius*.
- Geene, R. (ed.).** 2001. *Waterbird count of Zanzibar and Pemba Islands; Tanzania, January 1998*. Zeist: WIWO (Foundation Working Group International Wader and Waterfowl Research). WIWO-report 73, 85 p. Available from: Stichting WIWO, Fetha 23, NL-3633 CT Vreeland, Netherlands. Price Dfl. 30 + 15 admin. costs, postal giro account 2666009). Most numerous waders: *Dromas ardeola*, *Calidris ferruginea*, *Numenius phaeopus*, *Xenus cinereus*.
- Gerasimov, Y., Artukhin, N., Gerasimov, N. & Lobkov, E.** 1999. Status of shorebirds in Kamchatka, Russia. *Stilt* 34: 31–34. (Kamchatka Inst. Ecol., Far-east Branch Russ. Acad. Sci., Rybakov 19A, Petropavlovsk-Kamchatsky, 683024 Russia, e-mail: bird@svyaz.kamchatka.su.).
- Gerasimov, Yu. & Gerasimov, N.N.** 2000. The importance of the Moroshechnaya River estuary as a staging site for shorebirds. *Stilt* 36: 20–25.
- Gerasimov, Yu.N. & Gerasimov, N.N.** 2000. Information on the northward migration of Great Knot *Calidris tenuirostris* in Kamchatka, Russia. *Stilt* 36: 35–38.
- Gerasimov, Yu.N. & Gerasimov, N.N.** 2001. Records of northward migration of Dunlin *Calidris alpina* through Kamchatka, Russia. *Stilt* 39: 37–40.



- Gerasimov, Yu. N.** 2001. Northward migration of shorebirds at Kharchinskoe Lake, Kamchatka, Russia. *Stilt* 39: 41–44.
- Gunnarsson, T.G.** 2001. Phenology of wader migration through Ölfusa estuary, S-Iceland. *Bliki* 22: 1–12. In Icelandic with English summary. (Sch. Biol. Sci., Univ. East Anglia, Norwich NR4 7TJ, UK, e-mail: tgg@hi.is). Gives timing of migration for *Calidris alpina*, *C. alba*, *Limosa limosa*, *Charadrius dubius*, *Tringa totanus*, *Phalaropus lobatus*.
- Gunnarsson, T.G., Potts, P.M., Gill, J. & Croger, R.** 2001. Studies of Icelandic Black-tailed Godwits. *Bliki* 22: 55–61. In Icelandic with English summary. Overview of current knowledge of *Limosa limosa islandica* and expectations for ongoing research.
- Heuer, J.** 2000. Erfolgreiche Brut des Rotschenkels *Tringa totanus* im nördlichen Harzvorland. [Successful breeding of Redshank *Tringa totanus* in the northern Harzvorland]. *Milvus Braunschweig* 19: 51–54. In German. (Am Güdecken 33, D-38667 Bad Harzburg, Germany). First breeding in 32 years.
- Hötker, H. & Frederiksen, M.** 2001. Estimation of total numbers of Pied Avocets *Recurvirostra avosetta* using a moulting site in the Danish Wadden Sea. *Ardea* 89(3): 537–541. (Naturschutzbund Deutschland, Goosstroot 1, D-24861 Bergenhusen, Germany, e-mail: nabu-inst.hoetker@t-online.de)
- Huettmann, F.** 2001. Summary of a trip to the Sea of Okhotsk to study migrating shorebirds: May 2000 on southern Sakhalin Island and August 2000 on western Kamchatka and Magadan region. *Stilt* 39: 65–71. (Dept. Biol. Sci., Simon Fraser Univ., 8888 Univ. Dr., Burnaby B.C., V5A 1S6 Canada, e-mail: huettman@sfsu.ca)
- Hulscher, J.B. & Bunskeoke, E.J.** 2001. Vijfentwintig jaar goudplevier tellingen in de Eempolders. [Twenty-five years of Golden Plover counts in the Eempolder (Netherlands)]. *Vogeljaar* 49(5): 199–203. In Dutch. (Rijksstraatweg 305, NL-9752 CE Haren, Netherlands). *Pluvialis apricaria*.
- Islam, M.S.** 2001. Southward migration of shorebirds in the Ganges delta, Bangladesh. *Stilt* 39: 31–36. (House No. 15/22, Munshi Bari, South Chartha, Comilla-3500 Bangladesh). 31 Wader species were recorded.
- Johnson, O.W., Bennett, A.J., Alsworth III, L., Bennett, L.A., Johnson, P.M., Morgart, J.R. & Kienholz, R.J.** 2001. Radio-tagged Pacific Golden Plovers: the Hawaii–Alaska link, spring destinations, and breeding season survival. *J. Field Ornithol.* 72(4): 537–546. (Dept. Ecol., Montana State Univ., Bozeman, MT 59717, USA). *Pluvialis fulva*.
- Kennerley, P.R. & Bishop, K.D.** 2001. Bristle-thighed Curlew *Numenius tahitiensis* on Manus, Admiralty Islands, Papua New Guinea. *Stilt* 38: 54–55. (16 Coppice Close, Melton Woodbridge, Suffolk IP12 1RX, UK).
- Krott, N.** 2000. Steppenkiebitz (*Chettusia gregaria*) am Heerter See. [Sociable Plover (*Chettusia gregaria*) at Heerter See]. *Milvus Braunschweig* 19: 69. In German. (Wilma-Bayer-Weg 8, D-31141 Hildesheim, Germany)
- Lane, B.A. & Rogers, D.I.** 2000. The Australian Painted Snipe *Rostratula benghalensis australis*: an endangered species? *Stilt* 36: 26–34. (Rogers: 340 Nink's Rd., St Andrews, Vic. 3761, Australia, e-mail: drogers@melbpc.org.au).
- Leader, P.J.** 2001. Kentish Plovers with a complete breast-band. *Brit. Birds* 94(5): 246–247. (60 Cypress Dr., Palm Springs, Hong Kong). *Charadrius alexandrinus dealbatus*, Hong Kong.
- Ma, M., Minton, C., Kraaijeveld, K., Jessop, R. & Gosbell, K.** 2001. Banding of the waders and terns between Australia and China. *Zool. Res.* 22(2): 166–168.
- McCrie, N.** 2000. A sighting of the Green Sandpiper *Tringa ochropus* at Darwin, Northern Territory. *Aust. Bird Watcher* 18: 229–232. (PO Box 41382, Casuarina, NT 0811, Australia). First confirmed record for Australia.
- Meininger, P.L., Arts, F.A. & Van Swelm, N.D.** 2000. *Kustbroedvogels in het Noordelijk Deltagebied: ontwikkelingen, knelpunten en potenties. [Coastal breeding birds in the Northern Delta area: trends, problems and potential].* (Rapport RIKZ/2000.052). Middelburg: Rijksinstituut voor Kust en Zee RIKZ [etc.], 238 p. In Dutch. (RIKZ, PO Box 8039, NL-4330 EA Middelburg, Netherlands). Development of breeding populations of coastal breeding birds in the 20th century in the Delta area, SW Netherlands. Waders: *Himantopus himantopus*, *Recurvirostra avosetta*, *Charadrius hiaticula*, *C. alexandrinus*.
- Meissner, W. & Kozirog, L.** 2001. Biometrics of Turnstone *Arenaria interpres* migrating in autumn through the Gulf of Gdansk region. *Ornis Svecica* 11: 181–188. (Dept. Vertebrat. Ecol. & Zool., Univ. Gdansk, Al. Legionow 9, PL-80-441 Gdansk, Poland, e-mail: biowm@univ.gda.pl)
- Meissner, W.** 2001. Spring migration of Common Snipe *Gallinago gallinago*, in the Gulf of Gdansk area (Poland). *Alauda* 69(3): 429–434.
- Minton, C. & Jessop, R.** 2000. Corrections to sightings of waders leg-flagged in North-Western Australia report number 6. *Stilt* 36: 39–41.
- Minton, C.** 2001. Summary of overseas sightings of waders leg flagged in Victoria and south eastern South Australia. *Stilt* 39: 45–47. (165 Dalgety Rd., Beaumaris 3139 Vic., Australia, e-mail: mintons@ozemail.com.au)
- Minton, C., Jessop, R., Collins, P. & Graham, D.** 2001. Sightings of waders leg-flagged in Victoria: report number 8. *Stilt* 39: 48–60.
- Moores, N.** 1999. A survey of the distribution and abundance of shorebirds in South Korea during 1998–1999: interim summary. *Stilt* 34: 18–29. (251 Nooha-Dong, Chongro-Gu, Seoul 110-042, Republic of Korea).
- Morrison, R.I.G., Gill, R.E. Jr., Harrington, B.A., Skagen, S., Page, G.W., Gratto-Trevor, C.L. & Haig, S.M.** 2001. *Estimates of shorebird populations in North America.* Canadian Wildlife Service Occasional Paper No. 104. Ontario: Environment Canada, 64 p. Population size of 57 shorebird species is described.
- Ogilvie, M. and the Rare Breeding Birds Panel** 2001. Rare breeding birds in the United Kingdom in 1999. *Brit. Birds* 94(8): 344–381. (Glencairn, Bruichladdich, Isle of Islay PA49 7UN, UK). Breeding waders: *Recurvirostra avosetta*, *Burhinus oedicnemus*, *Charadrius dubius*, *Calidris morinellus*, *Calidris temminckii*, *Limosa limosa*, *Numenius phaeopus*, *Tringa nebularia*, *Tringa ochropus*, *Tringa glareola*, *Phalaropus lobatus*.
- Oldekop, W., Melchert, F. & Hermenau, B.** 2000. 50 Jahre Limikolenbeobachtungen in der Umgebung Braunschweigs. [50 Years of wader observations in de Braunschweig area]. *Milvus Braunschweig* 19: 1–35. In German. (Bergiusstr. 2, D-38116 Braunschweig, Germany).
- Paton, D.C., Ziembicki, M., Owen, P., Hill, B. & Bailey, C.** 2000. Distribution and abundance of migratory waders and their food in the estuarine areas of the Murray



- Mouth and patterns in the composition of sediments. *Stilt* 37: 47. (Dept. Environm. Biol., Univ. Adelaide, SA 5005, Australia).
- Pearce-Higgins, J.W.** 2001. A model describing the exchange of individuals between Turnstone *Arenaria interpres* roosts on the North Wales coast. *Ring. & Migr.* 20(3): 209–212. (Sch. Biol. Sci., 3.239 Stopford Bldg., Univ. Manchester, Oxford Rd., Manchester M13 9PT, UK)
- Riegen, A.C.** 1999. Movements of banded Arctic waders to and from New Zealand. *Notornis* 46: 123–142. (231 Forest Hill Rd., Waiatarua, Auckland 8, New Zealand, e-mail: riegen@xtra.co.nz)
- Riegen, A.C.** 2000. Sightings of leg-flagged shorebirds from New Zealand report number 1: July 2000. *Stilt* 37: 28–31. *Calidris canutus*, *Limosa lapponica*, *Arenaria interpres*.
- Rogers, D., Battley, P., Russell, M. & Boyle, A.** 2000. A high count of Asian Dowitchers in Roebuck Bay, north-western Australia. *Stilt* 37: 10–12. (340 Nink's Rd., St Andrews, Vic. 3761, Australia, e-mail: drogers@melbpc.org.au). *Limnodromus semipalmatus*.
- Rogers, D.I., Boyle, A. & Hassell, C.** 2001. Occasional counts No. 5. Wader counts on Kidneybean Claypan and adjacent Roebuck Plains, northwestern Australia. *Stilt* 38: 57–63.
- Sangha, H.S.** 2000. Recent sightings of *Vanellus gregarius* (Pallas) at Tai Chhapar and Revasa, Rajasthan. *J. Bombay Nat. Hist. Soc.* 97(2): 278.
- Sanzenbacher, P.M. & Haig, S.M.** 2001. Killdeer population trends in North America. *J. Field Ornithol.* 72(1): 160–169. (USGS Forest & Rangeland Ecosyst. Sci. Ctr., 3200 SW Jefferson Way, Corvallis, OR 97331, USA). *Charadrius vociferus*.
- Savard, G.** 2000. A partial albino Semipalmented Plover at Saint-Fulgence, Quebec. *Birders J.* 9: 149. *Charadrius semipalmatus*.
- Scheiffarth, G.** 2001. Bar-tailed Godwits (*Limosa lapponica*) in the Sylt-Romo Wadden Sea: which birds, when, from where, and where to? *Vogelwarte* 41(1): 53–69. (Inst. Vogelforschung, Vogelwarte Helgoland, An der Vogelwarte 21, D-26386 Wilhelmshaven, Germany, e-mail: g.scheiffarth@t-online.de).
- Schrücke, V., Triplet; P. & Yesou, P.** 2001. (French contribution to the knowledge of western Palaearctic water birds overwintering in the Senegal River delta). *Alauda* 69: 135–148. In French with English summary. (Ministère de l'Amenagement du Territoire et de l'Environnement, 53 rue Russeil, F-44000 Nantes, France, e-mail: v.schrücke@onc.gouv.fr).
- Segelbacher, G.M.** 2000. Zum Durchzug von Limikolen am Rohrsee RV/Baden-Württemberg – eine Zusammenfassung der Jahre 1994–1998. [On the migration of waders at the Rohrsee RV/Baden-Württemberg – a sumary for the years 1994–1998]. *Orn. Jh. Bad.-Württ.* 16(1): 33–63. In German. (Mörikestr. 35, D-88285 Bodnegg, Germany).
- Singor, M.** 1999. Hooded Plover report No. 2: 1996–1999. The search continues. *Stilt* 35: 42. (149a Bishopsgate St., Carlisle 6101 WA, Australia). *Thinornis rubricollis*.
- Sundar, K.S.G.** 2000. The greyheaded lapwing, *Vanellus cinereus* (Blyth) in Kaliveli Tank, Tamil Nadu. *J. Bombay Nat. Hist. Soc.* 97(2): 277.
- Tomkovich, P.S. & Riegen, A.C.** 2000. Mixing of Red Knot populations in Australasia: some thoughts. *Stilt* 37: 25–26. (Zool. Mus., Moscow Lomonosov State Univ., Bol. Nikitskaya St. 6, Moscow 103009, Russia, e-mail: tomkovic@l.zoomus.bio.msu.ru). *Calidris c. canutus*, *C.c. rogersi*.
- Tree, A.J.** 1998. Movements of the Blacksmith Plover in south-central Africa. *Honeyguide* 44(4): 199–203. (PO Box 211, Bathurst 6166, South Africa, e-mail: tony.tree@xsinet.co.za). *Vanellus armatus*, three populations suggested. Local breeders which are nomadic; population shifts from arid areas during droughts; and more northerly migrants visiting and passing through during the summer rainfall season.
- Tree, A.J.** 1999. An analysis of the occurrence of some rare or scarce Palaearctic waders in Zimbabwe. *Honeyguide* 45(2): 120–126. *Charadrius mongolus*, *C. leschenaultii*, *Pluvialis squatarola*, *Arenaria interpres*, *Xenus cinereus*, *Tringa totanus*, *Calidris melanotos*, *C. alba*, *Limicola falcinellus*, *Gallinago media*, *Limosa limosa*, *L. lapponica*, *Numenius arquata*, *N. phaeopus*, *Phalaropus fulicaria*.
- Tree, A.J.** 2000. The changing status of the Curlew Sandpiper in Zimbabwe. *Honeyguide* 46(1): 19–22. *Calidris ferruginea*. With the advent of a mega-dam building the Curlew Sandpiper has increased from a rare visitor to a common southbound migrant with increasing over-summering and over-wintering. Initially mainly young birds were observed, but now an established adult population visits the area.
- Tree, A.J.** 2001. Kittlitz's Plover as an intra-African migrant. *Honeyguide* 47(1): 10–15. *Charadrius pecuarius* has adapted successfully to man-made aquatic habitats. Strong movements into southern Africa during summer rains with all populations from higher rainfall areas at least as far north as the Congo and Tanzania migrating. Extent of southward penetration dependent on availability of suitable water bodies. Migrating birds accumulate lipid deposits sufficient for non-stop flights of up to 2000 km. Coastal populations relatively sedentary, inland southern birds nomadic, northern birds migratory/nomadic.
- Tree, A.J.** 2001. The occurrence of Painted Snipe in Zimbabwe and neighbouring countries. *Honeyguide* 47(2): 165–169. *Rostratula benghalensis* has a distinct degree of seasonality in occurrence in lower rainfall areas and greater stability in Zambia. General pattern of movement probably north–south but south of Zambezi tendency for overlay of east–west movement. Possibility of more southerly birds leap-frogging more sedentary populations and originating further north in Africa. No proved examples of site-fidelity during ringing studies.
- Van der Have, T.M., Keijl, G.O., Mansoori, J. & Morozov, V.V.** 2001. *Searching for Slender-billed Curlews in Iran, January–February 2000*. Zeist: WIWO (Foundation Working Group International Wader and Waterfowl Research). WIWO-report 72, 69 p. Available from: Stichting WIWO, Fetha 23, NL-3633 CT Vreeland, Netherlands. Price Dfl. 20 + 15 admin. costs, postal giro account 2666009.
- Wierda, J.** 2001. Steltstrandloper in Noord-Holland in juli 2000. (Stilt Sandpiper in Noord-Holland in July 2000). *Dutch Birding* 23(1): 5–7. In Dutch with English summary. *Micropalama himantopus*, Netherlands.
- Wilson, J.** 2000. The northward movement of immature Eastern Curlews in the Austral winter as demonstrated by the Population Monitoring Project. *Stilt* 36: 16–19. (13/27 Giles St., Kingston, 2604 ACT, Australia). *Numenius madagascariensis*.



- Wilson, J.R.** 2000. Occasional counts No. 3. Wader counts on Eyre Island, South Australia. *Stilt* 36: 42–44.
- Wilson, J.R.** 2000. Occasional counts No. 4. Wader counts at Yantabulla Swamp (Cuttburra Basin), New South Wales, Australia. *Stilt* 37: 32–33.
- Wilson, J.R.** 2000. A survey of South Australian waders in early 2000. *Stilt* 37: 34–45.
- Wilson, J.R.** 2001. Population monitoring counts for winter 1999, summer 2000 and winter 2000. *Stilt* 38: 64–69.
- Van der Winden, J., Diadicheva, E.A., De Nobel, W.T. & Van Roomen, W.J.** (eds.) 2001. *Counts and ecology of waterbirds in the Sivash, Ukraine, August 1998*. Zeist: WIWO (Foundation Working Group International Wader and Waterfowl Research). WIWO-report 71, 118 p. Available from: Stichting WIWO, Fetha 23, NL-3633 CT Vreeland, Netherlands. Price Dfl. 30 + 15 admin. costs, postal giro account 2666009).
- Zhu, S.Y., Li, Z.W., Lu, J.Z., Shan, K. & Barter, M.A.** 2001. Northward migration of shorebirds through the Huang He delta, Shandong Province, in the 1997–1999 period. *Stilt* 38: 33–38. (Huang He National Nat. Res., Cao Zhou Rd., Dongying City, Shandong, Peoples Republic of China)

MISCELLANEOUS

- Augustin, J.C., Grabough, J.W. & Marshall, M.R.** 1999. Validating macroinvertebrate assumptions of the shorebirds management model for the Lower Mississippi Valley. *Wildl. Soc. Bull.* 27: 552–558.
- Bart, J., Battaglia, D. & Senner, N.** 2001. Effects of color bands on Semipalmated Sandpipers banded at hatch. *J. Field Ornithol.* 72(4): 521–526. (Snake River Field stn., USGS, 970 Luke St., Boise, Idaho 83706, USA). *Calidris pusilla*.
- Burton, N.H.K.** 2001. Reaction of Redshank *Tringa totanus* to colour-rings. *Ring. & Migr.* 20(3): 213–215. (BTO, The Nunnery, Thetford, Norfolk IP24 2PU, UK).
- Dekker, M.H.A., Piersma, T. & Sinninghe Damsté, J.S.** 2000. Molecular analysis of intact preen waxes of *Calidris canutus* (Aves: Scolopacidae) by gas chromatography/mass spectrometry. *Lipids* 35: 533–541. (Piersma: Neth. Inst. Sea Res. NIOZ, POBox 59, NL-1790 AB Den Burg, Texel, Netherlands, e-mail: theunis@nioz.nl).
- Gunn, T. & Pert, P.** 2001. Powdercoating of ABBS bands – an example using Crested Tern bands. *Stilt* 38: 56. (PO Box 228, Boronia, Vic. 3155, Australia, e-mail: petina.pert@nre.vic.gov.au).
- Jessop, R. & Collins, P.** 2000. The effects of cyclonic weather conditions on the bird life around Broome, Western Australia. *Stilt* 36: 11–15. (Philip Island Nature Park, PO Box 97, Cowes, Vic. 3922, Australia, e-mail: rjessop@penguins.org.au).
- Kooiker, G.** 2000. Empfehlungen zur Methodik von Brutbestandsaufnahmen beim Kiebitz *Vanellus vanellus*: Was zählen, wann und wie oft? (Recommendations on methods of performing a population census of Lapwings *Vanellus vanellus* during the breeding season). *Orn. Jh. Bad.-Württ.* 16(2): 203–207. In German with English summary. (Katharinenstr. 107, D-49078 Osnabrück, Germany).
- Kosinski, Z.** 1999. Effects of lake morphometry, emergent vegetation and shore habitat on breeding bird communities. *Acta Ornithol.* 34: 27–35. (Dept. Avian Biol. & Ecol., A. Mickiewicz Univ., Fredy 10, PL-61 701 Poznan, Poland)
- Lee, N.M. & Hockey, P.A.R.** 2001. Biases in the field estimation of shorebird prey sizes. *J. Field Ornithol.* 72(1): 49–61. (Percy FitzPatrick Inst. Afr. Ornithol., Univ. Cape Town, Rondebosch, South Africa 7701)
- Loyn, R.H., Dann, P. & McCulloch, E.** 2001. Important wader sites in the East Asian-Australasian Flyway: 1. Western Port, Victoria, Australia. *Stilt* 38: 39–53. (Arthur Rylah Inst., Dept. nat. Resourc. & Environm., PO Box 137, Heidelberg, Vic. 3084, Australia). Most numerous: *Calidris ruficollis*, *C. ferruginea*, *Numenius madagascariensis*.
- Paton, D.C., Ziembicki, M., Owen, P. & Heddle, C.** 2000. Disturbance distances for water birds and the management of human recreation with special reference to the Coorong region of South Australia. *Stilt* 37: 46. (Dept. Environm. Biol., Univ. Adelaide, SA 5005, Australia).
- Püsichel, H.** 1998. Schutzaun während der Brut beim Flussregenpfeifer (*Charadrius dubius*). (Fenced-in nesting site of the Little Ringed Plover (*Charadrius dubius*)). *Naturschutz südl. Oberrhein* 2: 205–206. In German with English summary. (Augustastr. 4, D-77654 Offenburg, Germany).
- Riegen, A.** 2000. Bivalve comes to grips with a Terek Sandpiper in China. *Stilt* 37: 22. (231 Forest Hill Rd., Waiatarua, Auckland 8, New Zealand, e-mail: riegen@xtra.co.nz) *Xenus cinereus*.
- Sinninghe Damsté, J.S., Dekker, M.H.A., Van Dongen, B.E., Schouten, S. & Piersma, T.** 2000. Structural identification of the diester preen-gland waxes of the Red Knot (*Calidris canutus*). *J. Nat. Prod.* 63: 381–384. (Piersma: Neth. Inst. Sea Res. NIOZ, PO Box 59, NL-1790 AB Den Burg, Texel, Netherlands, e-mail: theunis@nioz.nl).
- Thompson, D.R. & Dowding, J.E.** 1999. Site-specific heavy metal concentrations in blood of South Island Pied Oystercatchers *Haematopus ostralegus finschi* from the Auckland region, New Zealand. *Mar. Poll. Bull.* 38: 202–206.
- Wilson, J.R.** 2001. The Australasian Waders Studies Group Population Monitoring Project: where to now? perspectives from the chair. *Stilt* 39: 13–26. (13/27 Giles St., Kingston, 2604 ACT, Australia).

