

Recent publications on waders 80

compiled by HENK KOFFIJBERG

In trying to make these lists as complete as possible, and the next one even more complete and up to date, your help is necessary! Please send new titles, abstracts, reprints, copies, reports, and (important too!) omissions or corrections to: **Henk Koffijberg, Glenn Millerweg 102, NL-1311 RW Almere, The Netherlands.** koffijberg@wanadoo.nl

Remember to include, if possible, the address of the first author, and translations of titles if they are in other languages than English.

For help with this issue I am very grateful to S. Nebel, R.M. Ward, P. Triplet, B. Trolliet, K.-M. Exo and J. Elts. Translated titles from languages other than English are bracketed; with round brackets if the translation is by the author(s), square brackets if not.

N.B. A copy of your paper or abstract would be very helpful to enable me to add some keywords or give added annotation to the titles.

BREEDING

Baird, B. & Dann, P. 2003. The breeding biology of Hooded Plovers, *Thinornis rubricollis*, on Philip Island, Victoria. *Emu* 103(4): 323–328. (Philip Island Nature Pk., PO Box 97, Cowes, Philip Island, Vic 3922, Australia.)

Chandler, R. 2003. Breeding displays of Little Ringed Plover. *Brit. Birds* 96(7): 342–343. (4 Kings Rd., Oundle, Peterborough PE8 4AX, UK). *Charadrius dubius*.

Crossland, A. 2003. Observations at a breeding colony of Oriental Pratincole (*Glareola maldivarum*) near Songkhla Lake, southern Thailand. *Stilt* 43: 28–29. (Parks & Waterways Unit, Christchurch City Council, PO Box 267, Christchurch 8006, New Zealand.)
(andrew.crossland@ccc.govt.nz)

Cuervo, J.J. 2004. Nest-site selection and characteristics in a mixed-species colony of Avocets *Recurvirostra avosetta* and Black-winged Stilts *Himantopus himantopus*. *Bird Study* 51(1): 20–24. (Estacion Experimental de Zonas Aridas, Consejo Superior de Investigaciones Cientificas, Calle General Segura 1, E-04001 Almeria, Soain.)
(jcuerdo@eeza.csic.es)

Holt, S. & Whitfield, D.P. 2003. Timing of nest departure by broods of Dotterel *Charadrius morinellus*. *Bird Study* 50(1): 88–90. (34 Acre End St., Eynsham, Oxfordshire OX29 4PA, UK. sue@holtcress.freeserve.co.uk)

Hötker, H. 2002. Arrival of Pied Avocets *Recurvirostra avosetta* at the breeding site: effects of winter quarters and consequences for reproductive success. *Ardea* 90(3): 379–387. (Forschungs- und Technologiezentrum Westküste, Hafentörn, D-25761 Büsum, Germany.)
(NABU-Inst.hoetker@t-online.de)

Jaensch, R. 2003. Breeding by Australian Painted Snipe in the Diamantina Channel Country, south-western Queensland. *Stilt* 43: 20–22. (Wetlands International, c/o Queensland Herbarium, Mt Coot-tha Rd., Toowong, Qld. 4066, Australia. roger.jaensch@epa.qld.gov.au). *Rostratula benghalensis*.

Middleton, P. 2003. Population ecology, conservation and management of Little Ringed Plovers in South Yorkshire. *Brit. Birds* 96(7): 344–346. (7 Sandhill Grove, Grime-

thorpe, Barnsley, South Yorkshire S72 7AT, UK). *Charadrius dubius*.

Minton, C., Jessop, R., Collins, P. & Hassell, C. 2002. Year 2001 arctic breeding success, as measured by the percentage of first year birds in wader populations in Australia in the 2001/02 austral summer. *Stilt* 42: 32–36. (165 Dalgetty Rd., Beaumaris, Vic. 3193, Australia.)
(mintons@ozemail.com.au)

Minton, C., Jessop, R. & Collins, P. 2003. Variations in apparent annual breeding success of Red-necked Stints and Curlew Sandpipers between 1991 and 2001. *Stilt* 43: 30–33. *Calidris ruficollis*, *C. ferruginea*.

Nebel, S. & McCaffery, B.J. 2003. Vocalization activity of breeding shorebirds: documentation of its seasonal decline and applications for breeding bird surveys. *Can. J. Zool.* 81: 1702–1708. (Dept Biol. Sci., Simon Fraser Univ., 8888 Univ. Dr., Burnaby, BC V5A 1S6, Canada. snebela@sfu.ca). Data recorded for 15 species.

Pearce-Higgins, J.W. & Yalden, D.W. 2003. Golden Plover *Pluvialis apricaria* breeding success on a moor managed for shooting Red Grouse *Lagopus lagopus*. *Bird Study* 50(2): 170–177. (RSPB, Dunedin House, 25 Ravelston Terrace, Edinburgh EH4 3TP, UK.)
(james.pearce-higgins@rspb.org.uk)

Seymour, A.S., Harris, S., Ralston, C. & White, P.C.L. 2003. Factors influencing the nesting success of Lapwings *Vanellus vanellus* and behaviour of Red Fox *Vulpes vulpes* in Lapwing nesting sites. *Bird Study* 50(1): 39–46. (10 Augustine Rd., London W14 0HZ, UK.)
(aseymour uk@yahoo.co.uk)

Wallander, J. & Andersson, M. 2003. Reproductive tactics of the Ringed Plover *Charadrius hiaticula*. *J. Avian Biol.* 34(3): 259–266. (Dept Zool., Anim. Ecol., Göteborg Univ., Box 463, SE-405 30 Göteborg, Sweden.)
(johan.wallander@zool.gu.se)

Zöllner, T. 2003. Paarsysteme beim Kiebitz *Vanellus vanellus* und deren Auswirkungen auf die Brutpartner und ihre Gelege. (Mating system in the Lapwing *Vanellus vanellus* and its influence on breeding success). *Vogelwelt* 124(1): 35–44. In German with English summary. (Jüttenstr. 9, D-47445 Moers, Germany. zoellner.thomas@web.de)



BEHAVIOUR, ECOLOGY & EVOLUTION

- Aspinal, C.** 2003. Collared Pratincoles feeding at night. *Brit. Birds* 96(6): 302–303. (PO Box 45553, Abu Dhabi, United Arab Emirates). *Glareola pratincola*.
- Atkinson, P.W., Clark, N.A., Bell, M.C., Dare, P.J., Clark, J.A. & Ireland, P.L.** 2003. Changes in commercially fished shellfish stocks and shorebird populations in the Wash, England. *Biol. Conserv.* 114: 127–141.
- Ausden, M., Rowlands, A., Sutherland, W.J. & James, R.** 2003. Diet of breeding Lapwing *Vanellus vanellus* and Redshank *Tringa totanus* on coastal grazing marsh and implications for habitat management. *Bird Study* 50(3): 285–293. (RSPB, The Lodge, Sandy, Bedfordshire SG19 2DL, UK. malcolm.ausden@rspb.org.uk)
- Battley, P.F., Rogers, D.I., Piersma, T. & Koolhaas, A.** 2003. Behavioural evidence for heat-load problems in Great Knots in tropical Australia fuelling for long-distance flight. *Emu* 103(2): 97–103. (2/30 Bucklands Beach Rd., Auckland, New Zealand. philbattley@quicksilver.net.nz). *Calidris tenuirostris*.
- Corso, A.** 2003. Stone-Curlew taking Little Tern chicks. *Brit. Birds* 96(12): 655–656. (Via Camastra, I-10-96100 Siracusa, Italy). *Burhinus oedicnemus*.
- Durell, S.E.A. le V. dit, Goss-Custard, J.D. & Clarke, R.T.** 2001. Modelling the population consequences of age- and sex-related differences in winter mortality in the Oystercatcher, *Haematopus ostralegus*. *Oikos* 95: 69–77. (Ctr. Ecol. Hydrol. (CEH) Dorset, Winfrith Technol. Ctr., Dorchester, Dorset DT2 8ZD, UK. sld@ceh.ac.uk)
- Gillings, S.** 2003. Night feeding in Golden Plovers and Lapwings. *BTO News* 248: 14–15. *Vanellus vanellus*, *Pluvialis apricaria*.
- Gosbell, K. & Minton, C.** 2001. The biometrics and moult of Sanderling *Calidris alba* in Australia. *Stilt* 40: 7–22. (17 Banksia Court, Heathmont, 3135 Vic., Australia. kenbg@ozemail.com.au)
- Insley, H., Swann, B. & Rehfish, M.** 2004. Wader roosts and habitat loss. *BTO News* 251: 16–17. (BTO, The Nunnery, Thetford, Norfolk IP24 2PU, UK). Vulnerability to winter roost loss. Waders discussed: *Haematopus ostralegus*, *Calidris canutus*, *C. alpina*, *Tringa totanus*, *Limosa lapponica*.
- Jackson, D. & Plummer, T.** 2002. Shark Bay waders and seagrass: probings by visitors from afar. *Stilt* 41: 38–39. (RSPB, Scott. HQ., 25 Ravelston Trr., Edinburgh, EH3 4TP, UK. digger.jackson@rspb.org.uk). Use of seagrass wrack by feeding waders.
- Jessop, A.** 2003. Gut analysis of five wader species collected from the NW of Western Australia. *Stilt* 43: 3–13. *Calidris ruficollis*, *Charadrius leschenaultii*, *Calidris canutus rogersi*, *Numenius minutus*, *Calidris ferruginea*.
- Joest, R.** 2003. Junge Sabelschnäbler (*Recurvirostra avosetta L.*) in unterschiedlichen Klimazonen: physiologische und ethologische Anpassungen an ökologische Bedingungen in Norddeutschland und Südspanien. [Young Avocets in different climate zones: physiological and ethological adaptations in northern Germany and southern Spain]. Dissertation Christian-Albrecht University, Kiel. In German. (Freidrichstr. 12, D-33615 Bielefeld, Germany. ralfjoest@yahoo.de). Available online at: <http://e-diss.uni-kiel.de/math-nat.html>
- Johansson, T.** 2001. *Habitat selection, nest predation and conservation biology in a Black-tailed Godwit (Limosa limosa) population*. Ph.D. thesis Uppsala Univ. ISBN 91-554-4958-1.
- Kraaijeveld, K., Minton, C., Collins, P. & Jessop, R.** 2003. The first catch of Banded Stilts in Victoria: biometrics, moult and age/sex structure. *Stilt* 43: 14–19. (Dept Biol., Darwin Bldg., Univ. Coll. London, Gower St., London WC1E 6BT, UK). *Cladorhynchus leucocephalus*.
- Kraaijeveld-Smit, F., Minton, C., Jessop, R. & Collins, P.** 2001. Sexing criteria, age structure, biometrics, and moult of the Pied Oystercatcher, *Haematopus longirostris* in Victoria. *Stilt* 40: 29–37. (Dept Zool., Univ. Melbourne, Parkville 3010 Vic., Australia.) (f.kraaijeveld-smit@pgrad.unimelb.edu.au)
- Le Dréan-Quénéc'hdu, S. & Triplet, P.** 2004. (Influence of sedimentary characteristics of intertidal mud on the feeding habits of waders). *Alauda* 72(1): 11–21. In French with English summary. (Eco-ouest, 3 rue de la Janaie, F-35520 Melesse, France, sldq@club-internet.fr). Literature review of the influence of different types of sediments on the feeding behaviour of waders.
- Melville, D.S., Minton, C. & Collins, P.** 2003. Red-necked Phalaropes at Port Hedland saltworks. *Stilt* 43: 23–27. (Dovedal, R.D. 2 Wakefield, Nelson, New Zealand. david.melville@xtra.co.nz). Biometrics and moult of *Phalaropus lobatus*.
- Minton, C. & Collins, P.** 2003. Marsh Sandpipers can also swim when feeding. *Stilt* 43: 28. (165 Dalgetty Rd., Beaumaris, Vic. 3193, Australia.) (mintons@ozemail.com.au). *Tringa stagnatilis*.
- Nebel, S., Cloutier, A. & Thompson, G.J.** 2004. Molecular sexing of prey remains permits a test of sex-biased predation in a wintering population of Western Sandpipers. *Proc. R. Soc. Lond. B (suppl.)* 271: S321–S323. (Dept Biol. Sci., Simon Fraser Univ., 8888 Univ. Dr., Burnaby, BC V5A 1S6, Canada. snebela@sfu.ca). *Calidris mauri*.
- Owner, D. & Rohweder, D.A.** 2003. Distribution and habitat of Pied Oystercatchers (*Haematopus longirostris*) inhabiting ocean beaches in northern New South Wales. *Emu* 103(2): 163–169. (ERM Mitchell McCotter, PO Box 5279, Port Macquarie, NSW 2444, Australia).
- Piersma, T., Gessaman, J.A., Dekkinga, A. & Visser, D.** 2004. Gizzard and other lean mass components increase, yet Basal Metabolic Rates decrease, when Red Knots *Calidris canutus* are shifted from soft to hard-shelled food. *J. Avian Biol.* 35(2): 99–104. (Dept Mar. Ecol. & Evol., Royal Netherlands Inst. Sea Res. (NIOZ), PO Box 59, NL-1790 AB Den Burg, Texel, The Netherlands. theunis@nioz.nl)
- Rehfish, M.M., Insley, H. & Swann, B.** 2003. Fidelity of overwintering shorebirds to roost on the Moray Basin, Scotland: implications for predicting impacts of habitat loss. *Ardea* 91(1): 53–70. (Insley; 1 Drummond Pl., Inverness IV2 4JT, UK. hugh.insley@freeuk.com). *Haematopus ostralegus*, *Charadrius hiaticula*, *Calidris maritima*, *C. alpina*, *Numenius arquata*, *Tringa totanus*, *Arenaria interpres*, *C. canutus*, *Limosa lapponica*.
- Rohweder, D.A.** 2001. Nocturnal roost use by migratory waders in the Richmond River estuary, northern New South Wales, Australia. *Stilt* 40: 23–28. (Sch. Resourc. Sci. & Manage., Southern Cross Univ., PO Box 157 Lismore, NSW 2480 Australia). *Limosa lapponica*, *Numenius phaeopus*, *N. madagascariensis*, *Tringa nebularia*, *Heteroscelus brevipes*, *Calidris ferruginea*, *Himantopus himantopus*, *Vanellus miles*, *Pluvialis fulva*.



Rubega, M.A. & Oring, L.W. 2004. Excretory organ growth and implications for salt tolerance in hatchling American avocets *Recurvirostra americana*. *J. Avian Biol.* 35(1): 13–15. (Dept Ecol. & Evol. Biol., Univ. Connecticut, 75 North Eagleville Rd., U-3043, Storrs, CT 06269. rubega@uconnvm.uconn.edu).

Summers, R.W., Underhill, L.G., Nicoll, M., Strann, K.-B. & Nilsen, O. 2004. Timing and duration of moult in three populations of Purple Sandpipers *Calidris maritima* with different moult/migration patterns. *Ibis* 146(3): 394–403. (Lismore, Mill Crescent, North Kessock, Inverness IV1 3XY, UK. ron.summers@rspb.org.uk)

Szentirmai, I. & Székely, T. 2004. Diurnal variation in nest material use by the Kentish Plover *Charadrius alexandrinus*. *Ibis* 146(3): 535–537. (Dept Ethol., Eötvös Univ., H-1117 Budapest, Pazmany P. sétány 1/C, Hungary. szentirmai@ludens.elte.hu)

Triplet, P., Méquin, N., Prévost, A., Erlinger, D. & Sueur, F. 2003. (Daytime rhythm of activity of Eurasian Oystercatcher *Haematopus ostralegus*, Eurasian Curlew *Numenius arquata* and Dunlin *Calidris alpina* in winter in the Somme Bay (North France). *Alauda* 71(4): 459–468. In French with English summary. (SMACOPI RN Baie de Somme, Maison Ramsar, 1, Place de l'Amiral Courbet, F-80100 Abbeville, France.) (patricktriplet@baiedesomme.org)

Van Tuinen, M., Waterhouse, D. & Dyke, G.J. 2004. Avian molecular systematics on the rebound: a fresh look at modern shorebird phylogenetic relationships. *J. Avian Biol.* 35(3): 191–194. (Dept Biol. Sci., Stanford Univ., Stanford, CA 94305, USA. mvtuinen@stanford.edu)

Verkuil, Y. & Goeij, P. de 2003. (Do Reeves make different choices? Meadow selection by spring staging Ruffs *Philomachus pugnax* in Southwest Friesland). *Limosa* 76(4): 157–168. In Dutch with English summary. (Kamperfoeliestr. 63, NL-9713 RV Groningen, The Netherlands. y.verkuil@biol.rug.nl)

Verkuil, Y., Have, T.M. van der, Winden, J. van der & Chernichko, I.I. 2003. Habitat use and diet selection of northward migrating waders in the Sivash (Ukraine): the use of Brine Shrimp *Artemia salina* in a variably saline lagoon complex. *Ardea* 91(1): 71–83. (Royal Nethl. Inst. Sea Res. (NIOZ), PO Box 59, NL-1790 AB Den Burg, Texel, The Netherlands. yvonne_verkuil@hotmail.com). *Calidris alpina*, *C. ferruginea*, *Limocula falcinellus*.

Vernon, R. 2003. Feeding behaviour of migrant Ruff. *Brit. Birds* 96(9): 457–458. (16 Orchid Meadow, Pwllmeyric, Chepstow, Gwent NP16 6HP, UK). *Philomachus pugnax* feeding on caterpillars of *Hymonia recurviala*, Fuente-ventura, Canary Islands.

Watson, A. 2004. Woodcock feeding young. *Brit. Birds* 97(2): 98. (Ctr. Ecol. & Hydrol., Banchory, Kincardineshire AB31 4BW, UK). *Scolopax rusticola*.

Whitfield, D.P. 2003. Redshank *Tringa totanus* flocking behaviour, distance from cover and vulnerability to sparrowhawk *Accipiter nisus* predation. *J. Avian Biol.* 34(2): 163–169. (Scott. Nat. Heritage, 2 Anderson Pl., Edinburgh EH6 5NP, UK. phil.whitfield@snh.gov.uk)

Zharikov, Y. & Skilleter, G.A. 2004. A relationship between prey density and territory size in non-breeding Eastern Curlews *Numenius madagascariensis*. *Ibis* 146(3): 518–521. (Mar. Estuar. Ecol. Unit., Dept Zool. Entomol., Univ. Queensland, Brisbane, QLD 4072, Australia.) (yzharikov@zen.uq.edu.au)

DISTRIBUTION & FAUNISTICS

Antonov, A. 2003. A shorebird census of Schastya Bay and the Amur estuary, Sea of Okhotsk, Russia from 6 August–21 September 2002. *Stilt* 44: 52–55. (Khingansky Nat. Reserv., Amur Prov., Russia. hingan@amur.ru). 27 Wader species recorded, most numerous: *Calidris alpina*, *C. ruficollis*, *Numenius phaeopus*, *C. tenuirostris*, *Limosa lapponica*, *Charadrius mongolus*.

Barter, M.A., Du, J.J., Wang, H., Chen, Y.Q., Gao, Z.D., Cheng, H. & Li, C.R. 2002. Shorebird numbers in the Yancheng National Nature Reserve during the 2001 northward migration. *Stilt* 41: 27–34. (21 Chivalry Ave., Glen Waverley, Vic. 3150, Australia.) (markbarter@optusnet.com.au)

Barter, M.A. & Harris, K. 2002. Occasional count No. 6: shorebird counts in the South Australia-SW Queensland region in September–October 2000. *Stilt* 41: 44–47.

Barter, M.A., Riegen, A. & Xu, Q. 2003. Shorebird numbers in Bohai Wan during northward migration. *Stilt* 44: 3–8.

Clark, J.A. 2004. Ringing recoveries confirm higher wader mortality in severe winters. *Ring. & Migr.* 22(1): 43–50. (BTO, The Nunnery, Thetford, Norfolk IP24 2PU, UK)

Crossland, A.C. 2001. Long-term changes in numbers of Variable Oystercatcher (*Haematopus unicolor*) at two wintering sites in Canterbury, South Island, New Zealand. *Stilt* 40: 2–6. (Parks & Waterways Unit, Christchurch City Council, PO Box 267, Christchurch 8006, New Zealand. andrew.crossland@ccc.govt.nz). Substantial increase from 1988 onwards.

Crossland, A.C. 2002. Seasonal abundance of waterbirds at Tanah Merah Beach, a newly formed inter-tidal habitat on the edge of reclaimed land in Singapore. *Stilt* 42: 4–9. Most numerous: *Charadrius mongolus*, *Calidris ruficollis*, *Actitis hypoleucos*, *Charadrius peronii*, *Pluvialis squatarola*.

Dijken, L. & Koks, B. 2003. *Broedvogelmonitoring in het Nederlandse Waddengebied in 2002. [Monitoring breeding birds in the Dutch Wadden Sea area in 2002]*. SOVON-monitoring rapport 2003/03. Beek-Ubbergen: SOVON, 30 p. In Dutch with English summary. (SOVON, Rijksstraatweg 178, NL-6573 DG Beek-Ubbergen, Netherlands). Numbers and recent trends of breeding: *Himantopus himantopus*, *Haematopus ostralegus*, *Recurvirostra avosetta*, *Charadrius dubius*, *C. alexandrinus*, *C. hiaticula*, *Vanellus vanellus*, *Philomachus pugnax*, *Numenius arquata*, *Limosa limosa*, *Tringa totanus*.

Durell, S.E.A. le V. dit & Atkinson, P.W. 2004. Differential distribution of Oystercatchers *Haematopus ostralegus* overwintering on the Wash, east England. *Bird Study* 51(1): 76–82. (Ctr. Ecol. Hydrol. (CEH) Dorset, Winfrith Technol. Ctr., Dorchester, Dorset DT2 8ZD, UK.) (sld@ceh.ac.uk)

Elts, J., Kuresoo, A., Leibak, E., Leito, A., Lilleleht, V., Luigujo, L., Löhmus, A., Mägi, E. & Ots, M. 2003. (Status and numbers of Estonian birds, 1998–2002). *Hirundo* 16(2): 58–83. In Estonian with English summary. (Estonian Ornithol. Soc., pk. 227, EE50002 Tartu, Estonia. jaanus.elts@eoy.ee). Status, breeding numbers and trends for 23 wader species in Estonia.

Fefelov, I. 2003. Shorebirds of the Selenga River Delta, south-eastern Russia. *Stilt* 43: 40–41. (Res. Inst. Biol. Irkutsk Univ., PO Box 24, Irkutsk 664003, Russia.) (u000438@ic.isu.ru)



- Fischer, J. & Schweizer, M.** 2003. Ein adulter Weissbürzelstrandläufer *Calidris fuscicollis* im Chablais de Cudrefin/Fanel. (An adult White-rumped Sandpiper *Calidris fuscicollis* at Lake Neuchatel). *Ornithol. Beob.* 100(3): 247–250. In German with English summary. (Füllerichstr. 20, CH-3073 Gümligen, Switzerland.) (blackburnian@swissinfi.org)
- Fraser, P.A. & Rogers, M.J.** 2003. Report on scarce migrant birds in Britain in 2001. *Brit. Birds* 96(12): 626–649. (c/o Churchtown Cottages, Towednack, St Ives, Cornwall TR26 3AZ, UK). Waders: *Charadrius alexandrinus*, *Calidris temminckii*, *C. melanotos*, *Tringites subruficollis*, *Phalaropus lobatus*, *P. fulicarius*.
- Gerasimov, Yu.N. & Gerasimov, N.N.** 2002. Whimbrel *Numenius phaeopus* on Kamchatka, Russia. *Stilt* 41: 48–54. (Pacific Inst. Geograph., Russ. Acad. Sci, Petrovavlovsk, Kamchatka, Russia. bird@mail.kamchatka.ru). Migration and possible breeding.
- Gerasimov, Y.** 2003. Shorebird studies in north Kamchatka from July 5–August 12 2002. *Stilt* 44: 19–28.
- Gerritsen, G.J. & Tijssen, W.** 2003. (The importance of the Netherlands as staging area for Icelandic Black-tailed Godwit *Limosa limosa islandica* during spring migration in 2001 and 2002). *Limosa* 76(3): 103–108. In Dutch with English summary. (Veerallee 33, NL-8019 Ad Zwolle, The Netherlands. gerritgerritsen@hetnet.nl)
- Gillings, S. & Beaven, P.** 2004. Wintering farmland birds: results from mass-participation surveys. *Brit. Birds* 97(3): 118–129. (BTO, The Nunnery, Thetford, Norfolk IP24 2PU, UK). *Vanellus vanellus*.
- Gosbell, K., Collins, P. & Christie, M.** 2002. Wader surveys in The Coorong and coastal lakes of southeastern South Australia during February 2002. *Stilt* 42: 10–29. (17 Banksia Court, Heathmont, 3135 Vic., Australia.) (kenbg@ozemail.com.au)
- Hassell, C.J. & Rogers, D.I.** 2002. Painted Snipe nesting at Taylor's Lagoon near Broome, north-western Australia. *Stilt* 41: 14–21. (PO Box 3089, Broome, WA 6725, Australia. turnstone@wn.com.au). *Rostratula (benghalensis) australis*.
- Hedenström, A.** 2004. Migration and morphometrics of Temminck's Stint *Calidris temminckii* at Ottenby, southern Sweden. *Ring. & Migr.* 22(1): 51–58. (Dept Theor. Ecol., Ecol. Bldg., SE-223 62 Lund, Sweden.)
- Huettemann, F.** 2003. Shorebird migration on northern Sakhalin Island, Russia in early northern autumn. *Stilt* 43: 34–39. (Geograph. Dept, 2500 Univ. Dr. NW, Univ. Calgary, Calgary AB, T2N 1N4 Canada.) (falk@ucalgary.ca)
- Hulscher, J.B. & Verhulst, S.** 2003. (The rise and fall of the breeding population of Oystercatchers *Haematopus ostralegus* in Friesland in 1966–2000). *Limosa* 76(1): 11–22. In Dutch with English summary. (Zool. Lab., R.U. Groningen, PO Box 14, NL-9750 AA Haren, The Netherlands. j.hulscher@tiscali.nl)
- Islam, M.S. & Islam, M.Z.** 2002. Abundance of migratory shorebirds on the coast of Bangladesh. *Stilt* 41: 22–26. (House No-15/22, Munshi Bari South Chartha, Comilla 3500 Bangladesh. msislam2000@yahoo.com). 34 species counted during a mid winter count. *Charadrius alexandrinus* most abundant.
- Kaczorowski, G.** 2001. (First record of the Pacific Golden Plover *Pluvialis fulva* in Poland). *Notatki Ornitol.* 42(2): 143–144. In Polish with English summary. (Pukarow 40, PL-97-532 Zytno, Poland)
- Keates, A.** 2003. A survey of Bar-tailed Godwits in south-east Queensland. *Stilt* 44: 50–51. (33 Beach Rd., Nightcliff, NT 0761, Australia. tattlers@tpi.com.au). *Limosa lapponica baueri*, L.l. menzbieri.
- Koffijberg, K., Blew, J., Eskildsen, K., Günther, K., Koks, B., Laursen, K., Rasmussen, L.-M., Potel, P. & Südbbeck, P.** 2004. *High tide roosts in the Wadden Sea: a review of bird distribution, protection regimes and potential sources of anthropogenic disturbance*. A report of the Wadden Sea Plan Project 34. Wadden Sea Ecosystem No. 16. Wilhelmshaven: Common Wadden Sea Secretariat, 120 p. Detailed overview and distribution maps for each species.
- Kolodziejczyk, P. & Pola, A.** 2002. (First record of the Great Knot *Calidris tenuirostris* in Poland). *Notatki Ornitol.* 43(2): 113–118. In Polish with English summary. (Lotnicza 139/49, PL-54-132 Wrocław, Poland.)
- Kuresoo, A. & Luigujo, L.** 2003. The Great Snipe and its conservation in Estonia. *Hirundo*, suppl. 7: 29–56. (EPMU, Zool. Botaanika Inst., Riia 181, 51014 Tartu, Estonia. akuresoo@zbi.ee). *Gallinago media*, distribution, number and biology, threats and conservation.
- Leader, P.J. & Carey, G.J.** 2003. Identification of Pintail Snipe and Swinhoe's Snipe. *Brit. Birds* 96(4): 178–198. (Asia Ecol. Consultants Ltd., 127 Commercial Ctr., Palm Springs, Yuen Long, New Territories, Hong Kong). *Gallinago stenura*, *G. megala*. Most individuals are not safely identifiable in the field owing to the overlap in size, structure and plumage.
- Lei, G., Jiang, Y. & Yao, Y.** 2002. Waders of East Dongting lake National Nature Reserve, Hunan province, China. *Stilt* 41: 11–13. (East Dongting Natl. Nature Reserve, Hunan Province, P.R. China, 41400.) (glei1126@sina.com.cn). 17 species counted. *Tringa erythropus*, *Recurvirostra avosetta*, *Calidris alpina* most numerous.
- Ma, Z.J., Jing, K., Tang, S.M. & Chen, J.K.** 2002. Shorebirds in the eastern intertidal areas of Chongming Island during the 2001 northward migration. *Stilt* 41: 6–10. (Min. Educ. Key Lab. Biodiv. Sci. & Ecol. Engin., Inst. Biodiv. Sci., Fudan Univ., Shanghai, 200433, P.R. China). 44 904 shorebirds counted of 32 species. *Calidris alpina*, *C. canutus*, *Charadrius alexandrinus* most numerous.
- Ma, Z.J., Tang, S.M., Lu, F. & Chen, J.K.** 2002. Chongming Island: a less important shorebird stopover site during southward migration. *Stilt* 41: 35–37.
- Meissner, W.** 2003. Biometrics, length of stay and body mass increase of migrating Common Snipes *Gallinago gallinago* in the Gulf of Gdansk. *Vogelwelt* 124(1): 45–52. (Dept Vertebrat. Ecol. Zool., Univ. Gdansk, Al. Legionow 9, PL-80-441 Gdansk, Poland.) (biowm@univ.gda.pl)
- Minton, C.** 2002. Curlew Sandpiper transfers to the South African Flyway. *Stilt* 42: 37. (165 Dalgetty Rd., Beaumaris, Vic. 3193, Australia.) (mintons@ozemail.com.au). *Calidris ferruginea*.
- Minton, C., Jessop, R., Collins, P., Hassell, C., Deleyev, J. & Beasley, L.** 2002. Sightings of waders and terns leg-flagged in N.W. Australia: report number 7. *Stilt* 42: 38–51. Yellow-flagged waders seen in Korea, China, Taiwan, Russia, New Zealand, Japan.
- Minton, C., Jessop, R., Collins, P., Deleyev, J. & Beasley, L.** 2002. Sightings of waders leg-flagged in South Australia:



- report no. 1. *Stilt* 42: 52–55. Leg-flagged waders seen in Japan, New Zealand, Russia.
- Minton, C., Jessop, R., Collins, P., Deleyev, J. & Beasley, L.** 2002. Sightings of waders leg-flagged in Victoria: report no. 9. *Stilt* 42: 56–72. Leg-flagged waders seen in Japan, New Zealand, Russia, USA, China, Korea, Taiwan.
- Minton, C., Jessop, R., Collins, P. & Sitters, H.** 2003. Northwest Australia wader and tern expedition from 1 August to 1 November 1998. *Stilt* 43: 42–46.
- Minton, C., Jessop, R. & Collins, P.** 2003. Northwest Australia wader and tern expedition from 15 May to 3 June 2000. *Stilt* 43: 47–50.
- Minton, C., Collins, P. & Sitters, H.** 2003. Northwest Australia wader and tern expedition from 30 December 2000 to 30 January 2001. *Stilt* 43: 51–54.
- Minton, C., Jessop, R. & Collins, P.** 2003. Northwest Australia wader and tern expedition 15 September to 17 November 2001. *Stilt* 43: 55–67.
- Minton, C., Jessop, R. & Collins, P.** 2003. A summary of the results of the northwestern Australia wader and tern expedition from 28 June to 19 July 2003. *Stilt* 44: 64–72.
- Morozov, V.V.** 2004. Displaying Swinhoe's Snipe in eastern European Russia: a new species for Europe. *Brit. Birds* 97(3): 134–138. (Zool. Mus. Moscow State Univ., Bolshaya Nikitskaya St. 6, 125009 Moscow, Russia). *Gallinago megala*.
- Morrison, R.I.G., Ross, R.K. & Niles, L.J.** 2004. Declines in wintering populations of Red Knots in southern South America. *Condor* 106(1): 60–70. (Can. Wildl. Serv., Natl. Wildl. Res. Ctr., Carleton Univ., Raven Rd., ON K1A 0H3, Canada. guy.morrison@ec.gc.ca). *Calidris canutus rufa*.
- Nijland, F.** 2002. (Breeding populations of meadowbirds in the province of Friesland in 1999). *Limosa* 75(4): 169–172. In Dutch with English summary. (Weidevogelmeetnet Friesland, Emmakade 107, NL-8921 AH Leeuwarden, The Netherlands. wmf@planet.nl). Population estimates for *Haematopus ostralegus*, *Vanellus vanellus*, *Limosa limosa*, *Tringa totanus*, *Gallinago gallinago*, *Philomachus pugnax*.
- Ogilvie, M. & Rare Breeding Birds Panel.** 2003. Rare breeding birds in the United Kingdom in 2001. *Brit. Birds* 96(10): 476–519. (Glencairn, Bruichladdich, Isle of Islay PA49 7UN, UK). Waders: *Recurvirostra avosetta*, *Burhinus oedicnemus*, *Charadrius dubius*, *C. morinellus*, *C. maritima*, *Philomachus pugnax*, *Limosa limosa*, *Numenius phaeopus*, *Tringa nebularia*, *T. ochropus*, *T. glareola*, *Phalaropus lobatus*.
- Ozarowski, D.** 2000. (Distribution and abundance of breeding waders *Charadrii* at the Bay of Gdansk in 1991–1992). *Notatki Ornitol.* 41(2): 172–176. In Polish with English summary. (Rozewska 10/14, PL-81-055, Gdynia, Poland). *Haematopus ostralegus*, *Charadrius dubius*, *Ch. hiaticula*, *Vanellus vanellus*, *Calidris alpina*, *Gallinago gallinago*, *Tringa totanus*.
- Pugacewicz, E.** 2002. (Numbers and distribution of the Great Snipe *Gallinago media* in the Nizina Polnocnopodlaska lowland in 1976–2000). *Notatki Ornitol.* 43(1): 1–7. In Polish with English summary. (Botaniczna 3, PL-17-200 Hajnowka, Poland)
- Rehfisch, M., Austin, G. & Musgrove, A.** 2003. Wintering waders in decline. *BTO News* 248: 16–17. *Haematopus ostralegus*, *Recurvirostra avosetta*, *Charadrius hiaticula*, *Pluvialis squatarola*, *Calidris canutus*, *C. alba*, *C. maritima*, *C. alpina*, *Limosa limosa*, *L. lapponica*, *Numenius arquata*, *Tringa erythropus*, *T. totanus*, *T. nebularia*, *Arenaria interpres*.
- Rehfisch, M.M., Holloway, S.J. & Austin, G.E.** 2003. Population estimates of waders on the non-estuarine coasts of the UK and the Isle of Man during the winter of 1997–98. *Bird Study* 50(1): 22–32. (BTO, The Nunnery, Thetford, Norfolk IP24 2PU, UK). Decline of *Charadrius hiaticula*, *Calidris alba*, *C. maritima*, *Limosa lapponica*, *Arenaria interpres*.
- Riak, K.M., Ismail, A., Arshad, A. & Ismail, A.R.** 2003. Species composition and use of mudflat of Kapar, west coast of peninsular Malaysia by migratory shorebirds. *Stilt* 44: 44–49. (Dept Biol., Fac. Sci. & Environm. Stud., Univ. Putra Malaysia, 43400 UPM Serdang, Malaysia.) (khalidriak@yahoo.co.uk). Most abundant: *Charadrius mongolus*, *Tringa totanus*, *Limosa limosa*, *Numenius arquata*, *Limosa lapponica*, *Numenius phaeopus*.
- Rogers, M.J. & Rarities Committee.** 2003. Report on rare birds in Great Britain in 2002. *Brit. Birds* 96(11): 542–609. (2 Churchtown Cottages, Towednack, St. Ives, Cornwall TR26 3AZ, UK). Waders: *Himantopus himantopus*, *Charadrius vociferus*, *C. mongolus*, *Pluvialis dominica*, *P. fulva*, *Vanellus gregarius*, *Calidris pusilla*, *C. minutilla*, *C. fuscicollis*, *C. acuminata*, *Limicola falcinellus*, *Micro-palama himantopus*, *Gallinago media*, *Limnodromus scolopaceus*, *Numenius phaeopus hudsonicus*, *Bartramia longicauda*, *Tringa stagnatilis*, *T. melanoleuca*, *T. flavipes*, *T. solitaria*, *Xenus cinereus*, *Actitis macularia*, *Phalaropus tricolor*.
- Schweizer, M.** 2003. Seltene Vogelarten und ungewöhnliche Vogelbeobachtungen in der Schweiz im Jahre 2002. (Rare and unusual observations in Switzerland in 2002). *Ornithol. Beob.* 100(4): 293–314. In German with English summary. (Gartenstr. 2, CH-3176 Neuenegg, Switzerland. schw@bluewin.ch). Waders: *Glareola pratincola*, *Charadrius dubius*, *Chettusia gregaria*, *Calidris minuta*, *C. fuscicollis*, *Limicola falcinellus*, *Gallinago media*, *Tringa nebularia*, *Phalaropus lobatus*, *P. fulicarius*.
- Scott, I., Mitchell, P.I., Gudmundsson, G.A., Eaton, M., Ward, R.M. & Evans, P.R.** 2004. Using radio-transmitters to help monitor the spring migration of Sanderling *Calidris alba*: why do some birds stop in Iceland? *Bird Study* 51(1): 83–86. (Health Care Education Development Unit, St. Bartholomew Sch. Nursing Midwifery, City Univ., 20 Bartholomew Close, London, EC1A 7QN, UK.)
- Singor, M.** 2003. Wader observations at the Serpentine River Reserve, Western Australia. *Stilt* 44: 9–18. (149A Bishopsgate St., Carlisle 6101 WA, Australia.) (msingor@iprimus.com.au). Survey of waders wintering 1996–2003.
- Skewes, J.** 2002. Report on the 2001 population monitoring counts. *Stilt* 41: 55–61. (13 Waterloo Str., Heathmont, Vic. 3139, Australia.)
- Skewes, J.** 2003. Report on population monitoring counts, 2002. *Stilt* 44: 56–63. (13 Waterloo Str., Heathmont, Vic. 3139, Australia.)
- Smith, V.** 2002. Important wader sites in the Asian-Australian Flyway: Oyster Harbour and Princess Royal Harbour, Albany, Western Australia. *Stilt* 41: 40–43. (1 Karrakatta Rd., Goode Beach, Albany, WA 6330 Australia.)
- Stawarczyk, T.** 2000. (First record of the Spotted Sandpiper *Actitis macularia* in Poland). *Notatki Ornitol.* 41(4): 326–327. In Polish with English summary. (Mus. Przyrodnicze



- Univ. Wroclawskiego, Sienkiewicza 21, PL-50-335 Wroclaw, Poland. stawar@biol.uni.wroc.pl)
- Steiof, K.** 2003. Ergänzende Hinweise zur Bestandsaufnahme beim Waldwasserläufer *Tringa ochropus*. (Complementary notes on Green Sandpiper *Tringa ochropus* survey methods). *Vogelwelt* 124(1): 53–56. (Seestr. 60, D-13347 Berlin, Germany. k.steiof@web.de)
- Thyen, S. & Exo, K.-M.** 2003. [The importance of salt marshes of the German Wadden Sea off Lower Saxony for the reproduction of Redshanks *Tringa totanus*]. In: Schutz von Feuchtgrünland für Wiesenvögel in Deutschland. Tagungsbericht NABU, Bergenhausen, p. 20–26. In German with English summary. (Inst. Vogelforschung ‘Vogelwarte Helgoland’, An der Vogelwarte 21, D-26386 Wilhelmshaven, Germany.) (michael.exo@ifv.terramare.de)
- Tomkovich, P.S.** 2003. List of wader species of Chukotka, northern Far East of Russia: their banding and migratory links. *Stilt* 44: 29–43. (Zool. Mus. Moscow State Univ., Bolshaya Nikitskaya St. 6, Moscow, 125009, Russia. pst@zmmu.msu.ru).
- Trolliet, B.** 2003. Elements for a Lapwing (*Vanellus vanellus*) management plan. *Game & Wildl. Sci.* 20(1–2): 93–144. (Office Natl. de la Chasse et de la Faune Sauvage, Chanteloup, F-85340 Ile d’Olonne, France.) (bertrand.trolliet@oncfs.gouv.fr). Distribution and movements, population size and trends, population dynamics, habitat and ecology, hunting, legal status and conservation measures, and recommendations for management, conservation, research and monitoring.
- Tuule, E., Tuule, A. & Elts, J.** 2003. (Numbers and population dynamics of the Curlew in the surroundings of Saue, 1963–2002). *Hirundo* 16(1): 14–22. *Numenius arquata*. In Estonian with English summary. (Sütiste tee 3–61, EE13419, Tallinn, Estonia. etuule@hot.ee)
- Volet, B. & Burkhardt, M.** 2003. Übersicht über das Brutgeschehen und andere ornithologische Ereignisse 2002 in der Schweiz. (Rare and unusual records of breeding, migrating and wintering bird species in Switzerland, 2002). *Ornithol. Beob.* 100(4): 323–334. In German with English and French summary. (Schweizerische Vogelwarte, CH-6204 Sempach, Switzerland.) (id@vogelwarte.ch). Waders: *Charadrius dubius*, *Gallinago gallinago*.
- Wallace, D.I.M.** 2003. Shortish or long shot at snipe? *Brit. Birds* 96(8): 406–407. (Mount Pleasant Farm, Main Rd., Anslow, Burton-on-Trent, East Staffordshire DE13 9QE, UK). *Gallinago stenura*, *G. megala*. Comment on identification features.
- Ward, R.M., Evans, P.R. & O’Connell.** 2002. Study of long term changes in bird usage of the Tees Estuary. Report to English Nature. WWT Res. Dept, Slimbridge. (Wildfowl & Wetlands Trust, Res. Dept, Slimbridge, Glos GL2 7BT, UK. robin.ward@wwt.org.uk). Decline in wintering populations of *Calidris canutus*, *Pluvialis squatarola*, *Limosa lapponica*, *Calidris alpina*.
- Wilson, A. & Vickery, J.** 2003. Wet grassland waders in decline. *BTO News* 247: 12–13. *Haematopus ostralegus*, *Vanellus vanellus*, *Gallinago gallinago*, *Numenius arquata*, *Tringa totanus*.
- Wilson, J.R.** 2001. Wader surveys in The Coorong, South Australia in January and February 2001. *Stilt* 40: 38–54. (Granveien 46, 1911 Flateby, Norway).
- Wilson, J.R.** 2001. The January and February 2001 Victoria wader count. *Stilt* 40: 55–64.
- Zielinski, P., Mokwa, T., Krol, J., Malczyk, P., Nowak, D., Gorecki, D. & Włodarczak, A.** 2001. (Semipalmated Sandpiper *Calidris pusilla*, a new species for Poland. Identification problems). *Notatki Ornitol.* 42(4): 275–282. In Polish with English summary. (Biol. Stn. IE PAN, Nadwiślańska 108, PL-80-680 Gdańsk 40, Poland)

MISCELLANEOUS

- Bearhop, S., Ward, R.M. & Evans, P.R.** 2003. Long-term survival rates in colour-ringed shorebirds – practical considerations in the application of mark-recapture models. *Bird Study* 50(3): 271–279. (Graham Kerr Bldg., Div. Environm. Evol. Biol., Univ. Glasgow.) (sbearhop@udcf.gla.ac.uk). *Pluvialis squatarola*, *Arenaria interpres*, *Calidris alba*, *Numenius arquata*, *Limosa lapponica*.
- Bellebaum, J. & Boschert, M.** 2003. Bestimmung von Predatoren an Nestern von Wiesenlimikolen. (Determining predators at nests of meadow birds). *Vogelwelt* 124(2): 83–91. In German with English summary. (Dorfstr. 13, D-16248 Bölkendorf, Germany.) (Jochen.Bellebaum@t-online.de)
- Coles, R.K.** 2003. Nest protection of Little Ringed Plovers. *Brit. Birds* 96(7): 346. (12 Lennard Rd., Dunton Green, Sevenoaks, Kent TN13 2UV, UK). *Charadrius dubius*. Using a cage to avoid predation.
- Godfrey, T.** 2003. Nest protection of Little Ringed Plovers. *Brit. Birds* 96(9): 457. (7 Lynton Lane, Cassington, Oxfordshire OX29 4ES, UK). *Charadrius dubius* and nesting cages of bicycle baskets.
- Price, C. & Baker, J.** 2003. Wail and bark. *Wingspan* 13(4): 10–15. *Burhinus grallarius*, protection measures in NSW, Australia.
- Sykes, T.** 2003. Great Ringed Plover smothered by sand. *Brit. Birds* 96(9): 456–457. (Gibraltar Point Natl. Reserve, Skegness, Lincolnshire PE24 4SU, UK). *Charadrius hiaticula* nesting under a wire-mesh cage buried by wind-blown sand.
- Thyen, S. & Exo, K.-M.** 2003. (Ecological development of a re-aggregating clay pit in a salt marsh of the western Jade Bay). *Vogelkdl. Ber. Niedersachs.* 35: 143–150. In German with English summary. (Inst. Vogelforschung ‘Vogelwarte Helgoland’, An der Vogelwarte 21, D-26386 Wilhelmshaven, Germany.) (michael.exo@ifv.terramare.de)
- Thyen, S. & Exo, K.-M.** 2003. (Salt marsh succession at the coast of Lower Saxony, Germany: chance or risk for breeding birds?). *Vogelkdl. Ber. Niedersachs.* 35: 173–178. In German with English summary.
- Triplet, P., Sournia, A., Joyeux, E. & Le Dréan-Quénec'hdu, S.** 2003. (Human disturbances to wildlife: the case of waterbirds). *Alauda* 71(3): 305–316. In French with English summary. (SMACOPIRN Baie de Somme, Maison Ramsar, 1, Place de l’Amiral Courbet, F-80100 Abbeville, France. patricktriplet@baiedesomme.org).
- Weston, M. & Paton, S.** 2002. A survey of Victorian Hooded Plover counters. *Stilt* 42: 30–31. (Threatened Bird Network, Birds Australia Natl. Off., 415 Riversdale Rd., Hawthorn East, Vic. 3123, Australia). On survey techniques, motivations, methodology, and needs of counters.

