

sented gives corroborative evidence of at least a southerly movement of the species even on the part of first-year individuals.—DAYTON STONER, New York State Museum, Albany, New York.

Indigo Bunting Returns-1 and Returns-2 at Nashville, Tennessee

Year	Number of Birds Banded	Number Returning
1932	15	0
1933	16	0
1934	42	2
1935	51	17
1936 (to July 15th)	22	19

RECORD OF INDIVIDUALS

Band No.	Age—Sex	Date of Banding	Return-1	Return-2	Repeats
H73023	♀	July 6, 1933	September 9, 1934		7
H73038	♂	August 30, 1933	May 14, 1935	May 14, 1936	17
H73045	Ad. ♂	October 9, 1933	May 19, 1934	May 20, 1935	6
34-8106	" ♂	May 16, 1934	May 6, 1935		14
H73131	" ♂	June 13, 1934	May 13, 1935	June 5, 1936	28
H73153	" ♀	July 13, 1934	July 10, 1936		3
34-8139	Im. ♀	July 25, 1934	May 10, 1935		3
34-8142	" ♂	July 28, 1934	May 6, 1935	May 4, 1936	50
34-8147	" ♂	August 5, 1934	April 29, 1935	April 27, 1936	99
34-8148	" ♂	August 6, 1934	May 1, 1935		13
34-8151	" ♂	August 6, 1934	April 29, 1935	May 5, 1936	9
34-8162	" ♀	August 14, 1934	May 17, 1935		2
34-8169	" ♀	August 16, 1934	July 9, 1935	May 15, 1936	37
34-8180	Ad. ♀	August 21, 1934	July 1, 1936		5
34-8199	" ♂	August 31, 1934	July 28, 1935	July 4, 1936	2
34-8200	Im. ♀	August 31, 1934	September 11, 1935		..
34-8204	♀	September 9, 1934	June 10, 1935	June 11, 1936	12
34-8205	Ad. ♂	September 9, 1934	September 25, 1935		..
34-8213	" ♀	September 19, 1934	May 23, 1935		..
34-8226	Ad. ♂	September 25, 1934	July 12, 1935		1
34-87190	Yg. ♂	May 17, 1935	May 14, 1936		41
34-87194	Ad. ♂	May 17, 1935	May 14, 1936		40
34-87205	" ♂	May 13, 1935	May 6, 1936		20
34-87209	" ♂	May 20, 1935	May 26, 1936		6
34-87213	" ♂	May 23, 1935	May 5, 1936		19
34-87235	" ♂	July 3, 1935	April 30, 1936		17
35-7805	" ♀	July 28, 1935	May 5, 1936		15
35-7839	Im. ♂	August 21, 1935	May 4, 1936		31
35-7855	♀	August 26, 1935	June 11, 1936		18

Excepting a small percentage taken in water-drip traps, all captures of Indigo Buntings (*Passerina cyanea*) have been made with millet seed as bait. Potter type, modified Government sparrow, and top-opening funnel traps have been about equally successful. In 1934, they were attracted to the banding station in numbers when the millet seed ripened in a little patch that was planted as a lure, resulting in a considerable increase in banded birds for that year. None has been planted since then, but quantities are kept at the traps continually in various parts of the garden.—AMELIA R. LASKEY, Nashville, Tennessee.

A Pair of Northern Crested Flycatchers (*Myiarchus crinitus boreus*) Mated for Three Consecutive Years. The following nesting records were made at my banding station in Norristown, Pennsylvania:

A147214	—Adult banded	June 30, 1929
	Return-1	June 22, 1933
	" -2	July 8, 1934
	" -3	June 23, 1935
	" -4	June 27, 1936
34-105309	—Adult banded	July 8, 1934
	Return-1	June 23, 1935
	" -2	June 27, 1936

This pair of birds was taken on each of the years 1934, 1935 and 1936 in the nesting box while feeding half-grown young on the dates above given, being mated together for three consecutive years. A147214 is now at least 8 years old.