central southern region



STEPHEN J. STEDMAN

We arm conditions prevailed this winter, but rainfall was less uniform with some parts of the Region drier and others wetter than the "norm." Many new inland wintering records were set, presumably as a result of these conditions, the origin of which remains uncertain, but increased atmospheric carbon dioxide emissions might be part of the answer (see *Nature* 382: 146–149). Interesting avian records occurred offshore from Louisiana this winter; be sure to note the species you can encounter if you take a winter cruise on the Gulf of Mexico—or reside on an oildrilling platform for a few weeks.

LOONS THROUGH DUCKS

Mississippi obtained its first Red-throated Loon specimen when one was found dead in Harrison Jan. 24 (fide TLS*). One Redthroated and two Pacific loons in Henry, TN, Feb. 7 (†JRW, MW) visited a site where one of the latter was also noted Dec. 5 and Feb. 21. Alabama had 3 reports of Pacific Loons, including inland sightings in Marshall Dec. 10 (†GDJ, DGJ) and Laurence Jan. 23 (†SWM, JRW et al.). Single Eared Grebes in Washington and Clark, AR, Feb. 11 (MM et al.) and 15 (HP, MP), respectively, were locally rare as was a single inland in Marshall, AL, Dec. 3-17 (†RAR, LBR); one-three were in Panola, MS, Jan. 5-23 (GK, SK, WMD) while a single briefly visited Oktibbeha, MS, Jan. 23 (TLS, MS, MC). Western Grebes stayed in Faulkner and Hempstead, AR, Nov. 28-Feb. 6 (KN, LN et al.) and Dec. 5-6 (CM), respectively.

Alabama's 9th Sooty Shearwater appeared briefly in *Baldwin* on the Gulf Shores



Representing the third winter record for west Tennessee, this American Bittern was found in Black Bayou, Lake County, December 19, 1998, during the Reelfoot Lake CBC. Photograph/Jeff R. Wilson

CBC Jan. 2 (†GDJ); a small, unidentified shearwater (ALM) and at least one unidentified booby (RRR) were also sighted on that count. American White Pelicans were again abundant in Mississippi where 1430 were registered Dec. 19 in *Panola/Lafayette* (WMD, GK).

Western Tennessee's 3rd winter Am. Bittern was found on the Reelfoot L. CBC, Lake, Dec. 19 (JZ, NM et al.); only the 2nd locally and first in winter, another was noted in Putnam, TN, Dec. 9 (GKE, SJS); locally very rare in winter, one also visited Clark, AR, Dec. 31 (RHD). Six Great Egrets over the Gulf of Mexico 135 mi s. of Morgan City, LA, Feb. 13 (DNP) were somewhat unexpected though associated with a cold front. Rarely seen in n.w. Louisiana during January, and more rarely still by the dozen, 12 Snowy Egrets were in DeSoto Jan. 10 (JfT, JnT). Four Tricolored Herons in Nachitoches, LA, Feb. 6 (RJS, VL) were also unexpected winter voyageurs to n.w. Louisiana. Western Tennessee's first winter Cattle Egret was present in Lake Feb. 12 (WGC); one in Clark Feb. 15 (HP, MP) and two in Cleburne Feb. 18 (KN, LN) were considered early migrants in Arkansas; the 2nd sighting for Alabama's Tennessee Valley

was established by one in Colbert/ Lauderdale Jan. 24 (JH, m. ob.); another single in Caddo, LA, Jan. 31 (CL, JK, JoK) rounded out records of this northward-roving heron during the winter. Further evidence of the eastward expansion of Roseate Spoonbill was provided by two in *Plaquemines*, LA, Jan. 2 (DPM et al.).

Fulvous Whistling-Ducks were noted in 3 w. Louisiana parishes while reports of Black-bellied Whistling-Ducks were restricted to 2 sightings in Cameron, LA. Continuing a late fall sighting, an imm. Greater White-fronted Goose in Putnam, TN, Dec. 1-16 (WAW, BHS, m.ob.) was first locally; up to 17 in Okaloosa, FL, Dec. 1-Jan. 21 (DW, RAD, m.ob.) continued fall sightings and were locally rare; Alabama had 3 sightings of small groups; flocks of 5000 in Tunica, MS, Jan. 17 (JRW) and 8000 in Tallahatchie/Quitman, MS, Jan. 4 & 24 (FB, JeL) were certainly notable; groups of 25 & 2 in Iberville, LA, Feb 21 & 22 (DLD, SWC) were locally rare. A flock of up to 230 Snow Geese in Okaloosa, FL (RAD, m.ob.), through the period represented a locally unprecedented number, and 220 in Baldwin, AL, Jan. 2 (GDJ, DGJ) were considered a good coastal aggregation; 50,000 in

Tunica, MS, Jan. 17 (JRW) and 100,000 in Tallahatchie/Quitman Feb. 6 (GK, SK) represented large numbers at those locales. The species' search for wintering sites to accommodate its burgeoning continental population is probably indicated by these records. Continuing late fall sightings, an imm. Ross's Goose in Putnam, TN, Dec. 1-Jan. 12 (SJS, m.ob.) and an immature and three adults in Okaloosa, FL, Dec. 1-Feb. 23 (RAD, PCT, PB v.t., m.ob.) provided first local records; singles were noted in 3 inland Alabama counties while a group of three appeared in Baldwin Jan. 2 (GDJ, DGJ), the 3rd record for the Gulf Coast in that state. Mainly small groups showed up in 6 Mississippi counties with 250 in Tunica Jan. 17 (IRW) constituting by far the largest; a single in Orleans, LA, Jan. 24+ (DPM et al.) was rare at the location.

a superior to be a superior and the second second

A male Cinnamon Teal was probably a returnee from the previous winter to Orleans, LA, being noted late fall to Feb. 14 (GO et al.); two other males were in Cameron Dec. 20 (DP, JW, MWe, MA). Sixhundred Ring-necked Ducks in Lauderdale Feb. 8 (PAK) tied the high count for n.w. Alabama. An imm. Black and three Surf scoters stayed in Yell, AR, Jan. 16-30 (KN, LN) while two Blacks visited Baldwin, AL, Dec. 14 (GDJ, DGJ); Mississippi birders recorded Black Scoters in 3 countiesincluding two in Oktibbeha Dec. 13 (TLS, MS, m.ob.), one of which stayed until Feb. 12-and Surfs in three. Second locally in winter, a White-winged Scoter was noted briefly in Putnam, TN, Jan. 9 (SJS) while another showed up in Dyer, TN, Feb. 14 (WGC); a female in Yell, AR, Jan. 16-30 (KN, LN) created a clean sweep of scoters in one inland county in one season. Yell also hosted 350 Com. Goldeneyes Jan. 9 (KN, LN), a sizable flock evenly divided by sex. Sixteen Com. Mergansers in Henry, TN, Feb. 14 (JRW) represented a good count; seven were in Cleburne, AR, Jan. 5 (KN, LN). Mississippi had singles in Tunica and Yalobusha. An Alabama high count for Hooded Merganser was established Dec. 19 when 1057 were counted on the Wheeler N.W.R. CBC (fide GDJ).

RAPTORS THROUGH RAILS

Running late or wintering, an Osprey was noted in *Columbia*, AR, Dec. 19 (DC); late or wintering, singles were also sighted in 3 ınland Mississippi counties. Six Louisiana parishes hosted White-tailed Kites during the season, including s.e. *Plaquemines*. The mid-winter eagle survey in Tennessee turned up 244 in the west and 92 in the middle portion of the state, somewhat higher numbers than the long-term averages; Reelfoot L. was 95% iced over during the counting period (Jan. 1–15), reducing eagles counted there and probably increasing numbers on the Mississippi R. and Kentucky L./Tennessee R. (*fide* RMH). By the end of the period 7 active Bald Eagles nests were at Reelfoot L., TN (DH).

Judging from Regional CBC data (Table 1), Accipiters have been increasing nicely during the last decade. An imm. Broadwinged Hawk was studied at leisure in Plaquemines, LA, Jan. 31 (PW, DPM, RDP); early arriving, another was in Tipton, TN, Feb. 27 (†JRW). Fifth locally, a Swainson's Hawk ventured into Okaloosa, FL, Dec. 14 (†DW); two well-described individuals were in Vermilion, LA, Dec. 18 (CWi, AS, PC). A "Harlan's" Red-tailed Hawk returned to the same wintering site in Putnam, TN, for the 12th winter (SJS); others were reported in Fayette, Maury, and Shelby, TN. Very rare in Alabama, an ad. rufous-morph Red-tail was noted in Blount Dec. 6 (GDJ, DGJ). On the Mississippi Review List, a Rough-legged Hawk was identified in Sharkey Feb. 4 & 18 (PBH, CW; p.a.); another was notable in Nachitoches, LA, Jan. 10-Feb. 21 (RJS, m.ob.). Quite a few Golden Eagles were reported from Arkansas, Louisiana, and Mississippi. Back again in Calcasieu, LA, a Crested Caracara was noted Dec. 26 (JK, DR) and Feb. 20 (DCl, SC) while another was noted in Cameron, LA, Dec. 29 and Jan. 16 (JK, JoK, KF). Single Peregrines were observed over the Gulf of Mexico 135 mi s. of Morgan City, LA, Jan. 24 and Feb. 12 (DNP). The Yalobusha, MS, Prairie Falcon made it back for the 8th winter (GK, SK, FB et al.).

The northward migration of Sandhill Cranes in Tennessee peaked in mid-February for the 2nd consecutive year, about 2 weeks earlier than expected; in *Putnam*, TN, crane flocks were also noted for the first time in January, some going N and some S (SJS, JCS, GKE); sizable wintering crane flocks, including 300+ in *Obion* Dec. 16–Feb. 13 (WGC), persisted again in Tennessee (also see Appalachian Region); a new maximum count for the Alabama Gulf Coast was established when 35 were observed Jan. 2 in *Baldwin* (WJB, GB); other sightings took place in 4 Mississippi counties and 4 Louisiana parishes.

The first w. Tennessee winter record of King Rail was established Jan. 16–17 in *Lake* (JRW, WGC, NM); singles were also noted in *Leflore* and *Rankin*, MS, as well as *Nachitoches*, LA.

SHOREBIRDS THROUGH TERNS

Very rare inland, eight Semiplamated Plovers were counted in Jeff Davis, LA, Dec. 29 (DPM, PW, LCB, DP). A new Alabama maximum count of 35 Am. Oystercatchers took place in Mobile Feb. 8 (JoP, JP). Dyer, TN, provided the first w. Tennessee winter record of Am. Avocet when a single was noted Dec. 1-4 (WGC); also rare was one in Limestone, AL, Jan. 16 (HD). Alabama's 7th winter Whimbrel was noted in Mobile Feb 8 (JoP, JP); another was very rare inland in Cameron, LA, Dec. 29 (JFH). A Long-billed Curlew was noted Feb. 27 in Lafourche, LA (CK), where the species is scarce. Solitary Sandpipers were observed in 3 Louisiana parishes, where wintering is rare. A lingering Spotted Sandpiper in Putnam, TN, Dec 1-19 (SJS, BHS, LP, GKE) provided the first local winter record; one was rare in Jefferson, AR, Dec. 26 (RHD, JLY); still another was considered rare in Panola, MS, Jan. 9-Feb. 13 (GK, SK, WMD). Alabama's first inland winter Sanderling stayed the season in Limestone (SWM) where 10+ W Sandpipers were also located Dec. 27 (SWM et al.). Alabama's first Purple Sandpiper regaled a crowd in Baldwin Jan. 1-3 (JPe, TW; p.a. Alabama Bird Records Committee). "Impressive" is a good way to describe 300+ Stilt Sandpipers counted in Lafourche, LA, Jan. 17 (PW, DPM, MMy) Reliable inland records of Short-billed Dowitcher are rare, so 15-20 Dec. 29 in Jeff Davis, LA (DPM, PW), were notable. Longbilled Dowitchers seldom winter in Alabama, making a single Jan. 2 in Baldwin (GDJ, DGJ) a nice find while three in Limestone Jan. 22 provided the first record in n Alabama (SWM); flocks of 60 and 20 Dec 1 and Feb. 25, respectively, in Quitman, MS (FB), were probably migrants.

A dark-morph Pomarine Jaeger was well studied in Harrison, MS, Jan. 24 (JB, SJP et al.); three Pomarines in Cameron, LA, Dec 19 (SWC, DLD) and another there Dec. 21 (DP, JW) were the only others reported Staying Dec. 5–Jan. 23, a Laughing Gull was noteworthy in Pope/Yell, AR (KN, LN), another inland sighting took place in East Baton Rouge, LA, Jan. 1 (DCl, SC). An ad Franklin's Gull was noted Jan. 1 in Davidson, TN (CS), where rare; another was in Crawford, AR, Feb. 20 (KN, LN); and still another was in Noxubee, MS, Dec. 11 (TLS), five in Baldwin Dec. 2 (GDJ, DGJ) were notable while an adult stayed appropriately in Franklin, FL, Dec. 24-28 (†TL, DM, GM). Louisiana had singles in Cameron Dec. 19 (SWC, DLD) and Jeff Davis Dec. 29 (RB et al.).

central southern

Sixty-thousand Ring-billed Gulls in Yell, AR, Jan. 9 (KN, LN) must have taken a spell to count. Two California Gulls, still very rare in the Region, were in Cameron Dec. 21 (SWC, DLD). Lingering from fall, a Lesser Black-backed Gull stayed in Henry, TN, to Jan. 9 (JRW, m.ob.); one was found in Hancock, MS, Jan. 23 (JB, SJP); Louisiana birds included one in Cameron Dec. 20 (PW, MMy, CSo) and two in Jefferson Jan. 23-late February (PY); and four were noted in 3 Alabama counties including Lauderdale where a another rare larid, Glaucous Gull, also got counted Dec. 19 (DJS, MAB). Other Glaucous sightings included two in Jefferson during January and February (PY) and a single in Cameron Feb. 13 (JK, DR). Rare in Alabama, an imm. Great Blackbacked Gull was sighted Dec. 2 in Baldwin, AL, where an adult was seen the next day (GDJ, DGJ et al.). First ever in North America, a Gray-hooded Gull was sighted Dec. 26 in Apalachicola, FL (TL ph., DM ph) where it was seen by few. Unusual terns flocked to Louisiana: 30 Gull-billeds congregated in Lafourche, LA, Jan. 17 (PW, MMy, DPM); a single Sandwich stayed in Cameron Dec. 21 (PC); Common was represented by singles in John the Baptist Dec. 26 (JaB) and in Cameron Dec. 20 (MS, KF), and by three elsewhere in Cameron Dec. 19 (KF, BV).

ունըչեններումի ուլու, ու քրու պատոնքը երկ մին է երևույն հանգերել ու ուսել է երուլ չ

DOVES THROUGH HUMMINGBIRDS

Eurasian Collared-Doves showed up in Yell and Little River, AR, during December (KN, LN) and February (DA, JC), respectively; the species continued its dramatic expansion in Alabama (fide GDJ) where 134 Jan. 2 in Baldwin provided a Gulf Coast maximum. Small groups of up to 19 were reported in 3 Mississippi counties; 3 Louisiana parishes reported small flocks, but Jefferson boasted . . . er hosted . . . ok, tolerated 155 Feb. 17 (PY); if the increased number of collared-doves is having a negative effect on a competing columbid, Mourning Dove, in the Region, it is not yet evident (Table 1). Also increasing dramatically in Alabama, 89 White-winged Doves were counted in Baldwin Jan. 2 (PCT, BT, BCG, JoL), establishing a state maximum, and a single showed up in the Tennessee Valley in Morgan Jan. 20 (SH, PhC). Reports of White-wings emanated from 11 Louisiana parishes, an "unprecedented scatter" of sightings (SWC). An Inca Dove in Sevier, AR, Dec. 18 (CM) and eight there Feb. 22 (DA) were at a new site while other Arkansas reports came from Clark, Columbia, and Little River; the species is also becoming more common in Shreveport, LA (RJS). Continuing the theme of columbid conquest, Com. Ground-Doves, normally scarce in Louisiana, were reported from at least 6 parishes—but in mercifully small numbers.

Four Greater Roadrunners in *DeSoto*, LA, Jan. 10 (JfT, JnT) represented a notable concentration. *Jackson*, MS, hosted a Groove-billed Ani Feb. 2–20 (KH et al.); the only others were in *East Baton Rouge* and *Jefferson*, LA. Rare in Tennessee, Long-eared Owls were noted in *Henry* Dec. 5 and in *Lake* Dec. 19–Feb. 28 (JRW, m.ob.). Four Com. Nighthawks lingered in *Davidson*, TN, to at least Dec. 7 (CS), and one–four visited *Shelby*, TN, until Dec. 20 (LCC, MH); another stayed late in *Lowndes*, MS, Dec. 16–17 (JAJ, TLS).

Four Broad-billed Hummingbirds were banded in Louisiana, doubling the number previously banded (NN, DP). Buff-bellied Hummingbirds in Hancock, MS, Jan. 1-Feb. 18 and beyond (SJP et al.) and Harrison, MS, Feb. 19 (SJP, HM) were the only ones reported outside of Louisiana. Banded Jan. 18 (BC) in Harrison, MS, a 2nd-year male Ruby-throated Hummingbird present Jan. 14 to at least Feb. 28 was the only one reported away from Louisiana where numbers were unprecedentedly high (see Table 2, which does not account for all reports). A Black-chinned Hummingbird banded in Williamson, TN, Dec. 3 (CS, PM) established the 3rd state record. A Calliope Hummingbird banded by the Sargents in Davidson, TN, during winter 1997-1998 (FN 52: 210) returned for another gig in Music City, staying Nov. 10 until late February (CS, PM, EW); Louisiana experienced a "Calliope high" with about 20 present during the winter (SWC). Rufous Hummingbirds were banded and stayed the entire season in Davidson and Rutherford, TN (CS, PM) while another was banded in Wilson, TN, Dec. 4 (CS, PM); an ad. male stayed until frozen by a cold snap Dec. 23 in Washington, AR (DJ*), but another made it through the winter in Pulaski, AR, where it was banded (PF). Still others were noted in Cleburne and Faulkner, AR; Mississippi's only report came from Harrison where a male was noted Jan. 18 and Feb. 16 (SJP et al.).

FLYCATCHERS THROUGH WARBLERS

A Vermilion Flycatcher in *Hancock*, MS, provided a New Year's Day treat (SJP et al.); the only other reports came from *Ouachita* and *Rapides*, LA. The only Brown-crested

Flycatcher came from Plaquemines, LA, Jan 31 (DPM et al.). About the 4th Mississippi record, an Ash-throated Flycatcher remained in Jackson Feb. 6-28 onward (KH et al), 4 Louisiana parishes hosted others. Scissortailed Flycatchers were more numerous in Louisiana than usual. Rarely encountered in Tennessee in winter, a White-eyed Vireo was noted on the Memphis CBC Dec. 20 (†RP); two-three in Grenada, MS, Dec. 30 (FB, DeW) were also rare. Extremely rare was a Yellow-throated Vireo in Hancock, MS, Jan. 4 (SJP). Early-arriving swallows included two Purple Martins in Lowndes, MS, Feb. 11 (KK) and a Tree Swallow in Oktibbeha, MS, Feb. 26 (TLS); six of the latter were over the Gulf 135 mi s. of Morgan City, LA, Jan. 15 (DNP). A Bank Swallow in Cameron Dec. 20 (DLD, SWC) was amazin' The Rock Wren appearing in White, AR, last fall stayed until Mar. 1 (KN, LN). The only Bewick's Wrens were two in Little River, AR, Feb. 21 (JC, DA) and one in Evangeline, LA, Dec. 30 (MF).

A state maximum of 394 E. Bluebirds kept counters busy Jan. 2 in *Baldwin*, AL (*fide* GDJ). Where would you least expect six Am. Robins? You guessed it—135 m1 s of Morgan City, LA, Jan. 15 (DNP); Remsen evaluated robin numbers in Louisiana as "excellent"; I guess so. A Gray Catbird in *Putnam*, TN, Dec. 22–Jan. 6 (†RWS) provided the 2nd local winter record, and one in *Oktibbeha*, MS, Jan. 12–23 (MC, TLS) was rare. American Pipits beleaguered bird counters in *Limestone/Morgan*, AL, Dec. 19 when 1204 established a state maximum, 375 in *Lafayette*, MS, Jan. 23 (GK, SK) were considered a large group at that site.

Arkansas' first winter Tennessee Warbler showed up in Jefferson Dec. 26 (†GL, JL), another was in Cameron, LA, Dec. 12 (SJP, RDP, DPM). Three Louisiana parishes held Nashville, and two parishes Yellow warblers, both very rare in winter. Alabama's 3rd winter Cape May Warbler-and first in the Tennessee Valley-remained in Lauderdale Jan. 10-Feb. 28 onward (SW, m.ob.). Blackthroated Green Warblers were noted in Cameron and Plaquemines, LA. A Prairie Warbler in Putnam Dec. 16 (†SJS, †GKE, †JCS) provided Tennessee's first winter record (see Appalachian Region for particulars about the 2nd winter record). Palm Warblers wintered inland to the far northern limits of the Region in noteworthy numbers. American Redstarts appeared in 3 Louisiana parishes. Another Arkansas winter first occurred when an Ovenbird tarried in Prairie Dec. 22-Feb. 28+ (TS, RAS, ph DaC, OA), and Louisiana hosted Ovenbirds

Table 1. Christmas Bird Count Numbers for Accipiters and Two Columbids from 68 Central Southern Region CBCs Conducted All Ten Years 1988–1997										
Species	1988	1989	1990	1991	1992	1993	1994	1995	1996	1997
Sharp-shinned Hawk	194	157	176	232	216	222	230	245	229	253
Cooper's Hawk	98	70	85	128	133	134	149	169	152	175
Eurasian Collared-Dove	0	0	0	3	71	238	697	633	840	1562
Mourning Dove	15,925	17,172	22,407	19,672	22,049	20,300	26,217	22,741	21,544	19,016

and N. Waterthrushes in 4 parishes each. A Wilson's Warbler in *Limestone* Feb. 6 (DRC, RC) provided the first winter record in Alabama's Tennessee Valley. Yellow-breasted Chats showed up in 4 Louisiana parishes.

TANAGERS THROUGH FINCHES

Summer Tanagers were present in East Baton Rouge and Caddo, LA, the latter farther north than most sightings occur. The male W. Tanager in St. John the Baptist, LA, ın late February (BJ, RS, MWe, m.ob.) was a show stopper. The 2nd Alabama inland winter record for Lark Sparrow took place Dec. 19 in Lauderdale (JTG); others were in Cameron and LaSalle, LA. Eight Henslow's Sparrows in Bradley, Drew, and Hempstead Feb. 13–15 (WMS, JLY) continued a recent trend toward greater wintering numbers in Arkansas; another in Lee Dec. 19 (WDR, GEH) provided Alabama's first winter record in the Mountain Region; the banding project in Jackson, MS, once again scored big with 146 banded Jan. 10-Feb. 24 (MSW, SI, MD). Rarely reported in middle Tennessee and Arkansas, single Harris's Sparrows were noted in Lawrence Dec. 29-Jan. 20 (MoW) and Little River Feb. 21 (DA, JC), respectively. Found fittingly enough on Snow Bunting Pen., Davidson, TN, a Snow Bunting made one of the species' rare appearances in that state Jan. 10 (CS). Up to 500 Lapland Longspurs were noted throughout the period in Lake, TN (JRW, MG et al.) while 200+ were in Henry, TN, Jan. 9 (JRW); 386 in Limestone, AL, Dec. 27 (SWM) provided a good number at that location; Louisiana had records in 5 parishes Smith's Longspurs were monitored at the Stuttgart Airport in Prairie, AR, Jan. 16-17 (JZ, LZ, CB et al.) with up to 52 reported; in Sebastian, AR, 20 were noted Jan. 17 (TB, MP, DCh, MM) while 75 were present there Feb. 14 (MM, DMe).

Seldom well documented in winter, a Rose-breasted Grosbeak was in *East Baton Rouge* Jan. 1–16 (CoB, JBa). A basic-plumaged ad. male Indigo Bunting in *Baldwin*, AL, Dec. 14 (GDJ, DGJ) was late/rare; Louisiana had sightings in 6 parishes. Fifth In winter for Alabama, a Painted Bunting stayed in *Baldwin* Dec. 17–Jan. 7+ (ALM); Louisiana, again, had sightings in 6 parishes. First locally in winter, a female Baltimore Oriole wintered in Putnam, TN, Dec. 19-Feb. 28+ (†GKE, JCS, ES, JS); a male stayed in Davidson, TN, Dec. 18-Feb. 28+ (CS); another of unreported sex appeared on the Memphis CBC Dec. 20 (JZ, LZ). Farther south an ad. male Baltimore was noted Jan. 9 in Shelby, AL (PH), and a female was seen eating seed from a feeder in Harrison, MS, Jan. 6 (SJP). Louisiana hosted Baltimores in 6 parishes and Bullock's in two, but the oriole of the season was a Hooded Oriole in East Baton Rouge Feb. 6-28 onward (AD et al.). Winter finches were generally present in extremely low numbers or absent.

Acknowledgment: My thanks to Ginger K. Ensor for compiling the data in Table 1.

Contributors (subregional editors in boldface): Orapin Anurakdanthai, David Arbour, Mahlon Ayme, Coney Barre (CoB), Janie Barre (JBa), James Beck (JaB), Michael A. Beuerlein, L. C. Binford, Jerry Bird, Paul Blakeburn, Roger Breedlove, William J. Bremser, Greta Bremser, Fred Broerman, Tom Brooks, Carolyn Bullock, Joe Cambre, Steven W. Cardiff (Louisiana, scardif@unix1.sncc.lsu.edu), Bennett Carver, Phillip Casteel (PhC), David Chapman (DCh), Darrell Chatelain, Darren Clark (DCl), Susan Clark, Lula C. Coffey, David Cooper (DaC), P. Conover, Margaret Copeland, W. Glen Criswell, Dean R. Cutten, Raelene Cutten, Michelle Davis, W. Marvin Davis, Harry Dean, Antoinette DeBosier, Donna L. Dittman, Robert H. Doster, Robert A. Duncan (nw Florida), Ginger K. Ensor, Karen Fay, Marty Floyd, Perk Floyd, Ben C. Garmon, Jeff T. Garner, Mark Greene, Ken Hackman, David Haggard, Paul B. Hamel, Robert M. Hatcher, Martha Heinemann, Jim Hight, Geoff E. Hill, Susan Hollyday, James F. Holmes, Patricia Honeycutt, Siri Ibarguen, Debra G. Jackson, Greg D. Jackson (Alabama, gregdebi.jackson@prodigy.net), Jerome A. Jackson, Bobby Jacob, Douglas James, Keith Kimmerle, Cecil Kersting, Paul A. Kittle, Joe Kleiman, Joanne Kleiman (JoK), Gene Knight, Shannon Knight, Vicki LaFevers, Jeffrey Lee (JeL), Thomas Lewis, Jon Lloyd

Table 2. Hummingbird Totals on Louisiana Christmas Bird Counts (CBC)* and Totals Banded by Nancy Newfield (NN) and Dave Patton (DP) in Louisiana during Winter 1998–1999 (overlap exists between the two sets of data)								
Species	LA CBC Totals	NN/DP Banding Totals						
Broad-billed	1	4						
Buff-bellied	10	10						
Ruby-throated	20	25						
Black-chinned	26	31						
Archilochus sp.	3	-						
Calliope	5	4						
Broad-tailed	2	2						
Rufous	43	62						
Allen's	2	4						
Rufous/Allen's	53							
hummer sp.	4							
Total	169	142						
*Data supplied by	Steven W. Cardiff.							

(JoL), Guy Luneau, Joan Luneau, Charles Lyon, Steve W. McConnell, Portia McMillan, Douglas McNair, Helen Mead, David Mehlman (DMe), Gail Menk, Ann L Miller, Charles Mills, Mike Mlodinow, Nancy Moore, David P. Muth, Mac Myers (MMy), Nancy Newfield, Kenny Nichols, LaDonna Nichols, Glenn Ousset, Helen and Max Parker (Arkansas, hlparker@ CSWNET.com), Dave Patton, James Peavy (JPe), Lynda Perry, Stacy J. Peterson, John Porter (JoP), Jackie Porter, Mike Powers, Richard Preston, Robert D. Purrington, Robert R. Reid, David N. Purvis, Linda B Reynolds, Richard A. Reynolds, Dave Roark, W. Douglas Robinson, Marion Schiefer, Terence L. Schiefer (Mississippi, tschiefer@entomology.msstate.edu), Jeanne C. Schmitzer, Rosemary J. Seidler, William M. Shepherd, Damien J. Simbeck, Richard W. Simmers, Ruth Ann Singleterry, Terry Singleterry, Christopher Sloan, Curt Sorrels (CSo), Barbara H. Stedman, Stephen J Stedman, Ron Stein, Evelyne Stites, John Stites, Alison Styring, Mark Swan, Betsy Tetlow, Phil C. Tetlow, Jeff Trahan (JfT), Jean Trahan (JnT), Bill Vermilion, Winston A. Walden, Martha G. Waldron (w Tennessee, waldronm@ten-nash.ten.k12.tn us), Phillip Wallace, Donald Ware, Shirley Wayland, Melvin Weber (MWe), Ted Weems, Dennis Welch (DeW), Melinda Welton, Jim Whelan, Morris Williams (MoW), Jeff R. Wilson, Chris Witt (CW1), Mark S. Woodrey, Chris Woodson, Evelyn Wright, Peter Yaukey, J. Lyndal York, John Zempel, Linda Zempel.

Stephen J. Stedman, Department of English, Box 5053, Tennessee Technological University, Cookeville, TN 38505 (sstedman@ tntech.edu)