R ainfall was well above average in June, and temperatures were well above average in July, with numerous record temperatures set. As seems to always be the case in Florida in summer, either few observers were out in the field, or few people sent in reports, leaving the state with spotty coverage for the season.

Abbreviations/definitions: report (any observation); record (only those reports verifiable from photograph, videotape, or specimen evidence); N.S.R.A. (North Shore Restoration Area, *Orange*).

LOONS THROUGH SPOONBILLS

As is normal, a few single Common Loons were present: at St. Marks N.W.R., Wakulla 1 Jun (AB, DS), L. Ella, Leon 3-28 Jun (DS, RL, GM), and St. Teresa, Franklin 20 Jun (JM). Up to 10 Pied-billed Grebes were heard at Ten Thousand Islands N.W.R., Collier 7-13 Jun (TDo). Strong e. winds produced numerous observations of pelagic species along the Atlantic coast, with Bankert reporting 8 Cory's, 40+ Greater, and 2 Audubon's Shearwaters at Melbourne Beach, Brevard 23 Jun; Hope reporting a Greater and 15 Cory's Shearwaters at Boynton Inlet, Palm Beach 24 Jun; and Berney reporting 60 Cory's Shearwaters and 140+ unidentified large shearwaters off John Lloyd S.P., Broward 24 Jun. A significant die-off of shearwaters occurred along the Atlantic Coast in late Jun. In Volusia, one Sooty, 3 Cory's, and 291 Greaters were found dead or dying on beaches 21-27 Jun (MBr), 6 Greater Shearwaters were found dead at Canaveral N.S., Brevard 26 Jun (AV), and single dead Greaters were found at Amelia I., Nassau 24 Jun (PL) and Vilano Beach, St. Johns 26 Jun (DR). Two pelagic trips conducted 25 Jun, one from Miami (RT, RU) and one from Marathon, Monroe (LaM, BM, JBo et al.), found 17 and 50+ Cory's, one and 150+ Greater, and 33 and one Audubon's Shearwaters, respectively. Additionally, 21 Wilson's Storm-Petrels were seen on the Miami pelagic (RT, RU), and single Wilson's and Band-rumped Storm-Petrels were seen on the Marathon pelagic (LaM, BM, JBo et al.). Other storm-petrels included 2 Wilson's and one Band-rumped 28-32 km off Hillsborough Inlet, Broward 19 Jun (MBe). A single Leach's Storm-Petrel was reported from Melbourne Beach, Brevard 23 Jun (AB). A fishing trip 170 km off Ponce Inlet, Volusia found 40+ Cory's, 20+ Audubon's, and 2 Greater Shearwaters and 4 Wilson's Storm-Petrels 29 Jul (BW). Single Masked Boobies were at Boynton Inlet 11 & 24 Jun (BHo), while 2 Brown Boobies were off Marathon 25 Jun (LaM, BM, JBo et al.). About 30 Northern Gannets were at St. Marks N.W.R. 1 Jun (DS), with singles at Boynton Inlet 12 Jun (BHo), St. Teresa 19 Jun (JDo), Alligator Pt., Franklin 11 Jul (JDo, JM), and the Pasco

Florida



coast 20 Jul (KT).

The Polk phosphate mines usually have American White Pelicans during the summer, but 450 on 23 Jul was an impressive count (PT, CG). The peak of 663 Magnificent Frigatebirds at the ABC Is., Collier was well off last year's high count of 1629 (TDo). Hurricane Dennis swept numerous Magnificent Frigatebirds into the interior of the peninsula 10 Jul, with 5 at Newnans L., Alachua (ReR, [Br et al.), 2 at Ocala, Marion (AL), 15 at Brooksville, Hernando (KW), and singles at Micanopy, Alachua (C&ES) and Lorida, Highlands (PG). This storm also brought an impressive 50+ frigatebirds to Alligator Pt. 11 Jul (JDo). Two calling Least Bitterns were heard 13 Jun at Ten Thousand Islands N.W.R. (TDo), where singles had been heard in the past two years. A maximum of 28 Least Bitterns was at Lake Apopka N.S.R.A. 3 Jul (HR). A single Reddish Egret was inland to Gainesville, Alachua 3 Jun-11 Jul (JBr, BC), with coastal strays n. to Alligator Pt. 15 Jun (JDo) and St. Marks N.W.R. 8 Jul (TC). An ad. White-faced Ibis was at Springhill Rd. W.T.P., Leon 5 Jun (TC). Inland Roseate Spoonbills were at Homestead, Miami-Dade 12 Jul (RD) and the Polk mines 19 Jun and 10 Jul (PT, CG).

David J. Powell

10800 Brighton Bay Boulevard NE

Apt. 15207

St. Petersburg, Florida 33716

(vireo@vireos.com)

WATERFOWL THROUGH SKIMMER

Black-bellied Whistling-Ducks continue to thrive in the center of the state, with 10 at Lake Apopka N.S.R.A. 5 Jun (HR), 15 in Okeechobee 5 Jun (MK, GQ, RB), 20, including 5 young, at Orlando Wetlands Park, Orange 12 Jun (BA, CP), and 48 at Polk mines 19 Jun (CG, PT). Fulvous Whistling-Ducks are not doing as well, with the only reports from Robinson at Lake Apopka N.S.R.A., with a maximum of 8 on 28 Jul. A Bluewinged Teal was at St. Marks N.W.R. 2-5 Jun (AB, DS), with 2 others at Lake Apopka N.S.R.A. 26 Jun (HR). A male Ring-necked Duck was a surprise at Lake Apopka N.S.R.A. 24 Jun (BA, AV). A lone Redhead was found at St. Marks N.W.R. 1-3 Jun (DS, AB). A female Red-breasted Merganser was at Gullivan Key, Ten Thousand Islands N.W.R. 14 Jun (TDo). A male Masked Duck at Sarasota, *Sarasota* 6–7 Jun (TE et al.) represented a very rare central peninsula report.

Ospreys are common in Florida, but 102 at Lake Apopka N.S.R.A. 3 Jul (HR) made an especially high count. Hundreds of Swallowtailed Kites fed on abundant dragonflies at Lake Apopka N.S.R.A. in Jul, with a peak of 856 on 17 Jul (HR). At the Fisheating Creek, Glades roost, 1954 were counted 30 Jul (BP, PG et al.). White-tailed Kite is a scarce species in Florida, so two reports is about normal. Snail Kites nested at Inverness Pool, Citrus (FA), the northernmost nesting in many decades. Significant reports of Mississippi Kites were one at Merritt Island N.W.R., Brevard 1-4 Jun (TDu et al.), 5 near Green Cove Springs, Clay 15 Jun-17 Jul (LeM), and a pair at s. Jacksonville, Duval 7-17 Jul (JCo). Five Mississippis at Belleview, Marion 20 Jun (SB) were more expected. Cooper's Hawks continue to spread southward in the state, with many reported from the Tampa Bay area and others reported from Miami-Dade and Collier. Six reports of Short-tailed Hawk were received, about normal for this scarce species. The only American Kestrel report was one at Lake Apopka N.S.R.A. 26 Jun (HR).

A Black Rail 10 Jul at Polk mines (CG, PT) was a rare inland report. Robinson had an impressive 175 King Rails at Lake Apopka N.S.R.A. 31 Jul. An American Coot photographed feeding young 18 Jun established the first confirmed breeding for Pasco (KT). With the number of oversummering shorebirds in the state, the first southbound birds are often difficult to detect. A Black-bellied Plover at Little Estero Lagoon, Lee 21 Jun (CE) was likely the first "fall" migrant. A Wilson's and 3 Semipalmated Plovers at Crandon Beach, Miami-Dade 30 Jun (RD) were also thought to be fall arrivals. Ewell had 3 ad. and 23 young American Oystercatchers at Little Estero Lagoon 5 Jul, including 2 birds banded in North Carolina and one banded in Virginia. This is the 3rd consecutive year that banded oystercatchers have turned up at this location. An apparently oversummering American Avocet was at Snake Bight, Everglades N.P., Monroe 13 Jun (AB, EE), while 32 others were at Polk mines 23 Jul (PT, CG). The first reported Greater Yellowlegs for the fall was at Springhill Rd. W.T.P. 5 Jul (GM), while the first Lesser Yellowlegs was in mid-Pinellas 9 Jul (JF). Another migrant Lesser was 111 km off Ponce Inlet 29 Jul (BW). The only Solitary Sandpiper reported was one at Lake Apopka N.S.R.A. 23 Jul (HR). The first southbound Spotted Sandpipers were 6 at Springhill Rd. W.T.P. 21 Jul (GM). An apparent late northward migrant Whimbrel was found at Playalinda Beach, Brevard 3 Jun (AV). At Little Estero Lagoon, 7 Marbled Godwits 14 Jul SA Hurricane Dennis moved through the Straits of Florida and into the Gulf of Mexico in mid-Jul, bringing 2 Sooty Terns and the above-mentioned frigatebirds inland to Newnans L., Alachua 10 Jul (SC, GP). Another Sooty was inland at Tallahassee, Leon 11 Jul (MS). Dozier and Murphy had one Bridled and 11 Sooty terns at Alligator Pt. 10 Jul, and 24 Sooty and 3 Bridled Terns there 12 Jul. Another Sooty Tern was at Honeymoon Island S.P., Pinellas 10 Jul (EK).

Less obvious, but also apparently related to *Dennis*, was a significant influx of pelagic terns to the s. Atlantic coast 9 Jul. Hope, Berney, and others observed 120+ Bridled Terns, 100+ Sooty Terns, and 2 Brown Noddies at Boynton Inlet, while 2 Sooty Terns were off Crandon Beach (RD) and 6 Brown Noddies at Boot Key, *Monroe* (AW). In addition to the terns, one or 2 White-tailed Tropicbirds were at Boynton Inlet 9 Jul (BHo, MBe et al.). The effects of the storm were short-lived, as Diaz found only 2 Bridled and 5–6 Sooty Terns at Boynton Inlet 10 Jul, along with one Brown Noddy at Lloyd S.P. The last report from this event was a Bridled Tern at Jupiter Inlet Colony, *Palm Beach* 11 Jul (J&LH).

and an alternate-plumaged Red Knot among 150 oversummering basic-plumaged knots 5 Jul were thought to be fall migrants (CE). Leary observed 120 Red Knots, including one banded in Argentina, at Bird I., Nassau Sound, Duval 24 Jul, and 700 there, including 2 banded in Chile, by the end of the month. The first southbound Semipalmated Sandpipers were at Springhill Rd. W.T.P. 11 Jul (GM), while the first fall-arrival Western Sandpipers were 3 in alternate plumage at Little Estero Lagoon 5 Jul (CE). Ewell had his highest count of oversummering peeps at Little Estero Lagoon, with 160 Western/Semipalmated Sandpipers in basic plumage 5 Jul. An impressive 8100 Least Sandpipers were at Polk mines 23 Jul (CG, PT). White-rumped Sandpipers continue to move northward into early Jun, with 5 at Fort De Soto Park, Pinellas 1 Jun (LA) and 3 there 2 Jun (JG). Berney found 80+ Pectoral Sandpipers near Belle Glade, Palm Beach 31 Jul. Lingering Dunlins were seen at Ft. De Soto Park 1 Jun (AB, DS) and St. Marks N.W.R. 4-5 Jun (DS, AB). Two Stilt Sandpipers at Polk mines 23 Jul (PT, CG) were the only report. Alternate-plumaged Short-billed Dowitchers at Little Estero Lagoon 6 Jul (CE) and Crandon Beach 8 Jul (RD) were thought to be migrants. Three early Red Phalaropes were about 128 km off Ponce Inlet 29 Jul (BW).

Clark had 8000 Laughing Gulls, including many fledged young, and 2-6 Lesser Blackbacked Gulls at Huguenot Memorial Park, Duval in Jul. Two imm. Lesser Black-backed and 3 imm. Great Black-backed Gulls were at Crandon Beach 4 Jul (RD). Up to 4 Gullbilled Terns were at Polk mines 19 Jun-23 Jul (PT, CG), with 2 others at John Lloyd S.P. 11 Jun (MBe). Clark had 1100 Royal Terns at Huguenot Memorial Park during the end of Jul. Single Roseate Terns were at Lloyd S.P. 11 Jun (MBe) and Boynton Inlet 12 Jun (BHo). A surprising 55 Common Terns were inland at Polk mines 10 Jul (CG, PT). A single Arctic Tern was at Boynton Inlet 11 Jun (BHo). For a species that does not breed in Florida, 920 Forster's Terns at Polk mines 23 Jul (CG, PT) was an impressive mid-summer count. Up to 225 Least Terns were at Key Biscayne, Miami-Dade 15-21 Jul, an unusually high

number for this location (RD); at *Polk* mines, Leasts numbered 274 on 19 Jun (PT, CG). A fishing trip about 20 km off Hillsborough Inlet, *Broward* 19 Jun found 2 Bridled Terns (MBe). A Sooty Tern was picked up alive on a *Volusia* beach 25 Jun (MBr). A fishing trip 170 km off Ponce Inlet found 300+ Sooty Terns, 15+ Bridled Terns, and 4 Brown Noddies 29 Jul (BW).

The first southbound Black Tern was at Bald Pt., Franklin 4 Jul (JM). Large numbers of Black Skimmers were inland to Polk mines, with 412 on 19 Jun, 690 on 10 Jul, and 650 skimmers 23 Jul (CG, PT). Three hundred Black Skimmers were at Bird Is., Nassau Sound 24 Jul, where their colony was washed out by high spring tides (PL).

PIGEONS THROUGH FINCHES

White-crowned Pigeons are primarily birds of the Keys; 68 at Southern Glades W.M.A., Miami-Dade 12 Jul (RD) was an unusually high number on the mainland. White-winged Doves are becoming more common in the Tampa Bay area, with 29 at Brandon, Hillsborough 20 Jul (DP) the highest number reported. Burrowing Owls were present throughout the season at Kissimmee Prairie Preserve S.P., Okeechobee, where 19 was the highest count 13 Jul (PM). Very local nesters, Common Nighthawks nested unsuccessfully at Fort Clinch S.P., Nassau in Jun (PL). The only report of Antillean Nighthawks was of 3 at Cudjoe Key, Monroe 24 Jun (AB, MH). At least 2 Whip-poor-wills were heard and seen at Bald Pt. 5-16 Jun (JM, JDo); there are no breeding reports for Florida. Ruby-throated Hummingbirds are scarce in Florida away from the extreme north, so four reports from the mid-pen. were about normal. A Belted Kingfisher at St. Marks N.W.R. 8 Jun (JDo) may have been a summering bird. The first apparently returning Belted Kingfisher was one at Ocklawaha, Marion 30 Jun (AL). Single Western Kingbirds were in mid-Pinellas 2 Jun (JF) and Bald Pt. 5 Jun (JDo). Two Yellow-throated Vireos heard singing at Mead Garden, Winter Park, Orange 10 Jun+ (BA et al.) were at the s. edge of their range. A Redeyed Vireo at Sarasota 2 Jun (JDu, EM) was likely a late migrant; the first southbound Red-eyed was at Coconut Pt., Brevard 26 Jul (AB). A Black-whiskered Vireo was n. of normal in mid-Pinellas 6 Jun (JF). This species used to occur regularly in Pinellas but is now rare there.

A "few" Purple Martins at Tavernier, Monroe 24 Jun (BM) were thought to be southbound migrants. Noteworthy swallows at Lake Apopka N.S.R.A. were single Tree Swallows 6, 11, & 26 Jul, one or 2 Bank Swallows on seven dates during the season, and up to 3 Cliff Swallows 23-31 Jul (HR). Barn Swallows are continuing to expand as breeders in the state, so individuals at two locations in Everglades N.P. 4 & 5 Jun (BR) were intriguing. More expected, but still impressive, was the presence of at least 55 Barn Swallows on each of 14 dates at Lake Apopka N.S.R.A., with a maximum of 260 on 23 Jul (HR). A House Wren at Clearwater, Pinellas 26 Jun (EH) was only the 3rd reported in summer in the state. Single American Robins at Cedar Key, Levy 2 Jun and 6-11 Jul (DH) were interesting. An out-of-season Gray Catbird was at Lake Apopka N.S.R.A. 5-12 Jun (HR). Robinson observed the first Common Myna at Lake Apopka N.S.R.A. 11 Jul, further evidence of the expansion of this species in Florida.

A well-described Blue-winged Warbler at Ocklawaha Prairie R.A., Marion 21 Jul (AL) furnished only the 2nd summer report for Florida. The first southbound Yellow Warbler was at Cedar Key 25 Jul (DH). A Cerulean Warbler was at a water drip in s. Jacksonville 30 Jul (ICo). The first southbound Black-andwhite Warbler was at the Seminole Environmental Center 9 Jul (AV, LoM). An American Redstart at Gainesville 9 Jun (GP) was likely a late northbound migrant, while one at Kissimmee Prairie Preserve S.P. 16 Jul (PM) was the first southbound migrant. A Swainson's Warbler was heard in Tallahassee 26 Jun (GM). The first southbound Louisiana Waterthrushes were singles at Tallahassee 5 Jul (FR), Seminole Environmental Center 9 Jul (LaM, AV), and Newnans L., Alachua 10 Jul (JH, LD). The first southbound Hooded Warbler was in mid-Pinellas 25 Jul (JF). Blue Grosbeaks and Indigo and Painted Buntings were present throughout the season at Lake Apopka N.S.R.A., with maximum daily counts of 25, 27, and 3, respectively (HR). A male Varied Bunting that hit a window at Siesta Key, Sarasota 1 Jun (MvD; *Archbold Biological Station) provided the first record of the species for Florida. Small numbers of Dickcissels have been present at Lake Apopka N.S.R.A. in recent summers, but this summer's number of birds is unprecedented for Florida. Robinson found Dickcissels on all 14 of his surveys this year, with a maximum of 89 on 5 Jun and counts of 40+ until 20 Jul and 4 still present at the end of the season. Breeding was confirmed when a nest with eggs was found 19 May.

Out-of-season Bobolinks were at Lake Apopka N.S.R.A. 5 Jun (HR) and Cedar Key 18 Jun (RoR). Blackbird numbers always increase at Lake Apopka N.S.R.A. in Jul, but this year's numbers were nothing short of amazing, with counts of over 100,000 each for Red-winged Blackbirds, Common Grackles, and Boat-tailed Grackles, the maxima being 303,000 Red-winged Blackbirds and 117,000 Common Grackles 17 Jul and 167,000 Boat-tailed Grackles 14 Jul (HR)! Numbers at the roost rapidly decrease by mid-Aug-where do these birds come from and where do they go? Shiny Cowbirds were reported in good numbers, with singles at Alligator Pt. 1-7 Jun (JDo), Bald Pt. 17 Jun (JDo), and Cedar Key 2 Jul (DH), duos at Fort De Soto 2 Jun (LA) and Flamingo, Everglades N.P. 5 Jun (BR), up to 7 at Redlands, Miami-Dade through 3 Jul (RT, MF, RD), and up to 13 at Lake Apopka N.S.R.A. 19 Jun-28 Jul (HR). Up to 6 Bronzed Cowbirds, not yet confirmed to breed in Florida, were at Redlands, Miami-Dade through 12 Jul (RT, MF, RD). Orchard Orioles were present at Lake Apopka N.S.R.A. all season, with a maximum of 24 on 1 Jul (HR). An American Goldfinch lingered at Gainesville until 1 Jun (LD). House Finches were present in good numbers in the cen. peninsula, with one in mid-Pinellas in Jul (JF), up to 6 at Altamonte Springs, Seminole all season (PH), one at Lake Apopka N.S.R.A. 4 & 14 Jul (BA, HR), and up to 6, including a female feeding a fledgling, at Mead Garden (AV, BA, JCl).

Cited contributors (area compilers in bold): Bruce Anderson, Lyn Atherton, Frank Aumack, Andy Bankert, Mark Berney (MBe), Sam Benedetto, Robin Bjork, John Boyd (JBo), Michael Brothers (MBr), Judy Bryan (JBr), Bob Carroll, Joelyn Clifton (JCl), Julie Cocke (JCo), Steve Collins, Tom Curtis, Lloyd Davis, Robin Diaz, Terry Doyle (TDo), Jack Dozier (IDo), Jeanne Dubi (IDu), Thomas Dunkerton (TDu), Tommy Elliot, Erik Enbody, Charlie Ewell, Mark Faherty, Judy Fisher, Chuck Geanangel, Jill Gaetzi, Paul Gray, Jack & Liz Hailman, Mitchell Hait, Erik Haney, Bev Hansen (BHe), Dale Henderson, John Hintermister, Brian Hope (BHo), Paul Hueber, Marianne Korosy, Ed Kwater, Patrick Leary, Rob Lengacher, Angela Luzader, Lorne Malo (LoM), Larry Manfredi (LaM), Lenore McCullagh (LeM), Gail Menk, Edith Miller, Paul Miller, Brennan Mulrooney, John Murphy, Geoff Parks, Cheri Pierce, David Powell, Peggy Powell, Bill Pranty, Gallus Quigley, Bryant Roberts, Harry Robinson, Diane Reed, Ron Robinson (RoR), Rex Rowan (ReR), Fran Rutkovsky, David Simpson, Matt Smith, Catherine & Eugene Stoccardo, Pete Timmer, Roberto Torres, Ken Tracey, Raul Urgelles, Michelle van Deventer, Alex Vinokur, Bob Wallace, Kristen Wood.

