

Bruce H. Anderson

he season was rather mild, with barely freezing temperatures reaching south to Lake Okeechobee twice, causing topical damage to vegetation in outlying areas from Orlando southward. The drought in the peninsula continued, with a cumulative halfmeter precipitation deficit in some portions of the central and southern regions since 2005. The kinglet invasion that began in the fall continued, and countless flocks of American Robins and Cedar Waxwings reached well into southern Florida. State rarities included Thayer's Gull, Dovekie, Hammond's Flycatcher, Horned Lark, Mountain Bluebird, and Western Meadowlark. Given that up to 12% of Florida's 30,000+ Mottled Ducks are believed to be hybrids with Mallard (<http://www.myfwc.com/duck/Mottled/thep roblem.htm>), the Florida Fish and Wildlife Conservation Commission announced that it will step up its efforts to control Florida's feral Mallard population. In addition, the Commission will be initiating a study of Smoothbilled Ani, an endangered species in its limited United States range but one that has heretofore received little attention.

Abbreviations/definitions: F.O.S.R.C. (Florida Ornithological Society Records Committee); report (any observation); record (only those reports verifiable from photograph, videotape, or specimen evidence); UF (Florida State Museum).

WATERFOWL THROUGH CARACARA

Breeding throughout much of the peninsula, 458 Black-bellied Whistling-Ducks were n. at Gainesville, *Alachua* (C.B.C. data). In contrast, the greatest concentration of Fulvous Whistling-Ducks, a species that outnumbered

Black-bellieds a decade ago, was about 150 in Hendry (M. England). Geese were rare throughout. The only Greater Whitefronted Goose was at St. Marks N.W.R., Wakulla 16 Dec (C.B.C.). Snow Geese were in Leon (one; Glenda Simmons), Hernando (up to 4; A&BH et al.), Volusia (6; C.B.C.), and s. Brevard (one; C.B.C.). Rarely reported inland or Gulfside, a Brant at Gainesville 29 Dec (AK), was likely the same imm. as seen at at Ft. Island Gulf Beach, Citrus 2-22 Jan (m.ob.). Mallard × Mottled Duck hybrids were in Hamilton (C.B.C.), Orange (BHA, RD), Brevard (DF), Pinellas (C.B.C.), Martin

(C.B.C.), Lee (CE), Collier (DSu), Palm Beach (Robin Statfeld), and Pasco/Hernando (200; C.B.C.). Cinnamon Teal were in Polk (one; RM, DR), and in Brevard (2), as was one Bluewinged Teal × Cinnamon Teal hybrid (m.ob.). For the 3rd consecutive year, a Eurasian Teal was found in Florida, with one at St. Marks 6 Jan (JoH et al.). The first Greater Scaup to be verified in the s. peninsula was at Eco Pond, E.N.P., Monroe 30 Dec-13 Jan (ph. LM,

m.ob.). Up to 2 Common Eiders were in St. Johns 16 Dec and 3-4 Feb (m.ob.), with singles in Nassau 9 Dec (RoS) and Volusia 23 Dec (MBr). The three scoter species appeared on both coasts; the rarest, White-winged, was in Franklin (one; AB, JM), Volusia (up to one; MBr), and Brevard (one; C.B.C.). Surf Scoters were s. to Levy (2; C.B.C.) and Brevard (up to 2; Mitchell Harris, DF), whereas Black Scoters were s. to Brevard (fide DF) and Pasco (one; BAh, AB et al.). Longtailed Ducks included singles in Wakulla (AB, AW) and Brevard (fide DF), with 3 in St. Johns (ph. BW, BR). Rare were 3 Buffleheads inland at Lake

Apopka North Shore Restoration Area, Orange 19 Dec (BHA, DR) and 12 far s. in Miami-Dade 8 Jan (RT). A Common Goldeneye was again s. at Tangerine, Orange; this year brought an imm. male (BHA, DR). The Masked Duck discovered in the fall at Viera Wetlands, Brevard continued through 29 Dec and was joined by another 12-19 Dec (DF).

Red-throated Loons were in Gulf (one; C.B.C.), Franklin (up to 8; m.ob.), Citrus (one; MG), St. Johns (2; C.B.C.), Volusia (at

least one; MBr), and Brevard (at least one, BHA et al., ph. CG). The only Pacific Loon was one in Brevard 16-20 Dec (CG et al.). A Red-necked Grebe at Destin, Okaloosa 4 Feb (Charles Saleeby) was the 8th for the w. panhandle (BD). One Eared Grebe overwintered in Palm Beach (m.ob.), while 2 in Okaloosa since fall remained until 18 Dec (BD et al.) Other Eareds included singles in Manatee 10 Dec-24 Dec (IP, ID), s. Brevard 30 Dec (C.B.C.), and Volusia 25-27 Feb (MBr, MW) The only boobies were at Dry Tortugas N.P. Monroe: 10 Masked, 15 Brown, and one Redfooted Booby 3 Jan (Sonny Bass, Stuart Pimm). Inland, Brown Pelicans were in Orange (at least one; HR, Andrew Boyle, CP) and Polk (up to 3; TP, PT). Surprising were Magnificent Frigatebirds in numbers far n. in Indian River 6 Jan (26; Nancy Soucy) and Martin 30 Dec (38; C.B.C.); farther n. was one in Taylor/Dixie 5 Jan (one; RoS). Two Whitefaced Ibis were at St. Marks 11 & 15 Jan (AW, AB et al.), and one was at Lake Apopka 24 Jan (HR). Since Hurricane Wilma devastated Everglades N.P. in Oct 2005, the Greater Flamingo flock has been absent, with only an occasional individual being reported, as was the case in late Dec (C.B.C.).

The earliest reports of migrating Swallow-



The Masked Duck to the right, present from fall through 29 December 2006 at Viera Wetlands, Brevard County, Florida, was joined by the second Masked Duck 12 (here) through 19 December 2006. *Photograph by J. William Marr.*

tailed Kites and Short-tailed Hawks came from Flagler 8 Feb (Lenore McCullagh) and Pasco 1 Feb (KT), respectively. Two Short-taileds carried nesting material in Hillsborough 11 Feb (Mauri Peterson). Rare, White-tailed Kites were reported alone in Brevard (fide TDu, DF), Osceola (BHA, RM), Hardee (Barbara Singer et al.), and Miami-Dade (AB, LM), with 2 each in Okeechobee (PMi) and Palm Beach (BH, MBe et al.). A Mississippi Kite at Lake Woodruff N.W.R., Volusia 27 Jan




For the second consecutive year, a Semipalmated Sandpiper was verified in Florida, where a few likely winter annually. Photographed at Merritt Island National Wildlife Refuge, Brevard County on 26 January 2007, this was a first for the central region. *Photograph by Kevin Karlson*.

(MW, Diane Reed et al.) was early. Broadwinged Hawks wintered n. to Brevard (at least 2; DF, C.B.C.) and Pinellas (2; C.B.C.). A Swainson's Hawk was early in Hillsborough 25 Jan (Bob Landry). Single Golden Eagles were in s. Brevard (LoM), Orange 10 Jan and 4 Feb (HR), and Baker 10 Feb (L&RoS). At new sites near the n. extreme of their range, Crested Caracaras were in Orange, at Lake Apopka (HR), at Hal Scott Preserve (BAh), and in Seminole, where at Chuluota, a pair was nest building on 1 Jan (LoM, CP). A caracara at Florida Panther N.W.R., Collier 19 Dec (DSu) was s. of the species' usual range.

RAILS THROUGH DOVEKIE

Single Yellow Rails were at Hole-in-the-Donut, Everglades N.P., Miami-Dade 2 Jan (Mark Davis) and at Kissimmee Prairie Preserve S.P., Okeechobee 20 Feb (DS). Lone American Golden-Plovers visited Lake Apopka 17, 27, & 29 Dec and 28 Feb (HR) and near Brooksville, Hernando 16 Dec-8 Jan (Alex Aycrigg, MG et al.). Rare on the e. coast, a Snowy Plover was found in Duval 17 Dec (K. Dailey), for the 5th year. Up to 44 and 42 Piping Plovers wintered in Duval (C.B.C.) and Miami-Dade (RD), respectively. Black-necked Stilts wintered n. to Seminole and Orange (C.B.C.s); 10 stilts on C.B.C.s in Wakulla and St. Johns 16 Dec were late. Single Solitary Sandpipers were in Pasco (KT) and West Palm 30 Dec (C.B.C.) and in Martin 31 Dec (C.B.C.). A Willet inland at Viera 28 Jan (fide DF) was rare, as was a Long-billed Curlew at L. Kissimmee, Osceola 24 Jan, the first documented in Florida's interior (ph. Terry Donovan). Other Long-billeds were coastal in Gulf 27 Dec (one; C.B.C.), Franklin 14 Jan (one, possibly the Gulf bird; JM), and Levy 28 Dec and 13 Feb (up to 2; DH et al.). For a 2nd year, at least one Semipalmated Sandpiper was verified; Kevin Karlson photographed one at Merritt 1. 26 Jan. Purple Sandpipers were scarce this season on Atlantic jetties s. to



Afforing a first record for Dry Tortugas National Park and for Monroe County, Florida, this Purple Sandpiper was digiscoped on Fort Jefferson's moat wall 17 December 2006. *Photograph by Chris Skelton*.

Brevard (DF); Gulfside, where casual, 2 lone Purples were at Anna Maria 1., Manatee 22 Jan (Gavin Spooner) and on Ft. Jefferson's moat wall 17 Dec (ph. CS), a Dry Tortugas first! The only Red Phalaropes were 10 found about 64 km off Ponce Inlet, Volusia 29 Jan (MBr, ph. BW).

A Franklin's Gull visited Gainesville 17-19

Dec (AK et al.), while up to 4 were at Cutler Ridge, Miami-Dade 23 Dec (C.B.C.) and 4 & 20 Jan (AB, RT). An Iceland Gull was at Volusia landfill 11 Jan (BW), and another was at Perdido landfill, Escambia 13 Jan-17 Feb (ph. Laura Catterton, m.ob.), the 11th accepted for Florida (F.O.S.R.C.). In nearby Santa Rosa, a Lesser Black-backed Gull was discovered 16 Dec (fide BD); rare in the panhandle, their numbers have continued to increase in the peninsula, with 27 on the upper keys 15 Dec (C.B.C.) and 20 at Stock 1., Monroe 3 Jan (CG). Up to 3 apparent Herring Gull × Glaucous Gull hybrids and one Glaucous Gull were at the Volusia landfill 13 Jan+ (MW, ph. LA et al.) Other lone Glaucous were found in Okaloosa 5-14 Feb (John & Carol Grossa et al.), Duval 25 Dec and 28 Jan (BR), and St. Lucie 5 Feb (Jane Brooks). Rare Gulfside, at least 7 Great Black-backed Gulls wintered in Pinellas (LA), with singles at Little Estero Island Wildlife Conservation Area, Lee (DSu) and in Gulf (C.B.C.). Rare, an imm. Black-legged Kittiwake was found about 64 km off Ponce Inlet 29 Jan (ph. TDu et al.).

Usually absent in the n. peninsula, Gullbilled Terms were coastal with 5 at Talbot Island S.P., *Duval* 30 Dec (Robert Wears) and singles at Hernando Beach, *Hernando* 26 Dec (BPr, BAh et al.) and Merritt 1. 18 Dec

> (C.B.C.); inland, one was on the St. Johns R., Seminole/Volusia 31 Dec (Ken Bell et al.), and 4 were at Orlando Wetlands Park, Orange 1 Feb (BHA et al.). Also inland, Royal Terns were in Hernando (one; Clay Black et al.), e. Pasco 30 Dec (2; C.B.C.), and Orange 9 Feb (one; HR); up to 9 Royals were at two Polk locations 16 Dec-7 Jan, as were up to 4 Sandwich Terns (TP, PT et al.). Rare in winter, a Common Tern was at St. Vincent N.W.R., Franklin 28 Dec (C.B.C.). Two Sooty Terns at Dry Tortugas 3 Jan (C.B.C.) were early. One of the rarer finds this season was a Dovekie, seen in flight just beyond the surf at Canaveral N.S., Brevard 18 Dec (CG).

DOVES THROUGH HUMMINGBIRDS

Florida's introduced White-winged Doves breed inland at least as far n. in the peninsula as Gainesville but remain scarce along most of the coasts. White-wingeds in Escambia (C.B.C.), Franklin (JM, J. Cooke), Wakulla (C. Evans), Levy (C.B.C.), Suannee (RR), and Duval (Anne Turner), including 21 in Okaloosa (C.B.C.), were likely w. migrants. The provenance of 2 at Dry Tortugas





With almost a dozen accepted records for Florida, this Iceland Gull was a one-day-wonder at the Volusia landfill 11 January 2007. The dark tail band, secondaries, and tertials suggest potential admixture with Thayer's Gull genes. Photographs by Bob Wallace.

17-20 Dec (CS) is an even more speculative subject. Smooth-billed Anis were in Broward (6; m.ob.) and Palm Beach (10+; fide BHA). Lone Short-eared Owls were in Wakulla (AW, JM), Orange (HR), Brevard (RH), and three locations in Monroe, the latter likely from the Antilles. Single Lesser Nighthawks were at Alligator Pt., Franklin 4 Dec (JM), Lake Apopka 22 Dec (C.B.C.) and 12 Jan (HR), and near Royal Palm, Everglades N.P., Miami-Dade 18 Dec and 16 Jan (RD). A maximum of 5 Lessers at Flamingo 14 Jan (MBe) was fewer than in recent years. A Common Nighthawk at Gainesville 19-22 Dec was audio-recorded (ES, RR; v.r. to UF) and the 2nd verified in winter; one was heard at Royal Palm 5 Jan (AB, MBe et al.). Single Commons reported in Franklin 2 Dec (JM) and Orange 3 Dec (HR) were late. The first singing Chuck-will's-widow was heard 26 Feb at Sarasota (JP).

Fred Bassett banded 100 hummingbirds: 3 Buff-bellied (Escambia [1] and Santa Rosa [2]), 37 Ruby-throated (Escambia [6], Santa Rosa [6], Okaloosa [3], Walton [2], Bay [1], Franklin [3], Leon [11], Duval [1], Alachua [2], and Hillsborough [2]); 9 Black-chinned (Escambia [3], Santa Rosa [2], and Leon [4]); 5 Calliope (Walton [2], Leon [2], and Duval [1]); and an impressive 46 Rufous (Escambia [12], Santa Rosa [1], Okaloosa [6], Walton [1], Washington [1], Franklin [1], Leon [19], Alachua [4], and Hillsborough [1]). One of Bassett's 11 recaptures (2 Ruby-throated, 2 Black-chinned, a Calliope, and 6 Rufous) from previous years was a Rufous in Escambia banded four winters ago. The fall female Buffbellied Hummingbird at Naples, Collier remained through 12 Dec (Wayne Wauligman, Steve Carbol et al.). A Downy Woodpecker at Key West 10 Jan (Brian Gibbons, BM) and a Northern Flicker on nearby Boca Chica Key 11 Feb (CG) were unexpected.

FLYCATCHERS THROUGH WAXWINGS

Least Flycatchers were n. to Brevard (one; DF) and Orange, where 11 were at Lake Apopka 22 Dec (HR). Florida's 2nd Hammond's Flycatcher was discovered on the heels of the first identified only three months before, in Pinellas. Documented by Jeffrey Gordon, it remained at Hal Scott Conservation Area, Orange 28 Jan-2 Mar (m.ob.). Vermilion Flycatchers were widespread: Wakulla (2; C.B.C.), Alachua (3; Dominick Martino, JDe et al.), with singles in Columbia (4th year; DoR, RoS), Seminole (m.ob.), Collier (2nd year; AM et al.), and Monroe (Paul Bithorn, [Bo). Ash-throated Flycatchers were all but absent this season; one was at Lake Apopka 17 Dec-31 Jan (HR), and a pale Myiarchus at Bald Point S.P., Franklin 16 Dec may have been this species (JM). Single Brown-crested Flycatchers were at Royal Palm 15-16 Jan (RT et al.), Flamingo 13 Jan-7 Feb (JBo, RD), and Lake Apopka 10 Dec-11 Feb (HR). A Tropical Kingbird stopped at Lake Apopka 13 Dec, and a Cassin's Kingbird remained there 3 Dec-14 Feb (HR), where at least one has wintered since 1999. Western Kingbirds lingered in Franklin 14, 15, & 19 Jan (up to 2; Michael Hartley, JM, AW) and in Leon 21-24 Jan (one; FD). In the n. peninsula, Westerns were at Cedar Key 28 Dec-21 Feb (up to 17 mostly subsisting on Sabal Palm fruit; DH), in Citrus 28 Dec-mid-Jan (3; Daniel Smyth et al.), Flagler 30 Jan (3; Marilyn Bontly), and s. Marion, where up to 43 visited The Villages roost with up to 3 Scissor-tailed Flycatchers (Jim Dinsmore). The kingbird roosts in Orange and Polk held up to 45 Westerns and 7 Scissor-taileds (C.B.C., BHA, DR) and up to 13 Westerns and 2 Scissor-taileds (TP), respectively. A late Scissor-tailed was found in Okaloosa in mid-Dec (C.B.C.). Rare along the e. coast, single Scissor-taileds were in Brevard 11 Dec-10 Feb (fide DF) and St. Lucie 16 Dec (C.B.C.). An early White-eyed Vireo appeared at Key West with other newly arrived migrants, including 2 Blue-headed Vireos 13 Feb (CG). One Bell's Vireo was at Bradenton, Manatee 30 Dec-3 Jan (JD et al.), while one from fall at Southern Glades Wildlife Environmental Area, Miami-Dade remained throughout (m.ob.). Single Yellow-throated Vireos wintered in Orange (HR), Polk (C.B.C.), Monroe (C.B.C.), and at two locations in Miami-Dade (JBo, RD et al.). Two Yellow-throateds on the upper keys 15 Dec (C.B.C.), as well as singles in Escambia 16 Dec (C.B.C.) and Broward 17 Dec (C.B.C.), were late; the earliest to return was in Monroe 17 Feb (AB et al.).

Up to 7 Horned Larks remained at Bascom, Jackson 14 Jan-3 Mar (AB, ph. BW, m.ob.). The earliest Purple Martins were reported 7 Jan in Duval (Patrick Leary) and Sarasota (fide Tina Mossbarger). A Northern Roughwinged Swallow in Alachua 11 Feb (CB, JDe) was early. Cave Swallows were in Broward 17 Dec (11; C.B.C.), Palm Beach in late Dec (unspecified number; C.B.C.), Monroe 30 Dec (one; C.B.C.), Miami-Dade 23 Dec and 4, 5, & 14 Jan (up to 20; C.B.C., AB, JBo), and Brevard 21 Feb (2; DF et al.). The only Barn Swallows reported were singles in Brevard 17 Jan (DS) and Volusia 27 Feb (MW). Usually restricted to the Red Hills region of n. Leon, White-breasted Nuthatches dispersed to Osceola N.F., Baker 6 Dec (2; BR), Gainesville 27 Jan (one; Bob Holt), and New Smyrna 30 Dec (one; Jean McNeil). Brown Creepers and Winter Wrens were in better-than-average numbers this season; however, both species wandered as far s. as Alachua (R. Norton, ES, H. Warren, MM), except one creeper that reached Merritt I. 18 Dec (C.B.C.).

The fall kinglet invasion continued, with

Golden-crowned Kinglets s. in Volusia (one; CB.C.), Orange, Seminole, and Lake (19; LoM, Paul Hueber, C.B.C.), Brevard (4; C B.C.), Pasco (19; BAh, BPr, C.B.C.s), Pinellas (10; LA, RoS, C.B.C.s), Hillsborough (6; Ken Allen, M. Keith, BAh), and Polk (one; C B.C.). A record high of 874 Rubycrowneds was tallied on the Gainesville C B.C. 17 Dec. In extreme s. Florida, where rare, 6 Ruby-crowneds were in Miami-Dade (JBo) and 8 in Monroe, including 6 on the keys (CG, C.B.C.s) and one at Dry Tortugas 17-20 Dec (CS). A Mountain Bluebird photographed in a yard in Ft. Myers, Lee 30 Dec (ph. Susan Hooten) was Florida's 2nd. Almost as rare, a Swainson's Thrush on the Coot Bay, Everglades N.P. C.B.C. 30 Dec was sketched by observer Rafael Galvez. Hermit Thrushes were in greater numbers than usual, with 133 in Gainesville 13 Dec (C.B.C.); one was far s. at Boca Chica Key (CG). A late Wood Thrush was at Niceville 18 Dec (Pat Baker). In late Dec, American Robins invaded the Region; driving from Cedar Key to Pensacola, the Duncans reported: "It would be difficult to estimate numbers, but surely it must be in the millions." For several days, robins passed overhead from dawn to dusk in the Orlando area (BHA), and Robinson estimated 240,000 at Lake Apopka 29 Dec. Seven robins were reported on the keys, including one at Boca Chica Key 7 Dec (BM, CG). Also rare in the keys, an American Pipit was at Dry Tortugas 17-20 Dec (CS). The only Sprague's Pipit was at Eglin A.F.B., Okaloosa 3 Dec (Lenny Fenimore). Cedar Waxwings invaded the state in greater numbers than usual, with 12 s. to Key West 30 Dec (CG). Surprising was a Cedar over open ocean 40 km off New Smyrna, Volusia 29 Jan (MG et al.).

WARBLERS THROUGH FINCHES

Twenty-one species of warblers were reported this season. Nashville Warblers were in Alachua (one; B. Christensen), Pasco (one; D Gagne), Orange (one; BHA, BO et al.), Miami-Dade (3; RD, JBo), and Monroe (one; C B.C.). Late Northern Parulas lingered until 17 Dec in Hillsborough (BAh et al.) and Seminole (Paul Hueber), where another was the farthest n. reported to winter (BHA); the earliest to return were 4 males at Bill Baggs Cape Florida S.P., Miami-Dade 26 Jan (RD). Single Yellow Warblers were found n. to Alachua 8 Dec+ (J. Powell, AK et al.), Seminole 31 Dec, and s. Brevard 24 Jan (AB et al). Wintering Magnolia Warblers were in Collier (one; fide AM), Miami-Dade 16 & 23 Dec (4; C.B.C.s), and Monroe (2; C.B.C.); one at Key West 13 Feb was with other

newly arrived migrants (CG). Lone Blackthroated Blue Warblers in Orange 6 & 16 Dec (Andrew Boyle, C.B.C.) and Brevard 10-21 Feb (DF) were likely migrants. Although Yellow-rumped Warblers arrived on schedule, large numbers were not noted until after the new year; Lillian and Donald Stokes estimated that 10,000 flew over the Sanibel Lighthouse, Lee towards the mainland 28 Jan. Black-throated Green Warblers wintered n. to Orange (BO) and Brevard (Gerry Day, LoM et al.). A Yellow-throated Warbler and an American Redstart were seen at Key West 13 Feb with a flock of newly arrived migrants (CG). Prairie Warblers wintered in greater-than-normal numbers in the cen. peninsula; farther n., one was at Fort Walton Beach Sanitation Facility, Okaloosa (BD), and 2 were at Jacksonville (C.B.C.). A Yellow Palm Warbler wintered at Key West (ph. BM, CG). American Redstarts wintered n. to Citrus (2; P. Pilny, BAh) and Orange (3; Jeffrey Gordon, Tom Rodriguez, C.B.C.); one farther n. in Alachua 17 Dec (S. Robinson) was late. Two Worm-eating Warblers were in Monroe (C.B.C.). Although banding operations at Bill Baggs Cape Florida ceased for the winter months, wintering was suspected when a Swainson's Warbler, captured there several times during the fall, last on 3 Nov, was recaptured 12 Mar, shortly after operations resumed (RD). Lone Ovenbirds were in the panhandle in Okaloosa 18 Dec (Jim Kowalski) and Franklin 3 Feb (JM). Single Northern Waterthrushes wintered n. to Wakulla (AW et al.) and Alachua (AK); 21 were counted in Orange at Lake Apopka (C.B.C.). Single Louisiana Waterthrushes were at Miami, Miami-Dade 16 Dec (JBo, G. Jones, D. Jones), Gainesville 17 Dec (MM), Royal Palm 18 & 20 Dec and 5 Jan (RD, C.B.C., AB), and Coot Bay 30 Dec (C.B.C.). The first spring Louisianas were noted 25 Feb, with one at Lutz, Hillsborough (Anya & Tom Mason) and 2 at Lake Apopka (HR). Lone Hooded Warblers were in Pinellas at John H. Chesnut, Jr. Park 8-14 Jan (TA) and at Largo throughout the season (KN), and another was at Fakahatchee Strand Preserve S.P., Collier 4 Feb (AM). Wilson's Warblers were scarce, with singles in Okaloosa (Sandra Lefstad), Alachua (JoH, MM et al.), Volusia (BO), Orange (HR), and Miami-Dade (AB et al.); there were 3 in Pinellas (CoG) and 2 in Monroe (Key West and Dry Tortugas C.B.C.s). Lone Yellow-breasted Chats were in Franklin (JM), Bradford (C.B.C.), Orange (HR), and Pinellas (ph. Jay Jones), with duos in Alachua (MM, P. Colverson), Miami-Dade (RD), and Monroe (BM, C.B.C.). Twenty Summer Tanagers were re-

ported from 13 counties, from Okaloosa (Gene Estes) and Alachua (Ingrid Scales) s to Collier and Miami-Dade (C.B.C.). One banded male Summer in Leon returned for the 5th year (Peter Homann)! Western Tanagers pleased birders at Niceville 31 Dec (Robin & Tom Stiles), Pensacola 18 Jan (Paula Palmer), Paines Prairie Preserve S.P. Alachua 2 Dec (BRo, Patty Orr, Carolyn Retey), Gainesville 3 Dec (Ron Robinson), L. Disston, Flagler 10 Feb (Erik Haney, DG), St. Augustine 20 Feb (Geoff Carter), e of Naples 22 Feb (SM et al.), and Tallahassee at three locations: 19 Dec-21 Feb (male and female, FR, m.ob.); 23-26 Feb (female, J. Cavanagh); and 25 Feb (male, FD). A black-backed Western Spindalis, likely of the nominate Bahamian subspecies, was found at Bill Baggs Cape Florida S.P. 12 Dec and 9 &14 Feb (RD). At Fort Lauderdale, a green-backed Western Spindalis was seen 19 Jan-11 Feb (reappeared in Apr; m.ob) and was apparently of the subspecies townsendi.

While most sparrow species were not particularly common anywhere, 11 reached the southernmost peninsula and keys, where they are rare. Chipping Sparrows were reported s. to Collier (12; DSu) and Monroe (one; RD). Clay-colored Sparrows were in Orange (up to 4; HR), Hillsborough (2; Steve Backes, TA), Polk (5; C.B.C.), Sarasota (one, LA, JD), Miami-Dade (one; MBe et al.), and Monroe (up to 7, LM). A Vesper Sparrow was s. to Flamingo (JBo). Lone Lark Sparrrows were in Leon (GM, JCa), Hillsborough (TA), Manatee (JD et al.), and Sarasota (perhaps 2, JD, Margaret Dunson). Henslow's and Le Conte's Sparrows were widely distributed in the peninsula, with Henslow's located in 10 counties from Alachua (17; C.B.C.) to Okeechobee, where 33 were banded (Marianne Korosy). Le Conte's were in Hamilton (2, C.B.C.), Alachua (up to 4; JoH, CB, Peter Polshek et al.), Seminole (6; C.B.C.), Hernando (up to 3; BAh, DS et al.), and Miami-Dade (one; ph. LM). Gulfside, Saltmarsh Sharp-tailed Sparrows were at Cedar Key (one; C.B.C.), Fort Island Beach (one; RoS, BAh), and Sandy Key, Everglades N.P., Monroe (one; BM). A Nelson's Sharp-tailed Sparrow was far s. in Monroe (one; JBo, RT et al) Fox Sparrows were widely reported in the panhandle; however, 3 in Alachua reached their usual s. limit in the peninsula. Song Sparrows were s. to Sarasota and Polk/Highlands (C.B.C.s) and at Boca Chica Key (one; CG). Lincoln's Sparrows were in Alachua (3; CB, MM, William Bolte et al), Seminole (one; RH), Orange (up to 2; HR), Manatee (2; CoG, JD et al.), and at two Miami-Dade locations (2; AB et al., C.B.C.). Two Swamp Sparrows were s. to Flamingo (JBo). White-throated Sparrows were in above-average numbers in the cen. peninsula, wandering farther s. to Charlotte (one; RoS, BAh) and Monroe at Clive Key, Everglades N.P. (one; BM). By contrast, White-crowned Sparrows were scarce, with reports of lone birds in Alachua (BAh, RR, RoS), Brevard (DS), Pasco (KT), and Pinellas (RoS), up to 5 in Orange (HR), and up to 4 in Monroe (BM, RD, AB et al.). Dark-eyed Juncos were widespread and also in above-average numbers in the panhandle but were rare in the peninsula, with reports from n. Baker (one; BR), Columbia (one; C.B.C.), and Pinellas (3 in two locations; AB, LA, TA).

Rose-breasted Grosbeaks were in St. Lucie 16 Dec (C.B.C.), Okaloosa 18 Dec (Bob Larson), Pinellas 7 Jan (KN), Orange 16 Jan (BHA), and Leon 21 Feb (C. Clarke). Blue Grosbeaks were scarce, with only singles in Orange 1 Dec-4 Feb (HR) and Hillsborough 17 Dec (BAh). A Painted Bunting was far n. in Okaloosa 18 Dec (C.B.C.). Lone Dickcissels were in Orange 9 & 11 Feb (Joie Clifton) and Sarasota at two locations 2 Dec-24 Jan and 1 Feb (Edith Miller, Janet Carlock, Owen Comora); up to 3 were near Flamingo 14 Jan-11 Feb (BRo, LM). Not verified in Florida since 1977, a Western Meadowlark remained in n. Escambia 19 Jan-22 Feb (m.ob.) and was the 5th accepted for Florida. Single Yellow-headed Blackbirds were in Orange 3 & 6 Dec (HR) and Hernando 26 Dec (BAh, BPr et al.). Rusty Blackbirds were in good numbers: about 50 in Escambia (Bob & Lucy Duncan); 104 in Bay (C.B.C.); 100 in Leon (AW); 70 at two Wakulla locations (SM, AB, AW); up to 27 in Alachua (CB, RR et al.); 8 in Sumter (DG, Eric Haney); and one in Pinellas (BAh, BPr et al.). Brewer's Blackbirds were limited to the panhandle: in n. Escambia, about 75 in a mixed blackbird-species flock 19 Jan (B&LRD) and a pure flock of 122 on 4 Feb (B&LD, DoR, JM); 3 in Okaloosa 3 Jan (BD); and 2 in Bay 30 Dec (C.B.C.). Up to 7 Shiny Cowbirds were reported in Miami-Dade throughout (LM); singles were in Pasco 16 Dec (KT) and Highlands 28 Dec (C.B.C.). Only recently established in extreme s. Florida, 51 Bronzed Cowbirds were counted in Miami-Dade 16 Dec (C.B.C.); others were in Monroe (one; JBo), Collier (6 at two locations; AM), Hendry (3; Brent Bomkamp), Polk (5; C.B.C.), Hillsborough (2; D. Bowman et al.), and Orange 23 Dec (one; Steve Gast). Up to 2 Orchard Orioles were at Flamingo 30 Dec-27 Jan (JBo, ph. BM); one at Gainesville 14 Feb was early (Grace Kiltie). A female Bullock's Oriole lingered at Tallahassee 4 Jan–21 Feb (FR, m.ob.). Rare, Purple Finches were in Okaloosa (unstated number; C.B.C.), Bay (3; C.B.C.), Leon (12 at two locations, AW, G. Menk), Wakulla (one; G. Menk), and Alachua 12 Feb+ (one; Ron Robinson). House Finches have been slow to colonize the e. peninsula; not yet verified breeding, 30 were counted on three Volusia C.B.C.s. Gulfside, where House Finches are established s. to Sarasota, 2 were farther s. in Lee 11 Feb (Layne Hamilton). The only Pine Siskins were 2 in Escambia 3 Feb (RoS) and singles in Leon 30 Dec (C.B.C.) and Orange (HR).

Corrigenda: In the winter 2005-2006 report (N.A.B. 60: 220), the last date that the Cackling Goose was reported should read "4 Jan," not 14 Jan. In addition to the incorrect date, the caption of the photograph of the geese on the next page should read that the Cackling Goose would provide Florida's "third record," not second. In the same report, it was not a Bell's Vireo but a Yellow-throated Vireo that appeared at Bill Baggs Cape Florida S.P. 28 Feb.

Cited contributors and members of the Florida Ornithological Society Field Observations Committee: Brian Ahern (BAh), Bruce Anderson (BHA), Todd Arcos, Lyn Atherton, Andy Bankert, Mark Berney (MBe), Dave Bowman, John Boyd (JBo), Michael Brothers (MBr), Chris Burney, Jim Cavanagh (JCa), K. Dailey, John DeLuca (JDe), Robin Diaz, Fred Dietrich, Jeanne Dubi, Bob & Lucy Duncan (B&LD), Thomas J. Dunkerton (TDu), C. Evans, Charlie Ewell, David Freeland, Murray Gardler, Colin Gjervold (CoG), Carl Goodrich, Dave Goodwin, Jeffrey Gordon, Roy Halpin, Al & Bev Hansen (A&BH), Dale Henderson, John Hintermister (JoH), Brian Hope, Paul Hueber, Grace Kiltie, Andy Kratter, Bob Landry, Lorne Malo (LoM), Mike Manetz, Larry Manfredi, Sean McCool, Gail Menk, Paul Miller (PMi), Trey Mitchell, Roy Morris II, Brennan Mulrooney, John Murphy, Alan Murray, Kris Nelson, Brenda Orr, Jeff Palmer, Tom Palmer, Cheri Pierce, Peggy Powell, Bill Pranty (BPr), Dexter Richardson, Bob Richter, Dotty Robbins (DoR), Bryant Roberts (BRo), Harry Robinson, Rex Rowan, Fran Rutkovsky, Earl & Ingrid Scales (E&IS), David Simpson, Chris Skelton, Ron Smith (RoS), Doug Suitor (DSu), Pete Timmer, Roberto Torres, Ken Tracey, Bob Wallace, Meret Wilson, Andy Wraithmell.

Bruce H. Anderson, 2917 Scarlet Road Winter Park, Florida 32792, (scizortail@aol.com)



GET DOWN AND DIRTY FOR AMERICA.

And when you get you hands dirty in programs like National Public Lands Day, you help spruce up your beautiful lands. Last year, on this day, nearly 80,000 volunteers built trails, bridges, planted native trees and removed trash.

To find out how you can help, go to www.npld.com or call 800-VOL-TEER (800-865-8337).

HELPING HANDS FOR AMERICA'S LANDS



© 2004 Muench Photography, Inc.