

# ORNITOLOGIA NEOTROPICAL

EDITOR

RAYMOND McNEIL

## **INDEX TO VOLUME 15, 2004**

(Authors, key words, families, common names, latin names)

An International Journal of Neotropical Ornithology  
published by

**THE NEOTROPICAL ORNITHOLOGICAL SOCIETY**

The “S” prefix before page numbers refers to pages in the Proceedings of the VII<sup>th</sup> Neotropical Ornithological Congress held in Chile, 5–11 October 2003, and published as a supplement to Volume 15 (2004) of *Ornitología Neotropical*.

- Abril, M., S377–S384  
 Abundance, 93–102, 371–380  
 Accipiters, 331–347  
*Accipiter bicolor*, 31–40, 173–200, 381–398, 447–482, 501–512, S83–S99  
*Accipiter chilensis*, S83–S99  
*Accipiter chionogaster*, S83–S99  
*Accipiter collaris*, 31–40, S83–S99  
*Accipiter cooperi*, S267–S278, S83–S99  
*Accipiter erythronemius*, S83–S99  
*Accipiter gentilis*, S83–S99  
*Accipiter gundlachi*, S83–S99  
*Accipiter poliogaster*, S83–S99  
*Accipiter striatus*, , 501–512, S83–S99, S267–S278  
*Accipiter superciliosus*, 447–482, S83–S99  
*Accipiter ventralis*, S83–S99  
*Aeestrura mulsant*, S205–S213  
*Acropternis orthonyx*, 483–492  
*Actitis macularia*, 173–200, 381–398, 447–482  
 Adaptation, S191–S198  
*Adelomyia melanogenys*, 483–492  
*Agamia agami*, 447–482  
*Agelaioides badius*, 381–398  
*Agelaius phoeniceus*, 447–482  
*Agelaius thibius*, 271–278  
 Age-structured, 289–297  
 Aggressive interactions, 539–542  
*Aglaeactis cupripennis*, 483–492, S205–S213  
 Agriculture, S309–S315  
 Agricultural landscape, S117–S126  
 Agriculture, S301–S307  
*Agriornis montana*, 513–525  
*Agriornis murinus*, 381–398  
 Agroecosystems, 371–380, 493–500, S301–S307  
 Agrosystems, 65–70  
 Aillapan, L., S419–S434  
*Aimophila botterii*, 447–482  
*Aimophila rufescens*, 447–482  
*Aimophila sumichrasti*, S175–S181  
*Ajaia ajaja*, S223–S232  
 Alanota, G., S339–S347  
*Alectrurus risora*, S145–S157  
 Alegre, B., 103–110  
 Allozymes, S169–S174  
*Alopochelidon fucata*, 173–200  
 Altiplano, S339–S347  
 Altitudinal migrations, S205–S213  
 Alvarez, M. E., 501–512  
*Amaurolimnas concolor*, 447–482  
*Amaurospiza concolor*, 447–482  
*Amazilia amabilis*, 447–482  
*Amazilia candida*, 447–482  
*Amazilia chionogaster*, 381–398  
*Amazilia cupreicauda*, 173–200  
*Amazilia cyanocephala*, 447–482  
*Amazilia fimbriata*, 173–200  
*Amazilia rutila*, 447–482  
*Amazilia tzacatl*, 447–482  
*Amazilia versicolor*, 173–200  
*Amazilia viridifrons*, S175–S181  
*Amazona aestiva*, 381–398  
*Amazona amazonica*, 173–200  
*Amazona auropalliata*, 447–482  
*Amazona autumnalis*, 447–482  
*Amazona farinosa*, 447–482  
*Amazona finschi*, S175–S181  
*Amazona ocbrocephala*, 173–200  
*Amazona vittata*, 289–297, S455–S465  
 Amazonas, 265–267  
*Amazonetta brasiliensis*, 71–85, 173–200, 381–398  
 Amazonia, S317–S321  
 Amazonian gamebirds, S323–S328  
*Amblycercus bolosericus*, 447–482  
*Ammodramus humeralis*, 173–200  
*Ammodramus savannarum*, 447–482  
*Ampelion rubrocristatus*, 483–492  
*Anabacerthia variegaticeps*, 447–482

- Anairetes parulus*, 483–492, 513–525  
*Anas acuta*, 447–482  
*Anas andium*, 483–492  
*Anas bahamensis*, 381–398  
*Anas cyanoptera*, 381–398  
*Anas discors*, 447–482  
*Anas flavirostris*, 209–218, 543–545  
*Anas georgica*, 513–525, 543–545  
*Anas sibilatrix*, 513–525  
*Anas specularis*, 513–525  
Anatidae, 209–218  
Anatomical studies, S73–S81  
Anatomy, S73–S81  
Andean beliefs, S409–S417  
Andean foothills, 381–398  
Anderson, D. L., 447–482  
Andes, 381–398, 349–356, 403–406, 483–492, S191–S198  
*Andigena hypoglanca*, 527–537  
*Andigena laminirostris*, 527–537  
Andrade, A., 87–92  
*Androdon aequatorialis*, S191–S198  
*Anhinga anhinga*, 71–85, 173–200, 447–482  
*Anisognathus igniventris*, 483–492  
*Anisognathus somptuosus*, 483–492  
Anjos, L. dos, 71–85, S137–S144  
*Anodorhynchus hyacinthinus*, S279–S283  
*Anodorhynchus leari*, S279–S283  
*Anous stolidus*, S61–S72  
*Anous tenuirostris*, S61–S72  
Antagonists, S199–S204  
Antarctica, 163–171  
Antbird, S127–S135  
*Anthracothorax nigricollis*, 173–200  
*Anthracothorax prevostii*, 447–482  
*Anthus correndera*, 513–525  
*Anthus bellmayri*, 513–525  
*Anthus lutescens*, 173–200  
Antioquia, 413–415  
*Anurolimnas viridis*, 173–200  
*Aphanotriccus capitalis*, 447–482  
Aponte, O., 247–251  
*Aquila chrysaetos*, S83–S99  
*Ara ambigua*, 447–482  
*Ara ararauna*, 173–200, S279–S283  
*Ara chloroptera*, 173–200, S279–S283  
*Ara macao*, 173–200, 447–482, S279–S283  
*Ara severus*, 381–398  
*Aramides cajanea*, 173–200, 381–398, 447–482  
*Aramus guarauna*, 71–85, 173–200, 381–398, 447–482  
*Aratinga aurea*, 173–200  
*Aratinga finschi*, 447–482  
*Aratinga leucophthalma*, 71–85, 381–398, S117–S126  
*Aratinga nana*, 447–482  
*Aratinga pertinax*, 173–200  
*Aratinga solstitialis*, 173–200  
*Archaeopteryx*, S101–S116  
*Archilochus colubris*, 447–482  
Archosauria, S101–S116  
Archosauromorpha, S101–S116  
*Ardea alba*, 173–200, 381–398, 447–482  
*Ardea cocoi*, 71–85, 93–102, 103–110, 121–131, 139–141, 173–200, 381–398  
*Ardea herodias*, 447–482  
*Arenaria interpres*, 447–482  
Arendt, W. J., S455–S465  
Argentina, 145–153, 209–218, 271–278, 299–306, 307–311, 357–364, 371–380, 501–512, S145–S157, S309–S315, S367–S376  
*Arremon aurantirostris*, 447–482  
*Arremon flavirostris*, 381–398  
*Arremon taciturnus*, 173–200  
*Arremonops conirostris*, 447–482  
Artificial clutch, 201–208  
*Arundinicola leucocephala*, 71–85, 173–200  
*Asio flammeus*, 501–512  
*Asio stygius*, 173–200, 423–427, 447–482, 501–512  
Associations, 163–171  
*Asthenes baeri*, 145–153  
*Asthenes pyrrholaema*, 513–525  
*Asturina nitida*, 381–398, 447–482  
*Athene cunicularia*, 173–200, 381–398  
Atlantic forest, 547–548, S117–S126, S127–S135, S137–S144  
*Atlapetes latinuchus*, 483–492  
*Atlapetes leucopterus*, 483–492  
*Attila cinnamomeus*, 173–200

- Attila spadiceus*, 173–200, 447–482  
 Audio, S455–S465  
*Aulacorhynchus derbianus*, 527–537  
*Aulacorhynchus haematopygus*, 483–492  
*Aulacorhynchus prasinus*, 433–445, 527–537  
 Austral migration, S349–S355  
 Austral staging areas, S357–S365  
*Automolus infuscatus*, 173–200  
*Automolus leucophthalmus*, 71–85  
*Automolus ochrolaemus*, 173–200, 447–482  
*Automolus rubiginosus*, 173–200  
*Automolus rufipileatus*, 173–200  
 Avifauna, 173–200, 381–398, 447–482, S395–  
 S407, S419–S434, S435–S444  
*Aythya affinis*, 447–482  
 Azpiroz, A. B., 269–270  
*Baillonius bailloni*, 527–537  
*Baillonius*, 527–537  
 Baker, A. J., S357–S365  
 Bala, L. O., 357–364  
*Balaenoptera acutorostrata*, 163–171  
*Balaenoptera musculus*, 163–171  
 Balbuena, R. A., S367–S376  
 Barnett, J. M., 257–263  
 Barrio, J., 111–119  
*Bartramia longicauda*, 173–200, 381–398, 447–  
 482, S159–S167  
*Baryphthengus martii*, 447–482  
*Basileuterus bivittatus*, 381–398  
*Basileuterus coronatus*, 483–492  
*Basileuterus culicivorus*, 5–15, 173–200, 447–482  
*Basileuterus flaveolus*, 247–251  
*Basileuterus fulvicauda*, 447–482  
*Basileuterus leucoblepharus*, 5–15  
*Basileuterus nigrocristatus*, 483–492  
*Batara cinerea*, 381–398  
 Beauchamp, G., 407–411  
 Bechard, M. J., 447–482  
 Behavior, 283–284, S145–S157, S243–S250,  
 S329–S337  
 Belize, 155–162  
 Bern, C., S377–S384  
 Bertelli, S., S285–S291, S293–S299  
 Bildstein, K. L., S83–S99  
 Bill morphology, 543–545  
 Binocular, S233–S242  
 Binocularity, S233–S242  
 Biocultural conservation, S395–S407, S409–  
 S417  
 Biodiversity, S127–S135  
 Biogeography, 173–200  
 Bird abundance, 513–525  
 Bird guilds, S137–S144  
 Bird origin, S101–S116  
 Bird richness, 371–380  
 Birds, 483–492  
 Blanco, D. E., S159–S167, S367–S376  
 Blending, P. G., 501–512  
*Bolborhynchus lineola*, 483–492  
 Bolivia, 381–398, 399–402, S261–S266,  
 S339–S347  
*Bombycilla cedrorum*, 447–482  
 Bonaccorso, E., 483–492  
*Botaurus pinnatus*, 173–200  
 Bozinovic, F., S215–S222  
 Brabata, G., 237–246  
*Brachygalba lugubris*, 173–200  
 Branco, F., S267–S278  
 Braun, M. M., 173–200  
 Brazil, 5–15, 65–70, 71–85, 219–224, 265–267,  
 279–282, 423–427, 547–548, S73–S81,  
 S117–S126, S127–S135, S137–S144,  
 S279–S283, S367–S376  
 Breeding, 133–137, 365–370, S145–S157  
 Breeding behavior, 349–356  
 Breeding biology, 219–224, 253–256, 283–  
 284  
 Breeding colony, 133–137  
 Brennan, P. L. R., S329–S337  
 Brewer, G. L., 209–218  
 Brightsmith, D. J., 319–330  
 Brood parasitism, 331–347  
 Brooks, D. M., S323–S328  
*Brotogeris sanctithomae*, 319–330  
*Brotogeris chrysopterus*, 173–200  
*Brotogeris cyanoptera*, 319–330  
 Brumfield, R. T., 381–398  
*Buarremon torquatus*, 483–492  
*Bubo virginianus*, 173–200, 279–282, 447–482  
*Bubulcus ibis*, 173–200, 381–398, 447–482,

- S223–S232  
*Buccinanops globulosum*, 357–364  
*Bucco capensis*, 173–200  
*Bucco tamatia*, 173–200  
Budget, S1–S34  
*Burhinus bistriatus*, 173–200, S223–S232  
*Busarellus nigricollis*, 173–200, 447–482, S83–S99  
Bustamente, J., 513–525  
*Buteo albicaudatus*, 173–200, 447–482, S83–S99  
*Buteo albigula*, S83–S99  
*Buteo albonotatus*, 173–200, S83–S99, S267–S278  
*Buteo brachyurus*, 173–200, 447–482, 501–512, S83–S99  
*Buteo galapagoensis*, S83–S99  
*Buteo jamaicensis*, 447–482, S83–S99, S267–S278  
*Buteo leucorrhous*, 31–40, S83–S99  
*Buteo lineatus*, S83–S99  
*Buteo magnirostris*, 31–40, 173–200, 381–398, 447–482, 501–512, S83–S99  
*Buteo nitidus*, 173–200, S83–S99  
*Buteo platypterus*, 173–200, 447–482, S83–S99, S267–S278  
*Buteo poecilochrous*, S83–S99  
*Buteo polyosoma*, 483–492, 501–512, 513–525, S83–S99  
*Buteo regalis*, S83–S99  
*Buteo ridgwayi*, S83–S99  
*Buteo swainsoni*, S83–S99, S267–S278  
*Buteo ventralis*, S83–S99  
*Buteogallus aequinoctialis*, S83–S99  
*Buteogallus anthracinus*, 447–482, S83–S99  
*Buteogallus meridionalis*, 173–200, 381–398, S83–S99  
*Buteogallus subtilis*, S83–S99  
*Buteogallus urubitinga*, 71–85, 173–200, 381–398, 447–482, 501–512, S83–S99  
*Buthraupis montana*, 483–492  
*Butorides striatus*, 71–85, 173–200, 381–398  
*Butorides virescens*, 447–482  
Bylaws, S35–S40  
*Cacicus cela*, 41–52, 173–200  
*Cacicus haemorrhous*, 41–52, 173–200  
*Cacicus uropygialis*, 447–482  
*Cairina moscbata*, 71–85, 173–200, 447–482  
*Calidris alba*, 447–482  
*Calidris bairdii*, 447–482, 513–525, S377–S384  
*Calidris canutus*, 447–482, S357–S365  
*Calidris fuscicollis*, 173–200, 357–364, 381–398, 447–482, S377–S384  
*Calidris himantopus*, 173–200, 447–482  
*Calidris mauri*, 447–482, S385–S394  
*Calidris melanotos*, 173–200, 381–398, 447–482, S159–S167, S377–S384  
*Calidris minutilla*, 173–200, 447–482  
*Calidris pusilla*, 447–482  
*Calliphlox amethystina*, 173–200  
*Callonetta leucophrys*, 381–398  
*Campephilus guatemalensis*, 447–482  
*Campephilus imperiales*, S175–S181  
*Campephilus leucopogon*, 381–398  
*Campephilus melanoleucos*, 173–200  
*Campephilus rubricollis*, 173–200  
Camperi, A. R., 299–306  
Campo rupestre, 399–402  
*Campostoma imberbe*, 447–482  
*Camptostoma obsoletum*, 71–85, 173–200, 381–398, S117–S126  
*Campylopterus curvipennis*, 447–482  
*Campylopterus hemileucurus*, 447–482, S191–S198  
*Campylorhynchus rocurvoides*, 173–200  
*Campylorhynchus trochilrostris*, 71–85  
*Campylorhynchus griseus*, 173–200  
*Campylorhynchus yucatanicus*, S175–S181  
*Campylorhynchus zonatus*  
Canopy access, 41–52  
*Capito auratus*, 41–52  
*Capito niger*, 173–200  
Capitonidae, 283–284  
Capllonch, P., 501–512  
*Caprimulgus carolinensis*, 447–482  
*Caprimulgus cayennensis*, 173–200, S251–S260  
*Caprimulgus longirostris*, 483–492, 513–525  
*Caprimulgus maculicaudus*, 447–482  
*Caprimulgus parvulus*, 71–85  
*Caprimulgus rufus*, 381–398  
*Caprimulgus vociferus*, 447–482

- Capsiempis flaveola*, 173–200, 447–482  
 Captive breeding, S339–S347  
 Capture-recapture, S357–S365  
*Caracara cheriway*, 173–200, 447–482  
*Caracara plancus*, 381–398, 501–512, 513–525  
 Carbajal, M., S357–S365  
*Cardinalis phoeniceus*, 253–256  
*Carduelis cucullata*, 173–200  
*Carduelis magellanica*, 271–278, 371–380, 381–398, 483–492  
*Carduelis notata*, 447–482  
*Cariama cristata*, 381–398  
 Caribbean, 289–297  
 Carmona, R., 237–246  
*Carpodectes nitidus*, 447–482  
*Caryothraustes canadensis*, 173–200  
*Caryothraustes polio-gaster*, 447–482  
 Cashca Totoras, 483–492  
*Casiornis rufus*, 381–398  
 Castro, V., S409–S417  
*Casutitermes corniger*, 319–330  
*Catamenia inornata*, 483–492  
 Cataño-B, E. J. F., 413–415  
 Catari, J. C., 381–398, 399–402  
*Cathartes aura*, 31–40, 71–85, 173–200, 381–398, 447–482, 501–512, S83–S99, S267–S278  
*Cathartes burrovianus*, 173–200, 447–482, S83–S99  
*Cathartes melambrotus*, 173–200, S83–S99  
 Cathartidae, 219–224  
*Catharus fuscescens*, 173–200, 247–251  
*Catharus minimus*, 173–200, 447–482  
*Catharus ustulatus*, 381–398, 447–482  
*Catoptrophorus semipalmatus*, 447–482, S223–S232  
 Cavity adopter, S455–S465  
 Cavity nesting, S455–S465, S477–S483  
*Celens castaneus*, 447–482  
*Celens elegans*, 173–200  
*Celens flavescens*, S117–S126  
*Celens flavus*, 173–200  
*Celens loricatus*, 447–482  
*Celens torquatus*, 173–200  
*Celens undatus*, 173–200  
*Cephalopterus glabricollis*, 433–445  
*Cephalopterus ornatus*, 173–200  
*Cercibis oxycerca*, 173–200  
*Cercomacra cinerascens*, 173–200  
*Cercomacra laeta*, 173–200  
*Cercomacra tyrannina*, 173–200, 447–482  
 Cerrado, 219–224, 399–402, 423–427  
*Certhiaxis cinnamomea*, 173–200  
*Ceryle alcyon*, 447–482  
*Ceryle torquata*, 71–85, 173–200, 381–398, 447–482  
 Cetaceans, 163–171  
 Chaco, 381–398  
*Chaetocercus mulsant*, 483–492  
*Chaetura brachyura*, 173–200  
*Chaetura cinereiventris*, 173–200, 447–482  
*Chaetura pelagica*, 447–482  
*Chaetura spinicauda*, 173–200  
*Chaetura vauxi*, 447–482  
*Chalcostigma herrani*, S205–S213  
*Chalybura urochrysa*, 447–482  
*Chamaepetes unicolor*, 433–445  
*Charadrius alexandrinus*, 447–482  
*Charadrius collaris*, 71–85, 173–200, 447–482  
*Charadrius falklandicus*, 357–364, 513–525  
*Charadrius modestus*, S159–S167  
*Charadrius semipalmatus*, 447–482  
*Charadrius vociferus*, 447–482  
*Charadrius wilsonia*, 447–482, S223–S232  
*Chauna torquata*, 381–398  
 Chaves-Campos, J., 433–445  
*Chelidoptera tenebrosa*, 173–200  
 Chiappe, L. M., S101–S116  
 Chile, 313–317, S1–S34, S395–S407, S419–S434, S435–S444  
 Chilean cultural anthropology, S409–S417  
 Chiribiquete Natural National Park, S183–S190  
 Chiroptera, 87–92  
*Chiroxiphia pareola*, 173–200  
*Chlidonias niger*, 447–482  
*Chloephaga picta*, 513–525  
*Chlorestes notatus*, 173–200  
*Chloroceryle aenea*, 447–482  
*Chloroceryle amazona*, 71–85, 173–200, 447–482

- Chloroceryle americana*, 173–200, 447–482  
*Chloroceryle inda*, 173–200, 447–482  
*Chlorophanes spiza*, 41–52, 173–200, 447–482  
*Chlorornis riefferii*, 483–492  
*Chlorostilbon aureoventris*, 381–398  
*Chlorostilbon canivetii*, 447–482  
*Chlorostilbon mellisugus*, 173–200  
*Chlorostilbon olivaresi*, S183–S190  
*Chlorothraupis carmioli*, 447–482  
*Chondrohierax uncinatus*, 447–482, S83–S99  
*Chordeiles acutipennis*, 173–200, 447–482, S251–S260  
*Chordeiles minor*, 447–482  
*Chordeiles pusillus*, 173–200  
 Christoff, A. U., 65–70  
*Chrysolampis mosquitus*, 173–200  
*Chrysomus cyanopus*, 381–398  
*Chunga burmeisteria*, 381–398  
*Ciccaba nigrolineata*, 447–482  
*Ciccaba virgata*, 447–482  
 Cicchino, A. C., 299–306  
*Ciconia maguari*, 71–85, 381–398  
*Cinclodes excelsior*, 483–492  
*Cinclodes fuscus*, 483–492  
*Cinclus leucocephalus*, 483–492  
*Cinclus mexicanus*, 447–482  
*Cinnyerthia unirufa*, 483–492  
*Circus buffoni*, S83–S99  
*Circus cinereus*, 501–512, 513–525, S83–S99, S267–S278  
*Circus cyaneus*, 447–482  
*Cistothorus platensis*, 447–482, 483–492  
 Cladistics, S285–S291  
 Clams, 357–364  
*Claravis pretiosa*, 173–200, 381–398, 447–482  
 Cloud forest, 403–406  
 Clutch size, 155–162, S41–S60  
*Cnemarchus erythropygius*, 483–492  
*Cnemotriccus fuscatus*, 381–398  
*Coccyzus americanus*, 447–482  
*Coccyzus melacoryphus*, 381–398  
*Coccyzus minor*, 447–482  
*Cochlearius cochlearius*, 447–482  
 Codesido, M., 139–141  
*Coeligena iris*, 483–492  
*Coeligena lutetiae*, S205–S213  
*Coereba flaveola*, 173–200, 447–482  
*Colaptes campestris*, 71–85, 173–200, S117–S126  
*Colaptes melanochloros*, 71–85  
*Colibri coruscans*, 483–492, S205–S213  
*Colibri delphinae*, 447–482  
*Colinus cristatus*, 173–200  
*Colinus nigrogularis*, 447–482  
 Colombia, 17–30, 413–415, S183–S190, S349–S355  
*Colonia colonus*, 173–200, 447–482  
 Color vision, S215–S222  
*Columba cayennensis*, 71–85, 173–200, S117–S126  
*Columba fasciata*, 483–492  
*Columba livia*, 447–482  
*Columba picazuro*, 71–85, 371–380, S117–S126  
*Columba plumbea*, 173–200  
*Columba speciosa*, 173–200  
*Columba subvinacea*, 173–200  
*Columbina minuta*, 173–200, 447–482  
*Columbina passerina*, 173–200, 447–482  
*Columbina picui*, 71–85, 381–398  
*Columbina talpacoti*, 71–85, 173–200, 381–398, 447–482, S117–S126  
 Combined analysis, S293–S299  
 Community ecology, S323–S328  
 Condori, C., S339–S347  
 Cone, S223–S232, S251–S260  
*Conirostrum cinereum*, 483–492  
*Conirostrum speciosum*, 71–85, 173–200  
*Conopias alborittata*, 447–482  
*Conopias parva*, 173–200  
*Conopias trivirgata*, 71–85  
 Consequences, S83–S99  
 Conservation, 17–30, 71–85, 501–512, 547–548, S145–S157, S159–S167, S175–S181, S279–S283, S317–S321, S367–S376  
 Conservation assessment, 41–52  
*Constrictotermes cavifrons*, 319–330  
*Contopus cinereus*, 173–200, 447–482  
*Contopus cooperi*, 447–482  
*Contopus fumigatus*, 381–398  
*Contopus sordidulus*, 447–482  
*Contopus virens*, 447–482

- Contreras, J. R., 87–92  
 Cooperative breeding, 417–421  
*Coragyps atratus*, 71–85, 173–200, 381–398, 447–482, 501–512, S83–S99  
*Corapipo alera*, 447–482  
*Coruja suindara*, 65–70  
*Coryphospingus cucullatus*, 381–398  
*Corythopsis torquata*, 173–200  
*Coscoroba coscoroba*, 513–525  
 Costa Rica, 201–208, 433–445, S267–S278, S279–S283  
 Costs, S199–S204  
*Cotinga amabilis*, 447–482  
*Cotinga cayana*, 173–200  
*Cotinga cotinga*, 41–52, 173–200  
 Cotinga, 433–445  
*Coturnix japonica*, 201–208  
 Cowbirds, 331–347  
*Cranioleuca pyrrhophia*, 145–153, 381–398  
*Cranioleuca vulpina*, 71–85  
*Crax alector*, 173–200  
*Crax rubra*, 447–482  
*Creatus furcatus*, S61–S72  
 Crocodylia, S101–S116  
 Crops, 371–380  
*Crotophaga ani*, 71–85, 173–200, 381–398, S117–S126  
*Crotophaga sulcirostris*, 447–482  
*Crypturellus atrocapillus*, S317–S321  
*Crypturellus bartletti*, S317–S321, S323–S328  
*Crypturellus boucardi*, 447–482  
*Crypturellus casiquiare*, S285–S291  
*Crypturellus cinereus*, 173–200, S317–S321  
*Crypturellus erythropus*, 173–200  
*Crypturellus erythropus/undulatus*, 173–200  
*Crypturellus soui*, 173–200, 447–482, S317–S321, S323–S328  
*Crypturellus tataupa*, 381–398  
*Crypturellus undulatus*, 173–200, S317–S321, S323–S328, S329–S337  
*Crypturellus variegatus*, 173–200, S317–S321, S323–S328  
*Crypturellus cinereus*, S323–S328  
*Cyanerpes caeruleus*, 41–52, 173–200  
*Cyanerpes cyaneus*, 173–200, 447–482  
*Cyanerpes lucidus*, 447–482  
*Cyanerpes nitidus*, 41–52, 173–200  
*Cyanicterus cyanicterus*, 173–200  
*Cyanocompsa brissonii*, 247–251, 381–398  
*Cyanocompsa cyanooides*, 173–200, 247–251, 447–482  
*Cyanocorax beecheii*, S175–S181  
*Cyanocorax cayanus*, 173–200  
*Cyanocorax chrysops*, 381–398  
*Cyanocorax cristatellus*, S117–S126  
*Cyanocorax cyanomelas*, 381–398  
*Cyanocorax dickeyi*, S175–S181  
*Cyanocorax morio*, 447–482  
*Cyanolyca mirabilis*, S175–S181  
*Cyanolyca nana*, S175–S181  
*Cyanolyca turcosa*, 483–492  
*Cyclarhis gujanensis*, 173–200, 381–398, S117–S126  
*Cygnus melancoryphus*, 513–525  
*Cymbilaimus lineatus*, 173–200, 447–482  
*Cyphorhinus arada*, 173–200  
*Cyphorhinus phaeocephalus*, 447–482  
*Cypseloides cryptus*, 447–482  
*Cypsnagra hirundinacea*, 173–200  
 D'Amico, V. L., 357–364  
*Dacnis cayana*, 173–200, 447–482, S117–S126  
*Dacnis lineata*, 173–200  
 Daily mortality rate, 201–208  
*Daption capense*, 163–171  
*Daptrius americanus*, S83–S99  
*Daptrius ater*, 173–200, S83–S99  
*Darina solenoides*, 357–364  
 Darrieu, C. A., 299–306  
 Declination, S357–S365  
*Deconychura longicauda*, 447–482  
 Delgado-V., C. A., 413–415  
*Dendrocincla anabatina*, 447–482  
*Dendrocincla fuliginosa*, 173–200, 447–482  
*Dendrocincla homochroa*, 447–482  
*Dendrocolaptes certhia*, 173–200  
*Dendrocolaptes picumnus*, 173–200, 539–542  
*Dendrocolaptes sanctithomae*, 447–482  
*Dendrocygna autumnalis*, 447–482  
*Dendrocygna bicolor*, 381–398, 447–482  
*Dendrocygna viduata*, 71–85, 173–200



- Dendroica caerulescens*, 447–482  
*Dendroica castanea*, 447–482  
*Dendroica cerulea*, 447–482  
*Dendroica coronata*, 447–482  
*Dendroica discolor*, 447–482  
*Dendroica dominica*, 447–482  
*Dendroica fusca*, 447–482  
*Dendroica graciae*, 447–482  
*Dendroica magnolia*, 447–482  
*Dendroica palmarum*, 447–482  
*Dendroica pensylvanica*, 447–482  
*Dendroica petechia*, 173–200, 447–482  
*Dendroica striata*, 41–52  
*Dendroica tigrina*, 447–482  
*Dendroica virens*, 447–482  
*Dendrotyx barbatus*, S175–S181  
*Derophtus accipitrinus*, 173–200  
 DeVries, P. J., 403–406  
 Di Giacomo, A. G., S145–S157  
 Di Giacomo, A. S., S145–S157  
 Diaz, M., S215–S222  
 Diet, 53–64, 87–92, 103–110, 121–131, 299–306, 307–311, 413–415, 423–427, 493–500  
*Diglossa albilatera*, 483–492  
*Diglossa humeralis*, 483–492  
*Diglossa lafresnayii*, 483–492  
*Diglossopsis cyanea*, 483–492  
 Dimorphism, 271–278  
 Dinosauria, S101–S116  
*Diomedea irrorata*, S61–S72  
*Diopsittaca nobilis*, 173–200  
 Distribution, 71–85, 111–119, 139–141, 313–317, S159–S167  
*Diuca diuca*, 513–525  
 Diurnal foraging, 407–411  
 Diversity, 447–482  
*Dives dives*, 447–482  
 DNA fingerprinting, 417–421  
 Dohen, E. B., 201–208  
*Dolospingus fringilloides*, 173–200  
*Donacobius atricapillus*, 71–85  
*Doricha eliza*, S175–S181  
 Dry forest, 111–119  
*Dryocopus lineatus*, 173–200, 381–398, 447–482  
*Dumetella carolinensis*, 447–482  
 Durães, R., S127–S135  
*Dysithamnus mentalis*, 71–85  
*Dysithamnus occidentalis*, 349–356  
*Dysithamnus striaticeps*, 447–482  
 Ecological indicator, 501–512  
 Ecology, 31–40  
 Ectoparasite, 247–251  
 Ecuador, 349–356, 365–370, 403–406, 483–492, 539–542  
 Effects, S61–S72  
 Egg description, 349–356, 403–406  
 Egg, 201–208, 219–224, 365–370  
*Egretta caerulea*, 173–200, 447–482  
*Egretta rufescens*, 447–482  
*Egretta thula*, 71–85, 173–200, 381–398, 447–482  
*Egretta tricolor*, 447–482, S223–S232  
 El Niño effects, S61–S72  
*Elaenia albiceps*, 381–398, 483–492  
*Elaenia chiriquensis*, 173–200  
*Elaenia cristata*, 173–200  
*Elaenia flavogaster*, 71–85, 173–200, 447–482, S117–S126  
*Elaenia pallatangae*, 483–492  
*Elaenia parvirostris*, 381–398, S349–S355  
*Elaenia ruficeps*, 173–200  
*Elaenia spectabilis*, 71–85, 381–398  
*Elaenia strepera*, S349–S355  
*Elanoides forficatus*, 31–40, 173–200, 447–482, S83–S99, S267–S278  
*Elanus leucurus*, 173–200, 447–482, S83–S99  
*Electron carinatum*, 447–482  
*Electron platyrhynchum*, 447–482  
 Electroretinogram, S215–S222, S223–S232, S251–S260  
 Elevational migration, 433–445  
 Emberizidae, 17–30, 145–153  
*Emberizoides herbicola*, 173–200  
 Emerald Toucanet, 433–445  
*Empidonax albigularis*, 447–482  
*Empidonax alhorum*, 447–482  
*Empidonax flaviventris*, 447–482  
*Empidonax minimus*, 447–482  
*Empidonax traillii*, 447–482  
*Empidonax virens*, 447–482

- Empidonomus aurantioatrocristatus*, 381–398  
*Empidonomus varius*, 41–52, 71–85, 145–153, 173–200, 381–398, S117–S126  
 Endangered, 133–137  
 Endangered birds, S175–S181  
 Energetic offer, S205–S213  
*Ensifera ensifera*, S205–S213  
 Environmental ethics, S419–S434, S435–S444  
*Eremobius phoenicurus*, 513–525  
*Eriocnemis derbyi*, S205–S213  
*Eriocnemis luciani*, 483–492  
*Eriocnemis mosquera*, S205–S213  
 Escalante, 1–3, S175–S181  
 Espinosa G., L. A., S1–S34  
 Ethnohisrory  
 Ethno-ornithology, S395–S407, S409–S417, S419–S434  
*Eubucco versicolor*, 283–284  
*Eucometis penicillata*, 247–251, 447–482  
*Endocimus albus*, 447–482  
*Endocimus ruber*, S223–S232  
*Endromia elegans*, 513–525  
*Euparkeria*, S101–S116  
*Eupetomena macroura*, 173–200  
*Eupherusa poliocerca*, S175–S181  
*Euphonia affinis*, 447–482  
*Euphonia chlorotica*, 71–85, 173–200, 381–398  
*Euphonia chrysopasta*, 173–200  
*Euphonia cyanocephala*, 483–492  
*Euphonia finschi*, 173–200  
*Euphonia gouldi*, 447–482  
*Euphonia hirundinacea*, 447–482  
*Euphonia luteicapilla*, 447–482  
*Euphonia minuta*, 447–482  
*Euphonia rufiventris*, 41–52  
*Euphonia violacea*, 71–85, 173–200  
*Eurypyga helias*, 173–200, 447–482  
*Euscarthmus rufomarginatus*, 173–200  
 Evolution, S101–S116, S191–S198  
 Excavator, S455–S465  
 Extinction, S145–S157  
 Eye size, S243–S250  
*Falco columbarius*, 173–200, 447–482, S83–S99, S267–S278  
*Falco deiroleucus*, 173–200, 447–482, S83–S99  
*Falco femoralis*, 173–200, 447–482, 483–492, 501–512, S83–S99  
*Falco mexicanus*, S83–S99  
*Falco peregrinus anatum*, 501–512  
*Falco peregrinus*, 173–200, 447–482, S83–S99  
*Falco rufigularis*, 173–200, 381–398, 447–482, 501–512, S83–S99  
*Falco sparverius*, 173–200, 381–398, 447–482, 483–492, 513–525, S83–S99, S267–S278  
 Falconiformes, S83–S99, S261–S266  
 Falcons, S477–S483  
 Farmer, A., S377–S384  
 Faucett, R. C., 381–398  
 Favero, M., 121–131, 493–500  
 Fecal samples, 53–64  
 Feeding, 93–102  
 Feeding strategies, 357–364, S223–S232  
 Fernández, G., S385–S394  
 Fernández, M. J., S215–S222  
 Fernández, M., S377–S384  
 Ferrari, S., 103–110  
 Ferretti, V., 299–306  
 Field edges, 371–380  
 Finch, D. W., 173–200  
 Fish prey, 121–131  
 Fishery waste site, 93–102  
 Fledging, 365–370, S455–S465  
 Flock size, 407–411  
 Flores, C., S339–S347  
*Florisuga mellivora*, 173–200, 447–482  
 Flower mites, S199–S204  
*Fluvicola pica*, 71–85, 381–398  
 F-number, S243–S250  
 Fonseca, F. L. de S., 133–137  
 Fonseca, V. S. da S., 133–137  
 Food habits, 87–92, 279–282  
 Food resources, S61–S72  
 Foraging, 283–284  
 Foraging behavior, 5–15  
 Foraging methods, 357–364  
 Foraging strategies, 357–364  
 Forest fragment isolation, S137–S144  
 Forest fragment use, S137–S144  
 Forest fragmentation, S117–S126  
 Forested habitat, 307–311

- Formicariidae, 225–235, S127–S135  
*Formicarius analis*, 173–200, 447–482  
*Formicarius colma*, 173–200  
*Formicivora grisea*, 173–200  
*Formicivora melanogaster*, 381–398  
*Formicivora rufa*, 173–200  
*Forpus xanthopterygius*, 71–85  
 Fossil birds, S73–S81  
 Fraga, R. M., 139–141  
 François Vuilleumier Fund, 576  
*Frederickena viridis*, 173–200  
*Fregata ariel*, S61–S72  
*Fregata magnificens*, 447–482, S61–S72  
*Fregata minor*, S61–S72  
*Fregetta tropica*, 163–171  
 Fritz, J. S., 201–208  
 Frost, M., 429–432  
 Frugivory, S117–S126  
*Fulica americana*, 447–482  
*Fulica leucoptera*, 381–398, 513–525  
*Fulmarus glacialisoides*, 163–171  
 Fungal pathogens, S199–S204  
 Funk, W. C., 365–370  
 Furnariidae, 145–153  
*Furnarius leucopus*, 173–200  
*Furnarius rufus*, 71–85, 271–278, 371–380, 381–398  
*Galbalcyrhynchus purusianus*, 319–330  
*Galbula albirostris*, 173–200  
*Galbula cyanescens*, 319–330  
*Galbula dea*, 173–200  
*Galbula galbula*, 173–200  
*Galbula leucogastra*, 41–52, 173–200  
*Galbula ruficauda*, 71–85, 447–482  
 Galliformes, S301–S307  
*Gallinago delicata*, 447–482  
*Gallinago gallinago*, 513–525  
*Gallinago paraguayana*, 173–200  
*Gallinago undulata*, 173–200  
*Gallinula chloropus*, 381–398, 447–482  
 Gamebirds, S301–S307, S323–S328  
*Gampsonyx swainsonii*, 173–200, S83–S99  
 Gandini, P., 103–110  
 Garitano-Zavala, A., S339–S347  
 Gelis, R. A., 403–406  
 Genetic differentiation, S169–S174  
 Genetic distances, 527–537  
*Geositta antarctica*, 513–525  
*Geositta cunicularia*, 513–525  
*Geothlypis aequinoctialis*, 173–200, 381–398  
*Geothlypis flavovelata*, S175–S181  
*Geothlypis poliocephala*, 155–162, 447–482  
*Geothlypis semiflava*, 447–482  
*Geothlypis trichas*, 447–482  
*Geotrygon montana*, 173–200, 265–267, 447–482  
*Geotrygon violacea*, 447–482  
*Geranoaetus melanoleucus*, 501–512, 513–525, S83–S99  
*Geranoospiza caeruleascens*, 71–85, 173–200, 447–482, S83–S99  
 Gerhardt, R. P., S477–S483  
 Ghys, M. I., 493–500  
 Giaccardi, M., 93–102  
 Gill, V. A., S159–S167  
 Gimenes, M. R., 71–85  
 GIS, S127–S135  
 Gismondi, P., S339–S347  
*Glaucidium brasilianum*, 173–200, 381–398, 447–482, S477–S483  
*Glaucidium griseiceps*, 447–482  
*Glaucidium hardyi*, 173–200  
*Glaucidium jardi*, 483–492  
*Glaucidium sanchezi*, S175–S181  
*Glaucidium tucumanum*, 501–512  
*Glaucis aenea*, 447–482  
*Glycera americana*, 357–364  
*Glyphorhynchus spirurus*, 173–200, 447–482  
 Goes, R., 423–427  
 Goldfeder, S., 543–545  
 González, P. M., S357–S365  
 Goodship, N. M., S243–S250  
*Grallaria quitensis*, 483–492  
*Grallaria ruficapilla*, 483–492  
*Grallaria rufula*, 483–492  
*Grallaria squamigera*, 483–492  
*Grallaricula loricata*, 225–235  
*Grallaricula peruviana*, 403–406  
*Granatellus pelzelni*, 173–200  
 Grasslands, S301–S307  
 Grassland shorebirds, S159–S167

- Grau, E. T., 527–537  
 Grazing, S301–S307, S309–S315  
 Greeney, H. F., 349–356, 365–370, 403–406, 539–542  
*Griseotyrannus aurantioatrocristatus*, 41–52, 145–153  
 Guatemala, S477–S483  
 Guayana, 173–200  
 Guerschman, J. P., S367–S376  
*Guira guira*, 71–85, 381–398, S117–S126  
 Gulf of California, 237–246  
 Gutiérrez Z., A., S205–S213  
*Gygis alba*, S61–S72  
*Gymnocichla nudiceps*, 447–482  
*Gymnoderus foetidus*, 173–200, 539–542  
*Gymnopithys leucaspis*, 447–482  
*Gymnopithys rufigula*, 173–200  
*Habia fuscicauda*, 53–64, 447–482  
*Habia rubica*, 53–64, 447–482  
 Habitat, 111–119, S145–S157  
 Habitat conservation, S445–S454  
 Habitat heterogeneity, S127–S135  
 Habitat model, S367–S376  
 Habitat partition, 5–15  
 Habitat selection, S317–S321  
 Habitat use, S117–S126, S159–S167  
*Haematoderus militaris*, 173–200  
*Haematopus leucopodus*, 513–525  
*Haematopus palliatus*, 269–270, 447–482  
*Haliaeetus leucocephalus*, S83–S99  
*Halobaena caerulea*, 163–171  
 Hannelly, E. C., 403–406, 539–542  
*Hapaloptila castanea*, 539–542  
*Harpagus bidentatus*, 447–482, S83–S99  
*Harpagus diodon*, S83–S99  
*Harpia harpyja*, 447–482, S83–S99  
*Harpophalietus coronatus*, S83–S99  
*Harpophalietus solitarius*, 31–40, S83–S99  
 Hartz, S. M., 5–15., 279–282  
*Heliactin bilopha*, 173–200  
*Heliangelus viola*, 483–492  
*Heliomaster longirostris*, 173–200, 447–482  
*Heliornis fulica*, 447–482  
*Heliobryx aurita*, 173–200  
*Heliobryx barroti*, 447–482  
*Hellmayrea gularis*, 483–492  
*Helmitberos vermivora*, 447–482  
*Hemispingus superciliosus*, 483–492  
*Hemithraupis flavicollis*, 41–52  
*Hemithraupis guira*, 173–200, 381–398  
*Hemitriccus josephinae*, 173–200  
*Hemitriccus margaritaceiventer*, 381–398  
*Henicorbina leucosticta*, 173–200, 447–482  
 Hennessey, A. B., 283–284  
 Henri Pittier National Park, 225–235  
 Hernández, M. de los A., 357–364  
*Herpetotheres cachinnans*, 173–200, 381–398, 447–482, S83–S99  
*Herpsilochmus atricapillus*, 381–398  
*Herpsilochmus longirostris*, 71–85  
*Herpsilochmus rufomarginatus*, 173–200  
*Herpsilochmus stictocephalus*, 173–200  
*Herpsilochmus sticturus*, 173–200  
 Herrera, G., S215–S222  
*Hersilochmus dorsimaculatus*, 41–52  
*Heteronetta atricapilla*, 381–398  
 High elevations, S191–S198  
 High-Andean birds, S205–S213  
*Himantopus himantopus*, 71–85  
*Himantopus mexicanus*, 381–398, 447–482, S223–S232  
*Hirundinea ferruginea*, 71–85, 173–200  
*Hirundo rustica*, 173–200, 447–482  
 Höfling, E., 527–537, S73–S81  
 Hole-nesting, S455–S465  
 Honduras, 447–482  
*Hoploxypterus cayanus*, 173–200  
 Host, 247–251  
 Huallpara, S., S339–S347  
 Hummingbirds, S191–S198, S205–S213, S215–S222, S409–S417  
 Hummingbird-plant communities, S183–S190  
 Hummingbird-plant interaction, S199–S204  
 Hutt, M. B., 429–432  
*Hydropsalis torquata*, 173–200  
*Hylocharis chrysura*, 71–85, 381–398  
*Hylocharis cyanus*, 173–200  
*Hylocharis eliciae*, 447–482  
*Hylocharis sapphirina*, 173–200  
*Hylocichla mustelina*, 447–482

- Hyloctistes subulatus*, 447–482  
*Hylomanes momotula*, 447–482  
*Hylopezus dives*, 447–482  
*Hylopezus macularius*, 173–200  
*Hylopezus perspicillatus*, 447–482  
*Hylophilus aurantifrons*, 247–251  
*Hylophilus decurtatus*, 447–482  
*Hylophilus muscicapinus*, 173–200  
*Hylophilus ochraceiceps*, 447–482  
*Hylophilus pectoralis*, 173–200  
*Hylophilus thoracicus*, 173–200  
*Hylophylax naevia*, 173–200  
*Hylophylax naevioides*, 447–482  
*Hylophylax poecilinota*, 173–200, 265–267  
*Hylorchilus navai*, S175–S181  
*Hymenops perspicillata*, 381–398  
*Hypocnemis cantator*, 173–200  
Hypothesis testing, S41–S60  
*Ibyster americanus*, 173–200, 447–482  
*Icteria virens*, 447–482  
*Icterus bonana*, 331–347  
*Icterus cayanensis*, 71–85, 173–200, 331–347, 381–398  
*Icterus chrysater*, 331–347, 447–482  
*Icterus chrysocephalus*, 173–200, 331–347  
*Icterus cucullatus*, 331–347  
*Icterus dominicensis*, 331–347  
*Icterus fuertesi*, 331–347  
*Icterus galbula*, 331–347, 447–482  
*Icterus graccannae*, 331–347  
*Icterus graduacauda*, 331–347  
*Icterus gularis*, 331–347  
*Icterus icterus*, 331–347  
*Icterus jamacaii*, 331–347  
*Icterus leucopteryx*, 331–347  
*Icterus mesomelas*, 447–482  
*Icterus nigrogularis*, 173–200, 331–347  
*Icterus parisorum*, 331–347  
*Icterus pectoralis*, 331–347, 447–482  
*Icterus prosthemelas*, 447–482  
*Icterus pustulatus*, 331–347  
*Icterus spurious*, 331–347, 447–482  
*Icterus wagleri*, 331–347  
*Icterus*, 331–347  
*Ictinia mississippiensis*, 447–482, S83–S99, S261–S266, S267–S278  
*Ictinia plumbea*, 173–200, 447–482, S83–S99, S267–S278  
Incubation, 155–162, 349–356, 365–370, 539–542  
Index to volume 15, 549–574  
*Inezia caudata*, 173–200  
*Inezia inornata*, 381–398  
Infrared, S455–S465  
Insects, 121–131, 299–306  
Instrumental value, S435–S444  
Intrinsic value, S435–S444  
Inventory, 483–492  
*Iodopleura isabellae*, 41–52  
Isacch, J. P., S159–S167  
Islands, 71–85  
*Ixobrychus exilis*, 173–200  
*Ixobryncbus exilis*, 447–482  
Ixodoidea ectoparasite, 247–251  
*Jabiru mycteria*, 71–85, 173–200, 447–482  
*Jacamerops aureus*, 173–200, 319–330  
*Jacana jacana*, 71–85, 173–200, 381–398  
*Jacana spinosa*, 447–482  
Jaksic, F., S61–S72  
Jurado, A., 139–141  
Kelly, D. J., S243–S250  
Kelp Gull, 93–102  
Kester, C., S377–S384  
Kimura, R. K., 527–537  
Kindel, A., 5–15  
Kirwan, G. M., 257–263  
*Klais guimeti*, 447–482  
*Knipolegus aterrimus*, 381–398  
Kopij, G., 271–278  
Krabbe, N., 365–370  
Kusch, A., 313–317  
La Niña effects, S61–S72  
Lafkenche, S395–S407  
*Lafresnaya lafresnayi*, 483–492, S205–S213  
Laguna Kaucaya, 381–398  
*Lampornis amethystinus*, S199–S204  
*Lamprospiza melanoleuca*, 173–200  
Lanctot, R. B., S159–S167, S367–S376  
Land birds, 513–525, S61–S72  
*Lanio fulvus*, 173–200

- Lanio leucothorax*, 447–482  
*Laniocera hypopyrra*, 173–200  
*Laniocera rufescens*, 447–482  
Lank, D. B., S385–S394  
Lara, C., S199–S204  
Large frugivorous birds, 433–445  
Large fruit-eating birds, 433–445  
*Larosterna inca*, S61–S72  
*Larus argentatus*, 447–482  
*Larus atricilla*, 447–482  
*Larus dominicanus*, 93–102, 163–171, 513–525  
*Larus fuliginosus*, S61–S72  
*Larus heermanni*, 237–246  
*Larus livens*, 237–246  
*Larus maculipennis*, 493–500  
*Laterallus albigularis*, 447–482  
*Laterallus exilis*, 447–482  
*Laterallus melanophaius*, 381–398  
*Laterallus ruber*, 447–482  
*Lathrotriccus euleri*, 173–200, 381–398  
Latitudinal dependent, S385–S394  
Lebbin, D. J., 139–141  
*Legatus leucophaius*, 41–52, 173–200, 447–482  
*Lepidocolaptes albolineatus*, 173–200  
*Lepidocolaptes angustirostris*, 173–200, 381–398  
*Lepidocolaptes souleyetii*, 447–482  
*Lepidothrix serena*, 173–200  
*Leptasthenura aegithaloïdes*, 513–525  
*Leptasthenura platensis*, 145–153  
*Leptodon cayanensis*, 31–40, 173–200, 447–482, S83–S99  
*Leptodon forbesi*, S83–S99  
*Leptopogon amaurocephalus*, 447–482  
*Leptosittaca branickii*, 483–492  
*Leptotila cassini*, 447–482  
*Leptotila rufaxilla*, 173–200, 447–482  
*Leptotila verreauxi*, 71–85, 173–200, 381–398, 483–492, S117–S126  
*Lesbia nuna*, 483–492  
*Lesbia victoriae*, S191–S198, S205–S213  
*Lessonia rufa*, 513–525  
*Leucopternis albicollis*, 173–200, 447–482, S83–S99  
*Leucopternis kuhli*, S83–S99  
*Leucopternis lacernulata*, S83–S99  
*Leucopternis melanops*, S83–S99  
*Leucopternis occidentalis*, S83–S99  
*Leucopternis plumbea*, S83–S99  
*Leucopternis polionota*, S83–S99  
*Leucopternis princeps*, S83–S99  
*Leucopternis schistacea*, S83–S99  
*Leucopternis semiplumbea*, 447–482, S83–S99  
Leveau, C. M., 371–380  
Leveau, L. M., 371–380  
Life-history strategies, S385–S394  
*Limnodromus griseus*, 447–482, S223–S232  
Lindell, C. A., 201–208  
*Lipaugus unirufus*, 447–482  
*Lipaugus vociferans*, 173–200  
*Lobipes lobatus*, S61–S72  
Loiselle, B. A., S127–S135  
*Longisquama*, S101–S116  
Lopes, E. V., S137–S144  
Lopes, I. F., S169–S174  
Lopes, L. E., 423–427  
*Lophornis ornatus*, 173–200  
*Lophotrix cristata*, 173–200, 447–482  
*Lophotriccus galeatus*, 173–200  
*Lophotriccus pilaris*, 173–200  
*Lophotriccus pileatus*, 447–482  
*Lophotriccus vitiensis*, 173–200  
*Lophornis belenae*, 447–482  
Los Tuxtlas, 53–64  
*Loxia curvirostra*, 447–482  
Lozano, E., 237–246  
Lozano, J. C., S339–S347  
Luquillo Mountains, 289–297  
*Lurocalis semitorquatus*, 447–482  
Lysinger, M., 365–370  
Macaws, S279–S283  
*Machetornis rixosus*, 71–85, 381–398  
*Macronectes giganteus*, 163–171  
Maillard Z., O., 381–398, 399–402  
*Malacoptila fusca*, 173–200  
*Malacoptila panamensis*, 447–482  
Maldonado-Coelho, M., 547–548  
Male parental care, S329–S337  
Mamani, A. M., 381–398  
*Manacus candei*, 447–482  
*Manacus manacus*, 173–200

- Management, S279–S283, S455–S465  
 Maniraptora, S101–S116  
 Manu Biosphere Reserve, 31–40  
 Mapuche, S395–S407, S419–S434, S435–S444  
 Mar Chiquita Lagoon, 121–131  
 Maragliano, R., 271–278  
 Margarita Island, 253–256  
*Margarops fuscatus*, S455–S465  
*Margarornis squamiger*, 483–492  
 Marín, G., S223–S232, S251–S260  
 Martin, G. R., S233–S242  
 Martínez, W. E., 155–162  
 Massardo, F., S395–S407  
 Mauco, L., 121–131  
 McCracken, K. G., 543–545  
 McNeil, R., 407–411, S223–S232, S233–S242, S251–S260  
*Mecocerculus leucophrys*, 483–492  
*Mecocerculus stictopterus*, 483–492  
*Megalancosaurus*, S101–S116  
*Megaptera novaeangliae*, 163–171  
*Megarhynchus pitangua*, 173–200, 447–482  
*Megascops choliba*, 381–398  
*Megascops guatemalae*, 447–482  
*Melanerpes candidus*, 71–85, 173–200, 381–398, S117–S126  
*Melanerpes cruentatus*, 173–200  
*Melanerpes flavifrons*, 71–85  
*Melanerpes formicivorus*, 447–482  
*Melanerpes pucherani*, 447–482  
 Melo Ferreira, R. de, 423–427  
 Mendes de Carvalho Filho, E. P., 219–224  
 Mendonça-Lima, A. de, 5–15  
*Mesembrinibis cayennensis*, 71–85, 173–200, 447–482  
*Metallura tyriantbina*, 483–492, S205–S213  
 Metaphors, S435–S444  
 México, 53–64, 237–246, S175–S181  
 Mezquida, E. T., 145–153  
*Micrastur buckleyi*, S83–S99  
*Micrastur gilvicolis*, 173–200, S83–S99  
*Micrastur mirandollei*, S83–S99  
*Micrastur plumbeus*, S83–S99  
*Micrastur ruficollis*, 173–200, 447–482, S83–S99  
*Micrastur semitorquatus*, 173–200, 447–482, S83–S99  
*Microbates cinereiventris*, 447–482  
*Microbates collaris*, 173–200  
*Microcerculus bambla*, 173–200  
*Microcerculus philomela*, 447–482  
*Microchera albocoronata*, 447–482  
 Microclimate, 145–153  
 Microhabitat selection, 145–153  
*Microphygia schomburgkii*, 173–200  
*Microbopias quixensis*, 447–482  
 Microsatellites, S169–S174  
 Migration, 41–52, S83–S99, S267–S278, S377–S384  
*Milvago chimachima*, 71–85, 173–200, S83–S99  
*Milvago chimango*, 371–380, S83–S99  
*Mimus gilvus*, 173–200, 417–421  
*Mimus patagonicus*, 513–525  
*Mimus saturninus*, 71–85, 173–200, 417–421, S117–S126  
*Mimus triurus*, 381–398  
*Mionectes oleagineus*, 173–200, 447–482  
 Mitochondrial DNA, S169–S174  
*Mniotilta varia*, 447–482  
 Molecular, S293–S299  
 Molina, M., S339–S347  
*Molothrus aeneus*, 331–347, 447–482  
*Molothrus ater*, 331–347  
*Molothrus bonariensis*, 71–85, 173–200, 271–278, 331–347, 381–398  
*Molothrus oryzivorus*, 173–200, 381–398, 447–482  
 Molt, S205–S213  
*Momotus momota*, 173–200, 381–398, 447–482  
*Monasa atra*, 173–200  
*Monasa morphoeus*, 447–482  
 Monitoring, S349–S355, S357–S365, S455–S465  
 Montalti, D., 271–278  
 Monte desert, 145–153  
 Monteverde, 433–445  
*Morphnus guianensis*, 447–482  
*Morphnus guianensis*, S83–S99  
 Morphology, S191–S198, S285–S291, S293–S299  
 Morphometrics, 271–278

- Morrison, R. I., S357–S365  
Morton, E. S., 417–421  
Mosa, S. G., S309–S315  
Moskitia, 447–482  
Mouse-colored Antshrike, 265–267  
*Mus musculus*, 65–70  
*Muscisaxicola macloviana*, 513–525  
*Mycteria americana*, 71–85, 173–200, 447–482, S169–S174  
*Myiarchus crinitus*, 447–482  
*Myiarchus ferox*, 173–200, S117–S126  
*Myiarchus swainsoni*, 173–200, 381–398  
*Myiarchus tuberculifer*, 173–200, 447–482  
*Myiarchus tyrannulus*, 173–200, 381–398, 447–482  
*Myiobius barbatus*, 173–200  
*Myiobius sulphureipygius*, 447–482  
*Myioborus melanocephalus*, 483–492  
*Myiodynastes luteiventris*, 447–482  
*Myiodynastes maculatus*, 41–52, 71–85, 173–200, 447–482, S117–S126  
*Myiopagis caniceps*, 41–52  
*Myiopagis gaimardii*, 41–52, 173–200  
*Myiopagis viridicata*, 173–200, 381–398, 447–482  
*Myiophobus fasciatus*, 173–200, 381–398  
*Myiopsitta monachus*, 209–218, 371–380  
*Myiornis ecaudatus*, 173–200  
*Myiotheretes fumigatus*, 483–492  
*Myiotheretes striaticollis*, 483–492  
*Myiozetetes cayanensis*, 173–200  
*Myiozetetes granadensis*, 447–482  
*Myiozetetes similis*, 71–85, 447–482, S117–S126  
*Myornis senilis*, 483–492  
*Myrmeciza excelsa*, 447–482  
*Myrmeciza ferruginea*, 173–200  
*Myrmeciza longipes*, 173–200  
*Myrmoborus leucophrys*, 173–200  
*Myrmorchilus strigilatus*, 381–398  
*Myrmornis torquata*, 173–200, 447–482  
*Myrmothera campanisona*, 173–200  
*Myrmotherula axillaris*, 173–200, 447–482  
*Myrmotherula brachyura*, 173–200  
*Myrmotherula fulviventris*, 447–482  
*Myrmotherula guttata*, 173–200  
*Myrmotherula gutturalis*, 173–200  
*Myrmotherula longipennis*, 173–200  
*Myrmotherula menetriesii*, 173–200  
*Myrmotherula schisticolor*, 447–482  
*Nannopterum harrisi*, S61–S72  
Nassif Del Lama, S., S169–S174  
Natural history, 365–370  
Nearctic-Neotropical migration, S41–S60  
Nectar thieves, S199–S204  
Negro, J. J., 513–525  
*Nemosia pileata*, 71–85, 173–200, 381–398  
*Neomorphus geoffroyi*, 447–482  
*Neopelma aurifrons*, 547–548  
*Neopelma pallescens*, 173–200  
*Neotbraupis fasciata*, 173–200  
Neotropical birds, S349–S355  
Neotropical gamebirds, S323–S328  
Neotropical Ornithological Society, S1–S34, S35–S40  
Neotropical Region, S41–S60  
Neotropical rodents, 413–415  
Neotropics, S83–S99, S317–S321  
*Neoxolmis rubetra*, 513–525  
*Neoxolmis rufiventris*, 513–525  
Néquiz, V., S175–S181  
*Nesofregatta albigularis*, S61–S72  
Nest, 219–224, 257–263, 265–267, 283–284, 365–370, 423–427, 539–542  
Nest adoption, 209–218  
Nest architecture, 257–263  
Nest excavation, 319–330  
Nest management, S455–S465  
Nest orientation, 145–153  
Nest site, 319–330, 349–356  
Nest success, 201–208  
Nesting, 269–270, S455–S465  
Nestling, 155–162, 219–224, 365–370  
Neto, S. D'A., 547–548  
New records, 483–492  
News, 575–576  
Niche, 5–15  
Nocturnal foraging, 407–411  
Nocturnal vision, S223–S232  
*Nomonyx dominicus*, 381–398  
Non-breeding season, S385–S394



- Notharchus macrorhynchos*, 41–52, 173–200, 447–482  
*Notharchus tectus*, 41–52, 173–200  
*Nothocercus bonapartei*, S285–S291  
*Nothoprocta ornata*, S339–S347  
*Nothoprocta pentlandii*, 111–119, S309–S315  
*Nothura darwini*, S309–S315  
*Nothura maculosa*, 371–380  
*Notiochelidon cyanoleuca*, 483–492, 513–525  
*Notiochelidon murina*, 483–492  
 Novak, S. J., 447–482  
*Numenius boreales*, S159–S167  
*Numenius phaeopus*, 447–482  
*Nyctanassa violaceus*, 447–482  
*Nyctibius grandis*, 173–200, 447–482  
*Nyctibius griseus*, 173–200, 381–398, 447–482  
*Nycticorax nycticorax*, 71–85, 173–200, 381–398, 447–482  
*Nycticorax violaceus*, S223–S232  
*Nyctidromus albicollis*, 71–85, 173–200, 447–482, S233–S242, S251–S260  
*Nyctiphrynus ocellatus*, 447–482  
*Nystalus maculatus*, 381–398  
 O'Hara, P. D., S385–S394  
 Obituaries, 429–432  
*Oceanites gracilis*, S61–S72  
*Oceanites oceanicus*, 163–171  
*Oceanites tethys*, S61–S72  
*Oceanodroma castro*, S61–S72  
*Oceanodroma macrodactyla*, S175–S181  
*Ochthoeca frontalis*, 483–492  
*Ochthoeca rufipectoralis*, 483–492  
 Ocmin-Petit, A., S323–S328  
*Odontophorus gujanensis*, 173–200  
*Odontophorus melanotis*, 447–482  
 Oesterheld, M., S367–S376  
 Ojeda, V., 307–311  
 Olivo, C., S261–S266  
*Oncostoma cinereigulare*, 447–482  
*Onychorhynchus coronatus*, 173–200, 447–482  
 Open cup, S455–S465  
 Open nesting, S455–S465  
*Oporornis formosus*, 447–482  
*Oporornis philadelphia*, 447–482  
*Oreopholus ruficollis*, 513–525, S159–S167  
*Oreotrochilus chimborazo*, 483–492  
*Oreotrochilus leucopterus*, S215–S222  
 Organizing committees, S1–S34  
 Orgeira, J. L., 163–171  
 Orioles, 331–347  
*Ornithion inermis*, 173–200  
*Ornithion semiflavum*, 447–482  
 Ornithischia, S101–S116  
 Ornithology, S41–S60  
*Oroaetus isidori*, 31–40, S83–S99  
*Ortalis canicollis*, 381–398  
*Ortalis cinereiceps*, 447–482  
*Ortalis motmot*, 173–200  
*Orthopsittaca manilata*, 173–200  
*Oryzoborus angolensis*, 173–200  
*Oryzoborus crassirostris*, 173–200  
*Oryzoborus funereus*, 447–482  
 Osborn, S., 31–40  
*Otus albogularis*, 483–492  
*Otus choliba*, 173–200, 501–512  
*Otus guatemalae*, S477–S483  
*Otus watsonii*, 173–200  
 Outbound migration, S261–S266  
 Oviraptoridae, S101–S116  
 Owls, S477–S483  
*Oxyura dominica*, 173–200  
*Oxyura jamaicensis*, 381–398  
*Pachyptila desolata*, 163–171  
*Pachyramphus aglaiae*, 447–482  
*Pachyramphus cinnamomeus*, 447–482  
*Pachyramphus marginatus*, 41–52, 173–200  
*Pachyramphus minor*, 173–200  
*Pachyramphus polychopterus*, 173–200, 381–398, 447–482  
*Pachyramphus rufus*, 173–200  
*Pachyramphus validus*, S117–S126  
 Página de la Presidenta, 1–3  
*Pagodroma nivea*, 163–171  
 Palacios, A., S215–S222  
 Panama, 417–421  
*Pandion haliaetus*, 173–200, 381–398, 447–482, S83–S99, S267–S278  
 Pando-Vasquez, L., S323–S328  
*Panyptila cayennensis*, 173–200, 265–267, 447–482

- Parabaño, 399–402  
*Parabuteo unicinctus*, 381–398, S83–S99  
 Paraguana Peninsula, 269–270  
 Paraguay, 87–92, 139–141  
 Paraná river, 71–85  
*Paridirallus sanguinolentus*, 381–398  
*Paroaria capitata*, 71–85  
*Paroaria coronata*, 381–398  
*Paroaria gularis*, 173–200  
 Parque Nacional Médanos del Chaco, 87–92  
*Parula americana*, 447–482  
*Parula pitiayumi*, 173–200, 381–398  
*Passer domesticus*, 371–380, 513–525  
*Passerina caerulea*, 447–482  
*Passerina ciris*, 447–482  
*Passerina cyanea*, 447–482  
 Passerines, 145–153, 271–278  
 Pastures, 371–380  
*Patagioenas cayennensis*, 381–398, 447–482  
*Patagioenas flavirostris*, 447–482  
*Patagioenas nigrirostris*, 447–482  
*Patagioenas speciosa*, 447–482  
*Patagona gigas*, 483–492  
 Patagonia, 93–102, 103–110, 307–311, 313–317, 513–525  
 Patterns, S83–S99  
 Pedó, E., 279–282  
*Pelecanoides* sp., 163–171  
*Pelecanus occidentalis thagus*, 447–482, S61–S72, S61–S72  
*Penelope montagnii*, 483–492  
*Penelope purpurascens*, 433–445, 447–482  
*Penelope superciliaris*, S117–S126  
 Peninsula Valdés, 357–364  
*Pernostola rufifrons*, 173–200  
 Pereira, S. L., 527–537  
*Perissocephalus tricolor*, 173–200  
*Peroglossus*, 527–537  
 Perú, 31–40, 111–119, 319–330, S279–S283  
 Petra, M. V., 133–137  
*Petrochelidon pyrrhonota*, 447–482  
*Phacellodomus ruber*, 71–85  
*Phacellodomus rufifrons*, 381–398  
*Phaenostictus mcleanmani*, 447–482  
*Phaeochroa cuvieri*, 447–482  
*Phaeomyias murina*, 173–200, 381–398  
*Phaeoprogne tapera*, 71–85  
*Phaeothlypis flaveola*, 173–200  
*Phaeothlypis rivularis*, 173–200  
*Phaethon aethereus*, S61–S72  
*Phaethon rubricauda*, S61–S72  
*Phaethornis angusti*, 173–200  
*Phaethornis bourcieri*, 173–200, S183–S190  
*Phaethornis gyy*, S191–S198  
*Phaethornis longirostris*, 447–482, S191–S198  
*Phaethornis malaris*, S183–S190  
*Phaethornis pretrei*, 71–85  
*Phaethornis ruber*, 173–200  
*Phaethornis rupurumii*, 173–200  
*Phaethornis strigularis*, 447–482  
*Phaethornis superciliosus*, 173–200  
*Phaethornis surmatophorus*, S191–S198  
*Phaethornis yarucui*, S191–S198  
 Phaethornithinae, S183–S190  
*Phaetusa simplex*, 71–85  
*Phalacrocorax atriceps*, 103–110  
*Phalacrocorax bougainvilli*, S61–S72  
*Phalacrocorax bransfordensis*, 163–171  
*Phalacrocorax brasiliannus*, 71–85, 173–200, 381–398, 447–482  
*Phalacrocorax gaimardi*, S61–S72  
*Phalaropus tricolor*, 447–482  
*Phalcoboenus albogularis*, S83–S99  
*Phalcoboenus australis*, S83–S99  
*Phalcoboenus carunculatus*, 483–492, S83–S99  
*Phalcoboenus megalopterus*, S83–S99  
*Pharomachrus mocinno*, 433–445  
*Phencticus chrysogaster*, 483–492  
*Phencticus ludovicianus*, 447–482  
*Philohydor lictor*, 173–200  
 Philosophy of science, S41–S60  
*Philhydor erythrocerus*, 173–200  
*Philhydor pyrrhodes*, 173–200  
*Philhydor rufum*, 381–398  
*Phleocryptes melanops*, 381–398  
*Phoebetria palpebrata*, 163–171  
*Phoenicopterus chilensis*, 513–525  
*Phoenicopterus ruber*, 407–411  
*Phrygilus fruticeti*, 513–525  
*Phrygilus gayi*, 513–525

- Phrygilus unicolor*, 483–492  
 Phylogenetic constraints, S191–S198  
 Phylogenetic hypotheses, S73–S81  
 Phylogeny, 527–537  
 Physiographic structure, 111–119  
 Piaskowski, V. D., 155–162  
*Piaya cayana*, 173–200, 381–398  
*Piaya cayana*, 447–482  
*Picooides mixtus*, 381–398  
*Picooides scalaris*, 447–482  
*Piculus chrysochloros*, 173–200, 381–398  
*Piculus flavigula*, 173–200  
*Piculus rivoli*, 483–492  
*Piculus rubiginosus*, 447–482  
*Piculus simplex*, 447–482  
*Picumnus albosquamatus*, 71–85  
*Picumnus cirratus*, 71–85, 173–200  
*Picumnus olivaceus*, 447–482  
*Picumnus* sp., 381–398  
 Pigmentation, S285–S291, S293–S299  
*Pilberodius pileatus*, 173–200  
 Piñeiro, G., S367–S376  
*Pionites melanocephala*, 173–200  
*Pionopsitta caica*, 173–200  
*Pionopsitta haematotis*, 447–482  
*Pionus fuscus*, 173–200  
*Pionus maximiliani*, 71–85, 381–398  
*Pionus menstruus*, 173–200  
*Pionus senilis*, 447–482  
*Pionus seniloides*, 483–492  
*Pionus sordidus*, 483–492  
 Piper, W. H., 417–421  
*Pipile cumanensis*, 173–200  
*Pipra erythrocephala*, 173–200, 247–251  
*Pipra filicauda*, 247–251  
*Pipra mentalis*, 447–482  
*Pipra pipra*, 173–200  
*Pipreola riefferii*, 483–492  
*Piprites chloris*, 173–200  
*Piprites griseiceps*, 447–482  
*Piranga flava*, 173–200, 381–398, 447–482  
*Piranga olivacea*, 447–482  
*Piranga rubra*, 173–200, 447–482  
*Pitangus sulphuratus*, 71–85, 173–200, 271–278, 371–380, 381–398, 447–482, S117–S126  
*Pithys albifrons*, 173–200  
*Pitylus grossus*, 173–200  
 Pizo, M. A., S117–S126  
 Plant community, S205–S213  
*Platalea ajaja*, 71–85, 381–398, 447–482  
*Platyrinchus canrominus*, 447–482  
*Platyrinchus coronatus*, 447–482  
*Platyrinchus mystaceus*, 247–251  
*Platyrinchus saturatus*, 173–200  
*Plegadis chibi*, 381–398  
*Pluvialis dominica*, 173–200, 447–482, S159–S167  
*Pluvialis squatarola*, 447–482, S223–S232  
*Podager nacunda*, 173–200  
*Podilymbus podiceps*, 381–398, 447–482  
*Poecilatriccus sylvia*, 173–200, 447–482  
 Poems, S419–S434  
 Poetry, S419–S434  
 Pohl, N., S215–S222  
 Point counts, 513–525  
*Polioptila dumicola*, 271–278, 381–398  
*Polioptila guianensis*, 41–52  
*Polioptila plumbea*, 173–200, 447–482, 539–542  
 Pollen loads, S183–S190  
*Polyborus plancus*, 71–85, S83–S99  
 Polychaetes, 357–364  
*Polystictus pectoralis*, 173–200  
*Polytmus guainumbi*, 173–200  
*Polytmus theresiae*, 173–200  
*Poospiza melanolenca*, 381–398  
*Poospiza ornata*, 145–153  
*Poospiza torquata*, 145–153  
 Popper, S41–S60  
 Population, 289–297, S145–S157  
 Population modeling, 289–297  
 Population size, S357–S365, S367–S376  
*Porphyrio flavirostris*, 173–200  
*Porphyrio martinica*, 173–200, 381–398, 447–482  
*Porphyrospiza caerulescens*, 399–402  
 Porrás-Peñaranda, P., S267–S278  
 Port, J. L., 209–218  
*Porzana albicollis*, 173–200  
 Porzecanski, A. L., S293–S299  
 Post-breeding period, 371–380

- Povedano, H., 139–141  
 Predation, 65–70  
*Premnoplex brunnescens*, 483–492  
 Prevailing winds, 145–153  
 Proceedings of the VII<sup>th</sup> Neotropical Ornithological Congress, 575  
 Processes, S83–S99  
*Procnias alba*, 173–200  
*Procnias tricarunculata*, 433–445, 447–482  
*Progne chalybea*, 173–200, 447–482  
*Progne modesta*, 513–525  
*Progne subis*, 447–482  
*Progne tapera*, 173–200, 381–398  
*Propyrrhura auricollis*, 381–398  
*Protonotaria citrea*, 447–482  
*Psarocolius decumanus*, 173–200, 381–398  
*Psarocolius montezuma*, 447–482  
*Psarocolius viridis*, 173–200  
*Psarocolius wagleri*, 447–482  
*Pseudocolaptes boissonneautii*, 483–492  
*Pseudoleistes virescens*, 139–141, 271–278  
*Pseudoleites guiraburo*, 139–141, S117–S126  
*Pseudoscops clamator*, 381–398  
*Psophia crepitans*, 173–200  
*Pterocnemia pennata*, 513–525  
*Pterodroma alba*, S61–S72  
*Pterodroma brevirostris*, 163–171  
*Pterodroma macroptera*, 163–171  
*Pterodroma phaeopygia*, S61–S72  
*Pteroglossus aracari*, 173–200, 527–537  
*Pteroglossus beaubarnaesii*, 527–537  
*Pteroglossus bitorquatus*, 527–537  
*Pteroglossus castanotis*, 527–537  
*Pteroglossus inscriptus*, 527–537  
*Pteroglossus torquatus*, 447–482, 527–537  
*Pteroglossus viridis*, 173–200  
*Pterophanes cyanopterus*, 483–492, S205–S213  
 Puebla-Olivares, F., 53–64  
 Puerto Rico, 289–297  
*Puffinus griseus*, 163–171  
*Puffinus lherminieri*, S61–S72  
*Puffinus nativitatis*, S61–S72  
*Puffinus pacificus*, S61–S72  
*Pulsatrix perspicillata*, 173–200, 381–398, 447–482  
 Pupil aperture, S243–S250  
 Puyehue, S1–S34  
*Pygiptila stellaris*, 41–52  
*Pygoscelis adeliae*, 163–171  
*Pygoscelis antarctica*, 163–171  
*Pygoscelis papua*, 163–171  
*Pyrocephalus rubinus*, 173–200, 381–398, 447–482  
*Pyrrhura molinae*, 381–398  
*Querula purpurata*, 173–200  
 Quintana, R. D., 513–525  
*Quiscalus mexicanus*, 447–482  
 Rainforest birds, 41–52  
 Ramírez F., Y. M., S223–S232, S233–S242, S251–S260  
 Ramphastidae, 433–445, 527–537  
*Ramphastos sulfuratus*, 447–482  
*Ramphastos swainsonii*, 447–482  
*Ramphastos toco*, 381–398, 527–537  
*Ramphastos tucanus*, 173–200  
*Ramphastos vitellinus*, 173–200, 527–537  
*Ramphocaenus melanurus*, 173–200, 447–482  
*Ramphocelus carbo*, 71–85, 173–200, 247–251  
*Ramphocelus passerinii*, 447–482  
*Ramphocelus sanguinolentus*, 447–482  
*Ramphomicron microrhynchum*, S205–S213  
*Ramphotrigon ruficauda*, 173–200  
 Ramsar convention, S445–S454  
 Raptor, 501–512  
 Raptors, 31–40, S83–S99, S267–S278, S477–S483  
 Red list, S175–S181  
 Rejecters, 331–347  
 Relationships, 257–263  
 Reproduction, 155–162, 219–224, 237–246, S205–S213, S455–S465  
 Reproductive biology, 423–427  
 Reproductive success, 237–246, 253–256  
 Research on Neotropical birds, S41–S60  
 Resources selection, S205–S213  
 Retina, S223–S232, S251–S260  
*Rhinocrypta lanceolata*, 145–153  
 Rhinocryptidae, 145–153  
*Rhynchocyclus brevirostris*, 447–482  
*Rhynchocyclus fulvipectus*, 365–370

- Rhynchocyclus olivaceus*, 173–200, 247–251  
*Rhynchortyx cinctus*, 447–482  
*Rhynchotus rufescens*, S285–S291  
*Rhytipterna holerythra*, 447–482  
*Rhytipterna immunda*, 173–200  
*Rhytipterna simplex*, 173–200  
 Rio Grande do Sul, 133–137  
*Riparia riparia*, 173–200, 381–398, 447–482  
 Risks, S199–S204  
 Robaldo Guedes, N. M., S279–S283  
 Robbins, M. B., 173–200  
 Robichaud, L., S267–S278  
 Rocha, C. D., S169–S174  
 Rod, S223–S232, S251–S260  
 Rodríguez-Ferraro, A., 253–256, 269–270  
 Rohwer, V. G., 381–398  
 Rojas, L. M., S223–S232, S233–S242, S251–S260  
 Rojas-Nossa, S. V., S205–S213  
*Rollandia rolland*, 381–398  
 Roosts, 313–317  
 Rosero Lasprilla, L., S183–S190  
*Rostrhamus hamatus*, S83–S99  
*Rostrhamus sociabilis*, 71–85, 381–398, S83–S99  
 Rozzi, R., S395–S407, S419–S434, S435–S444  
*Rupornis magnirostris*, 71–85  
*Rynchops niger*, 71–85, 447–482  
*Sakesphorus canadensis*, 173–200  
 Salamanca, S267–S278  
*Saltator atriceps*, 447–482  
*Saltator aurantiirostris*, 381–398  
*Saltator coerulescens*, 173–200, 247–251, 447–482  
*Saltator grossus*, 447–482  
*Saltator maximus*, 447–482  
*Saltator similis*, S117–S126  
 San José Island, 237–246  
 Sánchez, G., 381–398  
 Sandy coastal plain, 279–282  
 Sankoff characters, S285–S291  
*Sarcoramphus papa*, 173–200, 219–224, 381–398, 447–482, 501–512, S83–S99  
*Sarkidiornis melanotos*, 381–398  
 Satellite image, S367–S376  
*Satenes modesta*, 513–525  
*Satrapa icterophrys*, S349–S355  
 Sauropoda, S101–S116  
 Savanna, 173–200  
*Sayornis nigricans*, 447–482  
 Sazima, M., S183–S190  
*Scardafella squammata*, 71–85  
 Scheibler, D. R., 65–70  
 Schelsky, W. M., S317–S321  
*Schiffornis turdinus*, 173–200, 447–482  
*Schistoclamys melanopsis*, 173–200  
 Schmitt, C. G., 381–398  
 Schmitt, D. C., 381–398  
 Scientific method, S41–S60  
 Scientific program, S1–S34  
*Sclerurus guatemalensis*, 447–482  
*Sclerurus rufigularis*, 173–200  
 Scolopacidae, S357–S365  
*Scytalopus latrans*, 483–492  
 Seabirds, 163–171, S61–S72  
 Sealy, S. G., 331–347  
 Seed dispersal, 53–64, S117–S126  
*Seiurus aurocapillus*, 447–482  
*Seiurus motacilla*, 447–482  
*Seiurus noveboracensis*, 447–482  
*Selenidera gouldii*, 527–537  
*Selenidera maculirostris*, 527–537  
*Selenidera spectabilis*, 433–445, 447–482, 527–537  
 Semi-collared Hawk, 31–40  
 Semidecidual forest, 5–15  
*Sephanoides sephanoides*, S215–S222  
*Serpophaga griseiceps*, 145–153  
*Serpophaga munda*, 381–398  
*Serpophaga subcristata*, 71–85  
*Setophaga ruticilla*, 447–482  
 Sexual dichromatism, 543–545  
 Shorebirds, S223–S232, S357–S365, S377–S384  
*Sialia sialis*, 447–482  
*Sicalis flaveola*, 71–85, 173–200, 381–398  
*Sicalis lebruni*, 513–525  
*Sicalis luteola*, 173–200, 371–380, 447–482  
*Sirystes sibilator*, 173–200  
*Sittasomus griseicapillus*, 381–398, 447–482  
 Soave, G. E., 299–306

- Solitary Eagle, 31–40  
 Song, 365–370  
 Sotar condi, S409–S417  
 South America, S159–S167, S349–S355, S367–S376  
 South Atlantic Ocean, 163–171  
 Southeastern Pacific, S61–S72  
 Southern Brazil, 71–85  
 Southern Rupununi, 173–200  
 Souza, S., 423–427  
 Specht, G. V. A., 219–224  
 Speciation, S41–S60  
 Species richness, S127–S135  
 Spectral reflectance, S215–S222  
*Speotyto cunicularia*, 71–85, 87–92, 501–512  
*Spheniscus humboldti*, S61–S72  
*Spheniscus mendiculus*, S61–S72  
*Sphyrapicus varius*, 447–482  
*Spizca americana*, 447–482  
*Spizcaetus ornatus*, 447–482, S83–S99  
*Spizcaetus tyrannus*, 173–200, 447–482, S83–S99  
*Spizcastur melanoleucus*, 173–200, 447–482, S83–S99  
*Spizella passerina*, 447–482  
*Spiziapteryx circumcinctus*, S83–S99  
*Sporophila americana*, 447–482  
*Sporophila bouvreuil*, 173–200  
*Sporophila caeruleascens*, 71–85, 381–398  
*Sporophila insulata*, 17–30  
*Sporophila intermedia*, 173–200  
*Sporophila leucoptera*, 173–200  
*Sporophila lineola*, 381–398  
*Sporophila minuta*, 17–30, 173–200  
*Sporophila nigricollis*, 173–200  
*Sporophila plumbea*, 173–200  
*Sporophila schistacea*, 447–482  
*Sporophila telasco*, 17–30  
*Sporophila torqueola*, 447–482  
 Stable isotope, S377–S384  
 Status, S175–S181  
*Steatornis caripensis*, S233–S242, S251–S260  
*Stelgidopteryx ruficollis*, 71–85, 173–200, 381–398, 447–482  
*Stelgidopteryx serripennis*, 447–482  
*Sterna caspia*, 447–482  
*Sterna fuscata*, S61–S72  
*Sterna fuscatum*, 447–482  
*Sterna hirundo*, 121–131  
*Sterna lunata*, S61–S72  
*Sterna maxima*, 447–482  
*Sterna nilotica*, 447–482  
*Sterna paradisaea*, 163–171  
*Sterna sandvicensis*, 447–482  
*Sterna superciliaris*, 71–85  
*Sterna vittata*, 163–171  
*Stigmatura bodytoides*, 381–398, 381–398  
 Stiles, F. G., 17–30, S191–S198, S205–S213, S349–S355  
 Stratford, J. A., 265–267  
 Stratification, 41–52  
 Strem, R., 381–398  
*Streptoprogne zonaris*, 173–200, 447–482, 483–492  
 Strigiformes, 87–92  
*Strix albitarsis*, 483–492  
*Strix nigrolineata*, S477–S483  
*Strix virgata*, S477–S483  
 Student participation, S1–S34  
*Sturnella loyca*, 513–525  
*Sturnella magna*, 173–200, 447–482  
*Sturnella militaris*, 173–200  
*Sturnella superciliaris*, 299–306  
 Stutchbury, B. J. M., 417–421  
 Stygian Owl, 423–427  
 Subandean region, 399–402  
 Subtropical montane forest  
*Suiriri suiriri*, 173–200, 381–398  
*Sula dactylatra*, S61–S72  
*Sula leucogaster*, S61–S72  
*Sula nebovicii*, S61–S72  
*Sula sula*, S61–S72  
*Sula variegata*, S61–S72  
 Sunlight, 145–153  
 Supertramp, S455–S465  
 Survey techniques, 513–525  
 Systematics, S73–S81  
*Symphonia globulifera*, S183–S190  
*Synallaxis albescens*, 145–153, 173–200, 381–398  
*Synallaxis azarae*, 483–492

- Synallaxis brachyura*, 447–482  
*Synallaxis frontalis*, 71–85, 381–398  
*Synallaxis scutata*, 381–398  
*Syrigma sibilatrix*, 381–398  
 Systematics, S293–S299  
*Tachornis squamata*, 173–200  
*Tachybaptus dominicus*, 381–398, 447–482  
*Tachycineta albilinea*, 447–482  
*Tachycineta albiventer*, 71–85, 173–200  
*Tachycineta bicolor*, 447–482  
*Tachycineta* sp., 381–398  
*Tachycineta thalassina*, 447–482  
*Tachyphonus coronatus*, S117–S126  
*Tachyphonus cristatus*, 41–52, 173–200  
*Tachyphonus delatrii*, 447–482  
*Tachyphonus luctuosus*, 447–482  
*Tachyphonus phoenicius*, 173–200  
*Tachyphonus rufus*, 173–200, 247–251  
*Tachyphonus surinamus*, 173–200  
*Taeniopygia guttata*, 201–208  
*Tangara cayana*, 71–85, 173–200, S117–S126  
*Tangara chilensis*, 173–200, 539–542  
*Tangara larvata*, 447–482  
*Tangara lavinia*, 447–482  
*Tangara nigrocincta*, 539–542  
*Tangara punctata*, 173–200  
*Tangara vassorii*, 483–492  
*Tangara velia*, 41–52, 173–200  
*Tapera naevia*, 71–85, 173–200, 447–482  
 Tapetum, S251–S260  
*Taraba major*, 173–200, 381–398, 447–482  
 Taxonomy, 17–30  
 Techniques, S329–S337, S455–S465  
 Tejada-Renjifo, J., S323–S328  
 Temperate forests, S395–S407, S419–S434, S435–S444  
 Temperate grasslands, S159–S167  
*Terentotriccus erythrurus*, 447–482  
*Terenura spodiopitula*, 173–200  
 Termitarium nesting, 319–330  
 Termitarium, 319–330  
 Termite, 319–330  
*Tersina viridis*, S117–S126  
 Teta, P., 87–92  
 Teul, M., 155–162, 155–162  
*Thalassarche chlororhynchos*, 163–171  
*Thalassarche exulans*, 163–171  
*Thalassarche melanophris*, 163–171  
*Thalassoica antarctica*, 163–171  
*Thalurania colombica*, 447–482  
*Thalurania furcata*, 173–200, S183–S190  
*Thamnistes anabatinus*, 447–482  
*Thamnomanes ardesiacus*, 173–200  
*Thamnomanes caesius*, 173–200  
*Thamnophilus amazonicus*, 173–200  
*Thamnophilus atrinucha*, 447–482  
*Thamnophilus caerulescens*, 381–398  
*Thamnophilus doliatus*, 71–85, 173–200, 381–398, 447–482  
*Thamnophilus murinus*, 173–200, 265–267  
*Thamnophilus punctatus*, 173–200  
*Thamnophilus ruficapillus*, 71–85  
*Thamnophilus sticturus*, 381–398  
 Theory, S41–S60  
*Theristicus candatus*, 173–200, 381–398, 513–525  
 Theropoda, S101–S116  
*Thinocorus rumicivorus*, 513–525  
*Thlypopsis sordida*, 71–85, 381–398  
 Thomas, R. J., S243–S250  
 Thompson, J. J., 289–297, S301–S307  
 Thraupidae, 53–64  
*Thraupis abbas*, 447–482  
*Thraupis bonariensis*, 381–398  
*Thraupis episcopus*, 173–200, 447–482  
*Thraupis palmarum*, 41–52, 71–85, 173–200  
*Thraupis sayaca*, 71–85, 381–398, S117–S126  
 Threatened birds, S175–S181  
*Tbrenetes ruckeri*, 447–482  
*Thripophaga fusciceps*, 257–263  
*Thripophaga macroura*, 257–263  
*Thryothorus atrogularis*, 447–482  
*Thryothorus coraya*, 173–200  
*Thryothorus euophrys*, 483–492  
*Thryothorus leucotis*, 173–200  
*Thryothorus maculipectus*, 447–482  
*Thryothorus modestus*, 447–482  
*Thryothorus nigricapillus*, 447–482  
*Thryothorus thoracicus*, 447–482  
*Tiaris fuliginosa*, 247–251

- Tiaris obscurus*, 381–398  
 Tick, 247–251  
*Tigrisoma lineatum*, 71–85, 447–482  
*Tigrisoma mexicanum*, 447–482  
 Tikal National Park, S477–S483  
 Tilarán, 433–445  
 Timing of first breeding, S385–S394  
 Tinamidae, 111–119, S285–S291, S293–S299,  
     S301–S307, S309–S315, S317–S321,  
     S323–S328, S329–S337, S339–S347  
*Tinamotis ingoufi*, 513–525  
*Tinamus guttatus*, S317–S321, S323–S328  
*Tinamus major*, 173–200, 447–482, S317–S321,  
     S323–S328, S329–S337  
*Tinamus tao*, S317–S321, S323–S328  
*Tityra cayana*, 173–200  
*Tityra inquisitor*, 173–200, 447–482  
*Tityra semifasciata*, 447–482  
*Todirostrum cinereum*, 71–85, 173–200, 447–482  
*Todirostrum pictum*, 173–200  
*Tolmomyias assimilis*, 173–200  
*Tolmomyias flaviventris*, 173–200  
*Tolmomyias poliocephalus*, 173–200  
*Tolmomyias sulphurens*, 173–200, 381–398,  
     447–482  
 Tomazzoni, A. C., 279–282  
*Tornatina candei*, 357–364  
 Torres, J., S377–S384  
 Toucan, 433–445  
 Traditional ecological knowledge, S395–S407  
 Trait, S455–S465  
 Transects, 513–525  
 Travaini, A., 513–525  
*Travisia olens*, 357–364  
 Trejo, A., 307–311  
*Trichothraupis melanops*, 381–398  
*Tringa flavipes*, 71–85, 173–200, 381–398, 447–  
     482  
*Tringa melanoleuca*, 173–200, 381–398, 447–  
     482, S223–S232  
*Tringa solitaria*, 173–200, 381–398, 447–482  
 Trochilinae, S183–S190  
*Troglodytes aedon*, 71–85, 271–278, 371–380,  
     381–398, 447–482, 513–525  
*Troglodytes musculus*, 173–200  
*Trogon clathratus*, 433–445  
*Trogon collaris*, 447–482  
*Trogon curucui*, 319–330, 381–398  
*Trogon massena*, 447–482  
*Trogon melanocephalus*, 447–482  
*Trogon melanurus*, 173–200, 319–330  
*Trogon personatus*, 483–492  
*Trogon rufus*, 173–200, 447–482  
*Trogon violaceus*, 173–200, 447–482  
*Trogon viridis*, 173–200  
 Trogon, 433–445  
 Trogonidae, 433–445  
 Troodontidae, S101–S116  
 Tropical montane forest, 31–40  
 Trujillo, A., 253–256  
*Tryngites subruficollis*, 381–398, 447–482, S159–  
     S167, S367–S376  
 Tucumán, 501–512  
*Turdus albicollis*, 173–200, 247–251  
*Turdus amaurochalinus*, 381–398, S117–S126  
*Turdus assimilis*, 201–208  
*Turdus fumigatus*, 247–251  
*Turdus fusater*, 483–492  
*Turdus grayi*, 447–482  
*Turdus leucomelas*, 71–85, 173–200, S117–S126  
*Turdus nudigenis*, 173–200, 247–251  
*Turdus rufiventris*, 71–85, 381–398, S117–S126  
*Tyrannetes stolzmanni*, 41–52  
*Tyrannetes virescens*, 173–200  
 Tyrannidae, 145–153  
*Tyrannopsis sulphurea*, 41–52, 173–200  
*Tyrannulus elatus*, 41–52, 173–200  
*Tyrannus albogularis*, 41–52, 173–200  
*Tyrannus melancholicus*, 41–52, 71–85, 173–200,  
     271–278, 381–398, 447–482, S117–S126  
*Tyrannus savana*, 41–52, 71–85, 173–200, 381–  
     398, 447–482, S117–S126, S349–S355  
*Tyrannus tyrannus*, 447–482  
*Tyrannus vociferans*, 447–482  
*Tyto alba*, 65–70, 307–311, 413–415, 447–482,  
     501–512  
 Underwood, T. J., 331–347  
*Upucerthia dumetaria*, 513–525  
 Urban waste site, 93–102  
*Uropelia campestris*, 173–200



- Uruguay, S367–S376  
 Use-availability, 111–119  
 Valdez, U., 31–40  
 Valencia Rodríguez, I. D., S445–S454  
*Vanellus chilensis*, 71–85, 173–200, 371–380, 381–398, 513–525, S223–S232  
*Vanellus resplendens*, 483–492  
 Vargas, P., S339–S347  
 Vargas, S., S339–S347  
 Vasconcelos, M. F. de, 547–548  
 Vegetation type, 111–119  
 Venezuela, 139–141, 225–235, 253–256, 269–270, 407–411  
*Veniliornis cassini*, 173–200  
*Veniliornis fumigatus*, 447–482  
*Veniliornis nigriceps*, 483–492  
*Veniliornis passerinus*, 71–85, 173–200  
*Veniliornis spilogaster*, S117–S126  
 Vereá, C., 225–235, 247–251  
*Vermivora chrysoptera*, 447–482  
*Vermivora peregrina*, 447–482  
*Vermivora pinus*, 447–482  
 Video camera, S455–S465  
 Vigilance, 407–411  
 VII<sup>th</sup> Neotropical Ornithological Congreso, S1–S34  
 VIII<sup>th</sup> Neotropical Ornithological Congress, 575  
 Vilella, F. J., S455–S465  
*Vireo flavifrons*, 447–482  
*Vireo flavoviridis*, 447–482  
*Vireo gilvus*, 447–482  
*Vireo griseus*, 447–482  
*Vireo olivaceus chivi*, 381–398  
*Vireo olivaceus*, 41–52, 173–200, 447–482, S117–S126  
*Vireo pallens*, 447–482  
*Vireo philadelphicus*, 447–482  
*Vireolanus leucotis*, 173–200  
*Vireolanus pulchellus*, 447–482  
 Vision, S215–S222, S223–S232, S233–S242  
 Visual constraints, S243–S250  
 Visual fields, S233–S242  
*Volatinia jacarina*, 71–85, 173–200, 381–398, 447–482, S117–S126  
 Vuilleumier Fund, S1–S34  
 Vuilleumier, F., S1–S34, S41–S60  
*Vultur gryphus*, 313–317, 501–512, S83–S99  
 Wajntal, A., 527–537  
 Walther, B. A., 41–52  
 Waste site, 93–102  
 Waterbirds, S61–S72, S445–S454  
 Weather, S261–S266  
 West Indies, 289–297  
 Wetlands, S445–S454  
 White, T. H., S467–S476  
 Wiedenfeld, D. A., 447–482  
 Wilson, R. E., 543–545  
*Wilsonia canadensis*, 139–141, 447–482  
*Wilsonia citrina*, 447–482  
*Wilsonia pusilla*, 447–482  
 Winker, K., 53–64  
*Xantopsar flavus*, 133–137  
*Xenopipo atronitens*, 173–200  
*Xenops minutus*, 173–200, 447–482  
*Xenops rutilans*, 381–398  
*Xenopsaris albinucha*, 173–200  
*Xiphocolaptes major*, 381–398  
*Xiphocolaptes promeropirhynchus*, 483–492  
*Xipholena punicea*, 173–200  
*Xipborhynchus guttatus*, 173–200  
*Xipborhynchus pardalotus*, 173–200  
*Xipborhynchus picus*, 173–200  
*Xipborhynchus susurrans*, 447–482  
*Xipborhynchus lachrymosus*, 447–482  
*Xolmis cinerea*, 173–200  
*Xolmis irupero*, 381–398  
*Xolmis velata*, S117–S126  
 Yahgan, S395–S407, S435–S444  
 Yorio, P., 93–102  
 Zanette, L., S137–S144  
 Zárate, R., 139–141  
*Zenaida auriculata*, 71–85, 173–200, 371–380, 381–398, 483–492, 513–525, S117–S126  
*Zenaida macroura*, 447–482  
*Zimmerius gracilipes*, 173–200  
*Zimmerius vilissimus*, 447–482  
*Zonotrichia capensis*, 71–85, 271–278, 371–380, 381–398, 483–492, 513–525, S117–S126  
 Zorzin, G., 219–224

## REVIEWERS FOR ORNITOLOGÍA NEOTROPICAL, VOLUME 15

Each manuscript submitted for publication in *Ornitología Neotropical* is generally reviewed by two specialists, often including a member of the editorial board, and frequently by a third one. The following individuals kindly reviewed one or more (\*) manuscripts for Volume 15: Rosana M. Aramburu, Adrián B. Azpiroz\*, Isabel Bellocq\*, Andrew Benneth, Richard Bierregaard, Keith L. Bildstein\*, J. S. Birdsley, Walter J. Bock, Roberto Cavalcanti, David A. Christie, José Maria Cardoso da Silva, Andrew Cockburn, J. A. Donazar, Jessica R. Eberhard, Michael R. Erwin, Marco Favero, Mercedes Foster, Rosendo Fraga\*, Esteban Frere, Patricia Gandini, Michael Gochfeld, Patricia González, Jürgen Haffer, Floyd E. Hayes, Juan Pablo Isaach, Fabián Jaksic\*, Leo Joseph, E. L. Kershner, Guy M. Kirwan, Oliver Komar, Nils Krabbe, Daniel F. Lane, Scott M. Lanyon, Catherine A. Lindell, Bette Loiselle, Manuel Marín\*, Miguel Ângelo Marini, J. S. Marks, Paul Martin, Ray-

mond McNeil\*, T. A. Melo-Junior, Michael J. Meyers, Eugene S. Morton\*, Luciano Nicolas Naka\*, Ulysse Pardiñas\*, Tomas Pärt, Y. Oniki, Mauro Pichorim, Augusto Piratelli, Edwin R. Price, Flavio Quintana, John H. Rappole, J. van Remsen, Robin Restall, Nathan H. Rice, R. S. Ridgely, Mark Robbins, K. V. Douglas Robinson\*, James J. Roper, Rosenberg, Roberto P. Schlatter\*, Karl-L. Schuchmann, Grace Servat, Marina Silva, Ralf Strewe, Marc Théry, Jean-Marc Thiollay, A. Travaini, Ana Trejo, A. C. Vallely, François Vuilleumier, K. Weidinger, David F. Whitacre\*, Clayton M. White, Andrew Whittaker, James W. Wiley\*, Edwin O. Willis\*, Joseph W. Wunderle Jr., Pablo M. Yorio\*, Guthric Zimmerman, Kristof Zyskowski\*. The abstracts were kindly corrected by Cristina Yumi Miyaki for the Portuguese, and by Elisa Bonaccorso and Diego Montalti for the Spanish. Page-setting was done by Elisa Muñoz and María Rojas G.