

Up-the-country boys, 89.	Winter creeper, 145.
Vison-la, 49.	Winter gannet, 27.
War bird, 95.	Winter gros-bec, 54.
Warloon, 4.	Winter gull, 14, 15.
Water-belcher, 49.	Winter ox-eye, 68.
Water buzzard, 29.	Winter robin, 154.
Water guinea or water guinea- hen, 59.	Wood-knocker, 98.
Wear hen, 2.	Woodpecker, Great black, 100.
Weaser, 30a.	Wood-pile quawker, 80.
Whale-bird, 61.	Wood wren, 144, 149.
Wheatbird, 125.	Woozer, 30a.
Wheeler, Yellow, 102.	Wop, 49, 54.
Whicker, Yellow, 102.	Wren, Wood, 144, 149.
Whip-tail, 12b.	Yellow-bird, 149.
White crane, 51, 55.	Yellow hammer, 138.
White gull, 15.	Yellow-legged goose, 45.
White hawk, 91.	Yellow-legs, 45.
White-tailed stib, 66.	Yellow-legs, Bastard, 65.
Whittaker, 102.	Yellow-shanked sandpiper, 71.
Wick-up, 102.	Yellow snipe, 62.
Winter birds, 68.	Yellow wheeler, 102.
Winter chip-bird, 118.	Yellow whicker, 102.
	Zin-zin, 31.

MORE RECORDS FROM THE "SHORES"
COLLECTION.

BY W. F. HENNINGER.

1. *Gallinago delicata*.

Sage and Bishop give this species as "very rare in summer or winter," while it is common in spring and fall. To their records of this species for the summer months must now be added another one, a ♂, taken July 9, 1875, at Suffield, Connecticut, by Dr. Shores (now No. 1356, coll. W. F. H.),

2. *Helodromas solitarius*.

To the three unusual records for this species in Connecticut, given by Sage and Bishop, Birds of Connecticut, page 61, a fourth one must now be added, a ♂, taken July 12, 1875, at Suffield, Connecticut, by Dr. Shores (now No. 1355, coll. W. F. H.). Dr. Shores' earliest record for this species is May 17, 1874, a ♂ taken, and his latest record is August 20, 1875, a ♂ taken.

3. *Oxyechus vociferus*.

Quite a number of records must be added to those given by Sage and Bishop in Birds of Connecticut, pages 65 and 66. Besides the one mentioned in March, 1917, issue of Wilson Bulletin, there are

the following: May 26, 1874, a ♂ taken; July 12, 1875, one seen; April 12, 1875, a flock seen, which was very wild, so that in spite of a long and tedious chase he could not procure any; September 1, 1877, a ♀ shot. All at Suffield, Conn.

4. *Accipiter velox*.

An additional winter record is December 1, 1874, a ♀, taken at Suffield, Conn.

5. *Empidonax flaviventris*.

A ♀ taken June 24, 1874, at Suffield, Conn. This record was inadvertently omitted in the report as given in the March, 1917, Wilson Bulletin, and it is earlier than any that Sage and Bishop give (Birds of Conn., p. 103).

6. *Euphagus carolinus*.

On May 11, 1874, Dr. Shores took one ♂ and two ♀♀ at Suffield, Conn. This is a late spring record for Connecticut, where this species is not common in spring, though common in the fall.

7. *Zonotrichia leucophrys*.

An additional record is May 23, 1875, a ♀ taken at Suffield.

8. *Melospiza georgiana*.

The very few winter records given by Sage and Bishop for this species show that it is a rare winter resident throughout the region. Dr. Shores took a specimen, a ♀, at Cranston, R. I., on December 3, 1874.

9. *Dendroica vigorsii*.

I have been requested to publish the Shores records for this species in full, so here they are: June 5, 1874, a ♀ taken; May 28, 1875, a ♂ taken; May 3, 1877, two ♀♀ taken (now Nos. 1530 and 1599, coll. W. F. H.); June 26, 1879, a pair taken. These are all from Suffield, Conn. June 22, 1880, two ♂♂, one ♀ taken at the Southwick Ponds, Mass.

10. *Wilsonia canadensis*.

I have also been asked to give the Shores records for this species: June 5, 1874, a ♀ taken; June 6, 1874, a ♀ taken; May 23, 1875, a ♂ taken (now No. 1491, coll. W. F. H.); May 27, 1876, a ♂ taken; May 22, 1878, a ♂ taken; May 29, 1878, a ♂ taken; August 30, 1879, a ♀ taken (now No. 1492, coll. W. F. H.); September 1, 1879, a ♀ taken; September 2, 1879, a ♀ juvenal taken (now No. 1493, coll. W. F. H.); September 6, 1879, two ♂♂ taken. All at Suffield, Conn.